

Journal of Mathematical Extension
Vol. 18, No. 8, (2024) (3)1-17
URL: <https://doi.org/10.30495/JME.2024.2919>
ISSN: 1735-8299
Original Research Paper

Screen Locally Conformal Lightlike Hypersurfaces of an Almost Para-Hyperhermitian Statistical Manifold

M.B. Kazemi Balgeshir*

University of Zanjan

S. Miri

Azarbaijan Shahid Madani University

M. Ilmakchi

Azarbaijan Shahid Madani University

Abstract. In this paper, we investigate lightlike hypersurfaces of an almost para-hyperhermitian statistical manifold of constant holomorphic sectional curvature. We obtain some relations between the induced objects of such lightlike hypersurfaces with a conformal shape operator on the screen distribution. Further, we give an example of a lightlike hypersurface of an almost para-hyperhermitian statistical manifold.

AMS Subject Classification: 53A15; 53C05; 60D05

Keywords and Phrases: Screen locally conformal, lightlike hypersurface, mixed 3-structure, statistical manifold

Received: November 2023; Accepted: December 2024

*Corresponding Author

1 Introduction

Information geometry is a branch of science to extract features from objects which has fascinating applications in machine learning and evolutionary biology. A statistical manifold is a differential manifold that specifies each point with a probability distribution. In 1987, Lauritzen defined the notion of statistical manifold as a generalization of a statistical model with the Fisher metric and the Amari-Chenstov tensor [2]. Statistical manifolds are geometric objects viewed as a Riemannian manifold which admits a torsion-free affine connection $\bar{\nabla}$ and its dual connection $\bar{\nabla}^*$ with respect to the metric \bar{g} .

In lightlike submanifolds, the normal vector bundle and the tangent bundle intersects each other which does not occur in non-degenerate submanifolds [1, 8, 11].

The theory of lightlike hypersurfaces has been subject of interest by many of authors [5, 6, 12]. C. Atindogbe and K.L. Duggal investigated screen locally conformal lightlike hypersurfaces and derived some classification theorems [3]. Lightlike hypersurfaces of a statistical manifold were discussed by O. Bahadir, M.M. Tripathi [4].

However, the conception of Sasakian structures was presented by Shigeo Sasaki and further H. Furuhashi developed this idea for statistical manifolds [9]. K.L. Duggal and B. Sahin surveyed real lightlike hypersurfaces of an indefinite quaternion Kaehler manifolds [7, 14]. The main properties of a para-quaternionic hermitian manifold were given in [10, 13]. We intend to use these conceptions to achieve equivalent results for lightlike hypersurfaces of a para-hyperhermitian statistical manifold whose holomorphic sectional curvature is constant.

The present work is organized as follows: Section 2, contains some basic definitions about statistical manifolds and manifolds with mixed 3-structures. In Section 3 we study lightlike hypersurfaces of an almost para-hyperhermitian manifold. The relations between the induced objects of such lightlike hypersurfaces are obtained in Section 3. In particular, we review screen locally conformal lightlike hypersurfaces of an almost para-hyperhermitian statistical manifold of constant holomorphic sectional curvature. Also, the last section is concluded with an example.

2 Preliminaries

2.1 Statistical manifolds

Suppose that $(\bar{\mathcal{M}}, \bar{\mathfrak{g}})$ is a semi-Riemannian manifold and $\bar{\nabla}$ is an affine connection on $\bar{\mathcal{M}}$ associated with the semi-Riemannian metric $\bar{\mathfrak{g}}$. We will review some main definitions about statistical manifolds based on [9].

Definition 2.1. The triple $(\bar{\mathcal{M}}, \bar{\nabla}, \bar{\mathfrak{g}})$ is termed as a statistical manifold if $\bar{\nabla}$ is torsion free and the equalization

$$(\bar{\nabla}_E \bar{\mathfrak{g}})(F, C) = (\bar{\nabla}_F \bar{\mathfrak{g}})(E, C) \quad (1)$$

is satisfied for all $E, F, C \in \Gamma(T\bar{\mathcal{M}})$.

The dual affine connection $\bar{\nabla}^*$ of $\bar{\nabla}$ is indicated by

$$E\bar{\mathfrak{g}}(F, C) = \bar{\mathfrak{g}}(\bar{\nabla}_E F, C) + \bar{\mathfrak{g}}(F, \bar{\nabla}_E^* C), \quad (2)$$

Denote by $\bar{\nabla} = \frac{1}{2}(\bar{\nabla} + \bar{\nabla}^*)$ the Levi-Civita connection associated with the metric $\bar{\mathfrak{g}}$.

Remark 2.2. For a statistical manifold $(\bar{\mathcal{M}}, \bar{\nabla}, \bar{\mathfrak{g}})$, we define a tensor $\mathcal{K} \in \Gamma(T\bar{\mathcal{M}}^{(1,2)})$ by $\mathcal{K}_E F = \frac{1}{2}(\bar{\nabla}_E F - \bar{\nabla}_E^* F)$ which satisfies

$$\mathcal{K}_E F = \mathcal{K}_F E, \quad \bar{\mathfrak{g}}(\mathcal{K}_E F, C) = \bar{\mathfrak{g}}(F, \mathcal{K}_E C). \quad (3)$$

for all $E, F, C \in \Gamma(T\bar{\mathcal{M}})$.

Let $\bar{\mathcal{R}}, \bar{\mathcal{R}}^*$ be the curvature tensor fields of $\bar{\nabla}$ and $\bar{\nabla}^*$, respectively. Then the statistical curvature tensor field of the manifold $(\bar{\mathcal{M}}, \bar{\nabla}, \bar{\mathfrak{g}})$ is characterized by $\bar{\mathcal{S}}(E, F)C = \frac{1}{2}\{\bar{\mathcal{R}}(E, F)C + \bar{\mathcal{R}}^*(E, F)C\}$ for all $E, F, C \in \Gamma(T\bar{\mathcal{M}})$.

A $(1, 1)$ -tensor field \mathfrak{X} which satisfies $\mathfrak{X}^2 = -\mathcal{I}d$ is called an almost complex structure on $\bar{\mathcal{M}}$. Let $\mathfrak{X} \in \Gamma(T\bar{\mathcal{M}}^{(1,1)})$ be an almost complex structure such that $\bar{\mathfrak{g}}(\mathfrak{X}E, \mathfrak{X}F) = \bar{\mathfrak{g}}(E, F)$. We put θ as a 2-form on $\bar{\mathcal{M}}$ defined by $\theta(E, F) = \bar{\mathfrak{g}}(E, \mathfrak{X}F)$. A statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla})$

furnished by an almost complex structure \mathfrak{X} satisfying $\bar{\nabla}\theta = 0$, is called a holomorphic statistical manifold. In addition, the following relations $\bar{\nabla}_E\mathfrak{X}F = \mathfrak{X}\bar{\nabla}_E^*F$ and $\bar{\mathcal{R}}(E, F)\mathfrak{X}C = \mathfrak{X}\bar{\mathcal{R}}^*(E, F)C$ are deducible for all $E, F, C \in \Gamma(TM)$.

Definition 2.3. A holomorphic statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \mathfrak{X})$ is supposed to be of constant holomorphic sectional curvature $\bar{c} \in \mathbb{R}$ if

$$\begin{aligned} \bar{S}(E, F)C &= \frac{\bar{c}}{4}\{\bar{\mathfrak{g}}(F, C)E - \bar{\mathfrak{g}}(E, C)F \\ &+ \bar{\mathfrak{g}}(\mathfrak{X}F, C)\mathfrak{X}E - \bar{\mathfrak{g}}(\mathfrak{X}E, C)\mathfrak{X}E + 2\bar{\mathfrak{g}}(E, \mathfrak{X}F)\mathfrak{X}C\} \end{aligned} \quad (4)$$

holds for all $E, F, C \in \Gamma(T\bar{\mathcal{M}})$.

2.2 Mixed 3-structure manifolds

An almost product structure \mathfrak{X} on a smooth semi-Riemannian manifold $\bar{\mathcal{M}}$ is a $(1, 1)$ -tensor field satisfying $\mathfrak{X}^2 = \mathcal{I}d$, where $\mathcal{I}d$ indicates the identity tensor field on $\bar{\mathcal{M}}$.

Definition 2.4. [10] Let $H = (\mathfrak{X}_i)_{i=1,2,3}$ be a local basis of subbundle of $End(T\bar{\mathcal{N}})$ of rank 3. Then, $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \mathfrak{X} = (\mathfrak{X}_i)_{i=1,2,3})$ is called an almost para-hypercomplex manifold if $\mathfrak{X}_1, \mathfrak{X}_2$ are almost product structure on $\bar{\mathcal{M}}$ and \mathfrak{X}_3 is an almost complex structure on $\bar{\mathcal{M}}$ which satisfies $\mathfrak{X}_1\mathfrak{X}_2 = -\mathfrak{X}_2\mathfrak{X}_1 = \mathfrak{X}_3$.

A semi-Riemannian metric $\bar{\mathfrak{g}}$ on $\bar{\mathcal{M}}$ satisfying

$$\bar{\mathfrak{g}}(\mathfrak{X}_1E, \mathfrak{X}_1F) = \bar{\mathfrak{g}}(\mathfrak{X}_2E, \mathfrak{X}_2F) = -\bar{\mathfrak{g}}(\mathfrak{X}_3E, \mathfrak{X}_3F) = -\bar{\mathfrak{g}}(E, F), \quad (5)$$

is called compatible to the almost para-hypercomplex structure $H = (\mathfrak{X}_i)_{i=1,2,3}$ for all $E, F \in \Gamma(T\bar{\mathcal{M}})$.

Definition 2.5. [10] A triple $(\psi_i, \zeta_i, \eta_i)_{i=1,2,3}$ of structures on $(\bar{\mathcal{M}}, \bar{\nabla}, \bar{\mathfrak{g}})$ satisfying

$$\psi_i^2 = \tau_i(-\mathcal{I} + \eta_i \otimes \mathfrak{z}_i), \quad \eta_i(\zeta_i) = 1 \quad \tau_1 = \tau_2 = -\tau_3 = -1 \quad (6)$$

is said to be a mixed 3-structure if $(\psi_1, \zeta_1, \eta_1)$ and $(\psi_2, \zeta_2, \eta_2)$ are almost paracontact structures with $\tau_i = 1$, and $(\psi_3, \zeta_3, \eta_3)$ is an almost contact

structure, that is $\tau_i = -1$. Here, ζ'_i s indicate the structure vector fields, η_i s are 1-forms on $\bar{\mathcal{M}}$ and ψ'_i s are (1,1)-tensor fields. Moreover, (6) yields

$$\begin{aligned} \eta_i(\zeta_j) &= 0, & \psi_i(\eta_j) &= \tau_j \zeta_k, \\ \psi_j(\zeta_i) &= -\tau_j \zeta_k \\ \eta_i \circ \psi_j &= -\eta_j \circ \psi_i = \tau_k \eta_k, \\ \psi_i \psi_j - \tau_i \eta_j \otimes \zeta_i &= -\psi_j \psi_i + \tau_j \eta_i \otimes \zeta_j = \tau_k \psi_k. \end{aligned} \quad (7)$$

where (i, j, k) is an even permutation of $(1, 2, 3)$.

A semi-Riemannian metric $\bar{\mathfrak{g}}$ on the smooth manifold $\bar{\mathcal{M}}$ is called compatible to the mixed 3-structure $(\psi_i, \zeta_i, \eta_i)_{i=1,2,3}$, if the relation

$$\bar{\mathfrak{g}}(\psi_i E, \psi_i F) = \tau_i [\bar{\mathfrak{g}}(E, F) - \varepsilon_i \eta_i(E) \eta_i(F)] \quad (8)$$

holds for any $E, F \in \Gamma(T\bar{\mathcal{M}})$, where $\varepsilon_i = \bar{\mathfrak{g}}(\zeta_i, \zeta_i) = \pm 1$, $i = 1, 2, 3$.

2.3 Lightlike real hypersurfaces

Let $(\bar{\mathcal{M}}, \bar{\mathfrak{g}})$ be an $(n+1)$ -dimensional semi-Riemannian manifold and $(\mathcal{M}, \mathfrak{g})$ be a hypersurface of $\bar{\mathcal{M}}$. If $\bar{\mathfrak{g}}$ is degenerate then the normal vector bundle $T^\perp(\mathcal{M})$ and tangent vector bundle $T\mathcal{M}$ have an intersection along a non-zero differentiable distribution $\text{rad}(T\mathcal{M})$ indicated by radical distribution. For a lightlike hypersurface \mathcal{M} of a semi-Riemannian manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}})$, we have $\text{rad}(T\mathcal{M}) = T^\perp(\mathcal{M})$. Denoted by $\mathfrak{s}(T\mathcal{M})$ the complementary subbundle of $T\mathcal{M}$ to $\text{rad}(T\mathcal{M})$, we have $T\mathcal{M} = T^\perp(\mathcal{M}) \oplus \mathfrak{s}(T\mathcal{M})$ [7]. Denote by $\text{tr}(T\mathcal{M})$ the complementary (but not orthogonal) vector bundle to $T\mathcal{M}$ in $T\bar{\mathcal{M}}$. We have the decomposition

$$T\bar{\mathcal{M}} = \mathfrak{s}(T\mathcal{M}) \perp (T^\perp(\mathcal{M}) \oplus \text{tr}(T\mathcal{M})) = T\mathcal{M} \oplus \text{tr}(T\mathcal{M}), \quad (9)$$

The Gauss-Weingarten formulas for a lightlike hypersurface \mathcal{M} of $(\bar{\mathcal{M}}, \bar{\mathfrak{g}})$, are given by

$$\bar{\nabla}_E F = \nabla_E F + \omega(E, F), \quad \bar{\nabla}_E \mathcal{L} = -A_{\mathcal{L}} E + \nabla_F^\perp \mathcal{L}. \quad (10)$$

Here, $\{\nabla_E F, A_{\mathcal{L}} E\}$ belong to $\Gamma(T\mathcal{M})$ and $\{\omega, \nabla_F^\perp \mathcal{L}\} \in \Gamma(\text{tr}(T\mathcal{M}))$. We set $\sigma(E, F) = \bar{\mathfrak{g}}(\omega(E, F), \zeta)$ and $s(E) = \bar{\mathfrak{g}}(\nabla_E^\perp \mathcal{L}, \zeta)$. From (10), we have the following formulas

$$\bar{\nabla}_E F = \nabla_E F + \sigma(E, F) \mathcal{L}, \quad \bar{\nabla}_E \mathcal{L} = -A_{\mathcal{L}} E + s(E) \mathcal{L} \quad (11)$$

for all $E, F \in \Gamma(T\mathcal{M}), \mathcal{L} \in \Gamma(tr(T\mathcal{M}))$ and $\zeta \in \Gamma(\mathfrak{rad}(T\mathcal{M}))$. Here, σ denotes the second fundamental form associated with $\bar{\nabla}$ and $A_{\mathcal{L}}$ is the shape operator on \mathcal{M} . Let denote by \mathcal{P} the projection morphism of $T\mathcal{M}$ on $\mathfrak{s}(T\mathcal{M})$. Then, the Gauss-Weingarten formulas for $\mathfrak{s}(T\mathcal{M})$ are given by

$$\nabla_E \mathcal{P}F = \nabla'_E \mathcal{P}F + \rho(E, \mathcal{P}F)\zeta, \quad \nabla_E \zeta = -A'_\zeta E + s'(E)\zeta \quad (12)$$

Here, $\{\nabla'_E \mathcal{P}F, A'_\zeta E\}$ belong to $\Gamma(\mathfrak{s}(T\mathcal{M}))$ and we have

$$\begin{aligned} \rho(E, \mathcal{P}F) &= \bar{\mathfrak{g}}(\nabla_E \mathcal{P}F, \mathcal{L}), & s'(E) &= \bar{\mathfrak{g}}(\nabla_E \zeta, \mathcal{L}), \\ s(E) &= -s'(E), \end{aligned} \quad (13)$$

for all $E, F \in \Gamma(T\mathcal{M}), \zeta \in \Gamma(\mathfrak{rad}(T\mathcal{M}))$ and $\mathcal{L} \in \Gamma(tr(T\mathcal{M}))$.

Moreover, we have

$$\bar{\mathfrak{g}}(\mathfrak{X}_i \zeta, \zeta) = \bar{\mathfrak{g}}(\mathfrak{X}_i \mathcal{L}, \mathcal{L}) = \bar{\mathfrak{g}}(\mathfrak{X}_i \zeta, \mathcal{L}) = 0, \quad \bar{\mathfrak{g}}(\mathfrak{X}_i \zeta, \mathfrak{X}_i \mathcal{L}) = \tau_i, \quad (14)$$

$\mathfrak{X}_i T^\perp(\mathcal{M})$ and $\mathfrak{X}_i tr(T\mathcal{M})$ are distributions on \mathcal{M} of rank 3 such that $\mathfrak{X}_i T^\perp(\mathcal{M}) \cap T^\perp(\mathcal{M}) = 0$ and $\mathfrak{X}_i tr(T\mathcal{M}) \cap T^\perp(\mathcal{M}) = 0, i = 1, 2, 3$. Thus, $\mathfrak{X}_i T\mathcal{M} \perp \oplus \mathfrak{X}_i tr(T\mathcal{M})$ is a vector subbundle of $\mathfrak{s}(T\mathcal{M})$, where $\{e, f, g\}$ denotes an even permutation of $\{1, 2, 3\}$. Besides, we have $\bar{\mathfrak{g}}(\mathfrak{X}_i \zeta, \mathfrak{X}_j \mathcal{L}) = 0$, which consequently implies that $\mathfrak{X}_i T^\perp(\mathcal{M}) \oplus \mathfrak{X}_i tr(T\mathcal{M})$ is a vector subbundle of $\mathfrak{s}(T\mathcal{M})$ of rank 6. Thus, there exists a non-degenerate distribution Δ_0 on \mathcal{M} such that $\mathfrak{s}(T\mathcal{M}) = \{\Delta_1 \oplus \Delta_2\} \perp \Delta_0$, where $\Delta_1 = \mathfrak{X}_1 \zeta \oplus \mathfrak{X}_2 \zeta \oplus \mathfrak{X}_3 \zeta$ and $\Delta_2 = \mathfrak{X}_1 \mathcal{L} \oplus \mathfrak{X}_2 \mathcal{L} \oplus \mathfrak{X}_3 \mathcal{L}$. Thus, the following decomposition

$$T\mathcal{M} = \{T^\perp(\mathcal{M}) \oplus_{orth} \Delta_0 \oplus (\Delta_1 \oplus \Delta_2)\}, \quad \Delta = \{T^\perp(\mathcal{M}) \oplus \Delta_1\} \oplus \Delta_0 \quad (15)$$

are obtained. Considering $\mathfrak{X}_i \zeta = \zeta_i$ and $\mathfrak{X}_i \mathcal{L} = \xi_i$, we define $\mu_i, \nu_i \in \Gamma(T\mathfrak{X}^{(0,1)})$ by

$$\mu_i(E) = \bar{\mathfrak{g}}(E, \zeta_i), \quad \nu_i(E) = \bar{\mathfrak{g}}(E, \xi_i). \quad (16)$$

Let \tilde{S} be the projection morphism of $T\mathcal{M}$ on Δ . Consequently, we may write

$$E = \tilde{S}E + \mu_i(E)\xi_i, \quad (17)$$

and

$$\mathfrak{X}_i E = \psi_i E + \mu_i(E) \mathcal{L} \quad (18)$$

for any $E \in \Gamma(T\mathcal{M})$, where $\psi_i E$, imply tangent part of $\mathfrak{X}_i E$. Applying \mathfrak{X}_i to (18) and using the fact that $\mathfrak{X}_i^2 = -\tau_i \mathcal{I}$, we have

$$\psi_i^2 E = \tau_i(-E + \mu_i(E) \xi_i). \quad (19)$$

From (16), (18) and (19), it concludes that

$$\begin{aligned} \nu_i(\xi_i) &= \nu_i(\zeta_j) = \mu_i(\xi_j) = 0, \quad \psi_i(\zeta_j) = \tau_j \zeta_k, \\ \mu_i \circ \psi_j &= -\mu_j \circ \psi_i = \tau_k \mu_k, \quad \psi_i \psi_j - \tau_i \mu_j \otimes \mu_i = -\psi_j \psi_i + \tau_j \mu_i \otimes \mu_j = \tau_k \psi_k \end{aligned}$$

where (e, f, g) is regarded as an even permutation of $(1, 2, 3)$ and $\tau_1 = \tau_2 = -\tau_3 = -1$. Then, the triple $(\psi_i, \zeta_i, \mu_i)_{i=1,2,3}$ is indicated as an almost contact mixed 3-structure on \mathcal{M} [7].

3 Lightlike Real Hypersurfaces of an Almost Para-Hyperhermitian Statistical Manifold

Consider $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$ as an almost para-hyperhermitian manifold furnished by a statistical structure $(\bar{\mathfrak{g}}, \bar{\nabla})$ on $\bar{\mathcal{M}}$. Supposing $(\mathcal{M}, \mathfrak{g})$ as a lightlike hypersurface of an almost para-hyperhermitian statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$, we have the Gauss-Weingarten formulas as follow

$$\begin{aligned} \bar{\nabla}_E F &= \nabla_E F + \sigma(E, F) \mathcal{L}, & \bar{\nabla}_E^* F &= \nabla_E^* F + \sigma^*(E, F) \mathcal{L}, \\ \bar{\nabla}_E \mathcal{L} &= -A_{\mathcal{L}}^* E + s(E) \mathcal{L}, & \bar{\nabla}_E^* \mathcal{L} &= -A_{\mathcal{L}} E + s^*(E) \mathcal{L}, \end{aligned} \quad (20)$$

respectively. Here, the induced connections on \mathcal{M} are indicated by ∇, ∇^* and σ, σ^* denote the second fundamental forms associated with $\bar{\nabla}, \bar{\nabla}^*$. Taking \mathcal{P} as the projection morphism of $T\mathcal{M}$ on $\mathfrak{s}(T\mathcal{M})$, the Gauss and Weingarten formulas for $\mathfrak{s}(T\mathcal{M})$ are given by

$$\begin{aligned} \nabla_E \mathcal{P} F &= \nabla'_E \mathcal{P} F + \rho(E, \mathcal{P} F) \zeta, & \nabla_E^* \mathcal{P} F &= \nabla'^* E \mathcal{P} F + \rho^*(E, \mathcal{P} F) \zeta, \\ \nabla_E \zeta &= -A_{\zeta}^* E + s'(E) \zeta, & \nabla_E^* \zeta &= -A_{\zeta} E + s'^*(E) \zeta, \end{aligned} \quad (21)$$

$\{A'_\zeta, A_\zeta^*\}$ are shape operators on $\mathfrak{s}(T\mathcal{M})$ and $\nabla'_E \mathcal{P}F, \nabla_E^* \mathcal{P}F, A'_\zeta E, A_\zeta^* E$ belong to $\Gamma(\mathfrak{s}(T\mathcal{M}))$. The induced geometric objects are related to each other in this way

$$\begin{aligned} \rho(E, \mathcal{P}F) &= \bar{\mathfrak{g}}(A_\mathcal{L}^* E, \mathcal{P}F), & \rho^*(E, \mathcal{P}F) &= \bar{\mathfrak{g}}(A_\mathcal{L} E, \mathcal{P}F) \\ \sigma(E, F) &= \bar{\mathfrak{g}}(A_\zeta^* E, F), & \sigma^*(E, F) &= \bar{\mathfrak{g}}(A'_\zeta E, F), \end{aligned} \quad (22)$$

for any $E, F \in \Gamma(T\mathcal{M}), \zeta \in \Gamma(\text{rad}(T\mathcal{M}))$ and $\mathcal{L} \in \Gamma(\text{tr}(T\mathcal{M}))$.

Remark 3.1. Note that the induced connection on a non-degenerate submanifold of a statistical manifold is statistical which is not true for a lightlike submanifold of a statistical manifold.

Using (20) and the relation (2), it yields

$$\begin{aligned} \nabla_E \bar{\mathfrak{g}}(F, C) + \nabla_E^* \bar{\mathfrak{g}}(F, C) &= \sigma(E, F)u(C) + \sigma(E, C)u(F) \\ &+ \sigma^*(E, F)u(C) + \sigma^*(E, C)u(F) \end{aligned} \quad (23)$$

for all $E, F, C \in \Gamma(T\mathcal{M})$ where u is a 1-form such that $u(E) = \bar{\mathfrak{g}}(E, \mathcal{L})$.

Using (20) and (21), we have the following formulas for the statistical curvature tensor fields

$$\begin{aligned} 2\bar{\mathcal{S}}(E, F)C &= 2\mathcal{S}(E, F)C - \sigma(F, C)A_\mathcal{L}^* E + \sigma(E, C)A_\mathcal{L}^* C \\ &- \sigma^*(F, C)A_\mathcal{L} E + \sigma^*(E, C)A_\mathcal{L} F \\ &+ \{\sigma(F, C)s^*(E) - \sigma(E, C)s^*(F) \\ &+ \sigma^*(F, C)s(E) - \sigma^*(E, C)s(F) \\ &+ (\nabla_E \sigma)(F, C) - (\nabla_F \sigma)(E, C) \\ &+ (\nabla_E^* \sigma^*)(F, C) - (\nabla_F^* \sigma^*)(E, C)\} \mathcal{L}, \end{aligned} \quad (24)$$

$$\begin{aligned} 2\mathcal{S}(E, F)\mathcal{P}C &= 2\mathcal{S}'(E, F)\mathcal{P}C + \rho(E, \mathcal{P}C)A'_\zeta F - \rho(F, \mathcal{P}C)A_\zeta^* E \\ &+ \rho^*(E, \mathcal{P}C)A'_\zeta F - \rho^*(F, \mathcal{P}C)A'_\zeta E \\ &+ \{(\nabla_E \rho)(F, \mathcal{P}C) - (\nabla_F \rho)(E, \mathcal{P}C) \\ &+ (\nabla_E^* \rho^*)(F, \mathcal{P}C) - (\nabla_F^* \rho^*)(E, \mathcal{P}C)\} \zeta \end{aligned} \quad (25)$$

where

$$\begin{aligned} ((\nabla_E^* \sigma)(F, C) &= \nabla_E \sigma(F, C) - \sigma(\nabla_E F, C) - \sigma(F, \nabla_E C), \\ (\nabla_E^* \sigma^*)(F, C) &= \nabla_E^* \sigma(F, C) - \sigma(\nabla_E^* F, C) - \sigma(F, \nabla_E^* C), \\ (\nabla_E \rho)(F, C) &= \nabla_E \rho(F, C) - \rho(\nabla_E F, C) - \rho(F, \nabla_E C), \\ (\nabla_E^* \rho^*)(F, C) &= \nabla_E^* \rho^*(F, C) - \rho^*(\nabla_E F, C) - \rho^*(F, \nabla_E C) \end{aligned} \quad (26)$$

with $\mathcal{S}(E, F)C = \frac{1}{2}\{\mathcal{R}(E, F)C + \mathcal{R}^*(E, F)C\}$ and $\mathcal{S}'(E, F)C = \frac{1}{2}\{\mathcal{R}'(E, F)C + \mathcal{R}'^*(E, F)C\}$ for all $E, F, C \in \Gamma(TM)$.

Furthermore, from (20), we derive

$$\begin{aligned} \sigma(E, \xi_i) &= \rho(E, \zeta_i), & i = 1, 2, 3 \\ \nabla_E \xi_i &= -\psi_i A_{\mathcal{L}} E + s^*(E) \xi_i, & \nabla_E^* \xi_i = -\psi_i A_{\mathcal{L}}^* E + s(E) \xi_i \\ \nabla_E \zeta_i &= -\psi_i A_{\mathcal{L}}' E + s'^*(E) \zeta_i, & \nabla_E^* \zeta_i = -\psi_i A_{\mathcal{L}}'^* E + s'(E) \zeta_i \end{aligned} \quad (27)$$

Definition 3.2. [3, 4] Let $(\mathcal{M}, \mathfrak{g}, \nabla)$ be a lightlike real hypersurface of an almost para-hyperhermitian statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$. It is said that $(\mathcal{M}, \mathfrak{g})$ is

1. totally umbilical with respect to $\bar{\nabla}$ and $\bar{\nabla}^*$ if there exist smooth functions κ and κ^* on a neighborhood \mathcal{U} such that $\sigma(E, F) = \kappa \mathfrak{g}(E, F)$ and $\sigma^*(E, F) = \kappa^* \mathfrak{g}(E, F)$, respectively.
2. totally geodesic with respect to $\bar{\nabla}$ and $\bar{\nabla}^*$ if $\sigma = \sigma^* = 0$
3. screen locally conformal with respect to $\bar{\nabla}$ and $\bar{\nabla}^*$ if the shape operators $\{A_{\mathcal{L}}, A_{\mathcal{L}}'\}$ and $\{A_{\mathcal{L}}^*, A_{\mathcal{L}}'^*\}$ are related by

$$A_{\mathcal{L}} E = \gamma A_{\mathcal{L}}' E, \quad A_{\mathcal{L}}^* E = \gamma^* A_{\mathcal{L}}'^* E, \quad (28)$$

for all $E, F \in \Gamma(TM)$. Here, γ, γ^* are smooth functions on a neighborhood \mathcal{U} in \mathcal{M} which do not vanish.

Definition 3.3. It is said that $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$ of real dimension $4n \geq 8$ is of constant holomorphic sectional curvature \bar{c} if and only if

$$\begin{aligned} \bar{\mathcal{S}}(E, F)C &= \frac{\bar{c}}{4} \{ \bar{\mathfrak{g}}(F, C)E - \bar{\mathfrak{g}}(E, C)F \\ &+ \sum_{i=1}^3 \tau_i [\bar{\mathfrak{g}}(\bar{\mathfrak{X}}_i F, C) \bar{\mathfrak{X}}_i E - \bar{\mathfrak{g}}(\bar{\mathfrak{X}}_i E, C) \bar{\mathfrak{X}}_i F + 2\bar{\mathfrak{g}}(E, \bar{\mathfrak{X}}_i F) \bar{\mathfrak{X}}_i C] \} \end{aligned} \quad (29)$$

holds for E, F, C on $\Gamma(\bar{\mathcal{M}})$.

Lemma 3.4. *Let $(\mathcal{M}, \mathfrak{g}, \nabla)$ be a lightlike real hypersurface of an almost para-hyperhermitian statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$ of constant holomorphic sectional curvature \bar{c} . Then, we conclude that*

$$\begin{aligned} & \frac{\bar{c}}{2} \{ \mu_i(F) \bar{\mathfrak{g}}(\bar{\mathfrak{X}}_i E, C) - \mu_i(E) \bar{\mathfrak{g}}(\bar{\mathfrak{X}}_i F, C) - 2\mu_i(C) \bar{\mathfrak{g}}(E, \bar{\mathfrak{X}}_i F) \} \quad (30) \\ & = 2\bar{\mathfrak{g}}(\bar{\mathcal{S}}(E, F)C, \zeta) = \sigma(F, C)s^*(E) - \sigma(E, C)s^*(F) \\ & + \sigma^*(F, C)s(E) - \sigma^*(E, C)s(F) \\ & + (\nabla_F \sigma)(E, C) - (\nabla_F \sigma)(E, C) \\ & + (\nabla_E^* \sigma^*)(F, C) - (\nabla_F^* \sigma^*)(E, C) \end{aligned}$$

and

$$\begin{aligned} & \frac{\bar{c}}{2} \{ \bar{\mathfrak{g}}(F, C)u(E) - \bar{\mathfrak{g}}(E, C)u(F) + \bar{\mathfrak{g}}(\bar{\mathfrak{X}}_i E, C)\nu_i(F) \quad (31) \\ & - \bar{\mathfrak{g}}(\bar{\mathfrak{X}}_i F, C)\nu_i(E) - 2\bar{\mathfrak{g}}(E, \bar{\mathfrak{X}}_i F)\nu_i(C) \} \\ & = 2\bar{\mathfrak{g}}(\bar{\mathcal{S}}(E, F)C, \mathcal{L}) = 2\bar{\mathfrak{g}}(\bar{\mathcal{S}}(E, F)C, \mathcal{L}) \\ & = -\sigma(F, C)\bar{\mathfrak{g}}(A_{\mathcal{L}}^* E, \mathcal{L}) + \sigma(E, C)\bar{\mathfrak{g}}(A_{\mathcal{L}}^* F, \mathcal{L}) \\ & - \sigma^*(F, C)\bar{\mathfrak{g}}(A_{\mathcal{L}} E, \mathcal{L}) + \sigma^*(E, C)\bar{\mathfrak{g}}(A_{\mathcal{L}} F, \mathcal{L}) \end{aligned}$$

for all $E, F, C \in \Gamma(TM)$, $\mathcal{L} \in \text{ltr}(TM)$ and $\zeta \in \Gamma(\text{rad}(TM))$.

Proof. By taking the inner product with ζ and \mathcal{L} to (24) and using (29), we get (30) and (31), respectively. \square

Proposition 3.5. *Let $(\mathcal{M}, \mathfrak{g}, \nabla)$ be a lightlike real hypersurface of an almost para-hyperhermitian statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$ of constant holomorphic sectional curvature \bar{c} , then we obtain*

$$\begin{aligned} & \frac{\bar{c}}{2} \{ \mu_i(F)\nu_i(E) - \mu_i(E)\nu_i(F) \} = \rho(F, A'_\zeta E) - \rho(E, A'_\zeta F) \quad (32) \\ & + \rho^*(F, A'^*_\zeta E) - \rho^*(E, A'^*_\zeta F) \\ & - \sigma(F, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}}^* E, \mathcal{L}) + \sigma(E, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}}^* F, \mathcal{L}) \\ & - \sigma^*(F, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}} E, \mathcal{L}) + \sigma^*(E, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}} F, \mathcal{L}) \\ & - 2ds(E, F) - 2d^*s^*(E, F) \end{aligned}$$

for all $E, F \in \Gamma(TM)$.

Proof. Putting $C = \zeta$ into relation (31), we get

$$\begin{aligned} \frac{\bar{c}}{2}\{\mu_i(F)\nu_i(E) - \mu_i(E)\nu_i(F)\} &= 2\bar{\mathfrak{g}}(\mathcal{S}(E, F)\zeta, \mathcal{L}) \\ &\quad - \sigma(F, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}}^*E, \mathcal{L}) + \sigma(E, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}}^*F) \\ &\quad - \sigma^*(F, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}}E, \mathcal{L}) + \sigma^*(E, \zeta)\bar{\mathfrak{g}}(A_{\mathcal{L}}F, \mathcal{L}) \\ &\quad - 2ds(E, F) - 2d^*s^*(E, F) \end{aligned} \quad (33)$$

for all $E, F \in \Gamma(T\mathcal{M})$. However, using (20) and (21), we obtain

$$\bar{\mathfrak{g}}(\mathcal{R}(E, F)\zeta, \mathcal{L}) = \bar{\mathfrak{g}}(A_{\zeta}^*F, A_{\mathcal{L}}E) - \bar{\mathfrak{g}}(A_{\zeta}^*E, A_{\mathcal{L}}F) - 2ds(E, F) \quad (34)$$

where $ds(E, F) = \nabla_E s(F) - \nabla_F s(E) - s[E, F]$.

Similarly,

$$\bar{\mathfrak{g}}(\mathcal{R}^*(E, F)\zeta, \mathcal{L}) = \bar{\mathfrak{g}}(A_{\zeta}'F, A_{\mathcal{L}}^*E) - \bar{\mathfrak{g}}(A_{\zeta}'E, A_{\mathcal{L}}^*F) - 2d^*s^*(E, F) \quad (35)$$

with $d^*s^*(E, F) = \nabla_E^* s^*(F) - \nabla_F^* s^*(E) - s^*[E, F]$. Then, from (33), (34) and (35), the assertion follows. \square

Theorem 3.6. *Let $(\mathcal{M}, \mathfrak{g}, \nabla)$ be a lightlike real hypersurface of an almost para-hyperhermitian statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$ of constant holomorphic sectional curvature \bar{c} . If ξ_i , ($i = 1, 2, 3$), are eigenvectors of A_{ζ}' and A_{ζ}^* , then we get*

$$\begin{aligned} \frac{\bar{c}}{4}\{\mu_i(F)u(E) - \mu_i(E)u(F) - 2\tau_i\bar{\mathfrak{g}}(E, \bar{\mathfrak{X}}_iF)\} \\ = E(\beta_i + \beta_i^*)\nu_i(F) - F(\beta_i + \beta_i^*)\nu_i(E) \\ + \beta_i\bar{\mathfrak{g}}(E, (\psi_i A_{\mathcal{L}} + A_{\mathcal{L}}\psi_i)F) \\ + \beta_i^*\bar{\mathfrak{g}}(E, (\psi_i A_{\mathcal{L}}^* + A_{\mathcal{L}}^*\psi_i)F) \\ - 2\bar{\mathfrak{g}}(E, (A_{\mathcal{L}}\psi_i A_{\zeta}' + A_{\mathcal{L}}^*\psi_i A_{\zeta}^*)F) \\ + 2(\beta_i - \beta_i^*)\{s(E)\nu_i(F) - s(F)\nu_i(E)\} \end{aligned} \quad (36)$$

for all $E, F \in \Gamma(T\mathcal{M})$ where $\beta_i = \tau_i\mu_i(A_{\zeta}'\xi_i)$ and $\beta_i^* = \tau_i\mu_i(A_{\zeta}^*\xi_i)$.

Proof. From $A_{\zeta}^*\xi_i = \beta_i^*\xi_i$, $i = 1, 2, 3$ and using relation (27), we obtain

$$\begin{aligned} \bar{\mathfrak{g}}((\nabla_E^* A_{\zeta}^*)F, \xi_i) &= (E\beta_i^*)\nu_i(F) - \beta_i^*\bar{\mathfrak{g}}(F, \psi_i A_{\mathcal{L}}^*E) \\ &\quad + \bar{\mathfrak{g}}(A_{\zeta}^*F, \psi_i A_{\mathcal{L}}^*E), \end{aligned} \quad (37)$$

Similarly, we have

$$\begin{aligned}\bar{\mathfrak{g}}((\nabla_E A'_\zeta)F, \xi_i) &= (E\beta_i)\nu_i(F) - \beta_i\bar{\mathfrak{g}}(F, \psi_i A_\mathcal{L}E) \\ &\quad + \bar{\mathfrak{g}}(A'_\zeta F, \psi_i A_\mathcal{L}E)\end{aligned}\quad (38)$$

So, putting $C = \xi_i$, $i = 1, 2, 3$ into (30) and using equations (37) and (38), the assertion follows. \square

Theorem 3.7. *Let $(\mathcal{M}, \mathfrak{g}, \nabla)$ be a screen locally conformal lightlike real hypersurface of an almost para-hyperhermitian statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \mathfrak{X} = (\mathfrak{X}_i)_{i=1,2,3})$ of constant holomorphic sectional curvature \bar{c} . If the vector fields ξ_i, ζ_i $i = 1, 2, 3$ are eigenvalues of the shape operators $A_\mathcal{L}, A_\mathcal{L}^*$ then, we have*

$$\begin{aligned}\frac{\bar{c}}{2} &= \frac{\lambda_i \lambda_i^* - \lambda_i \alpha_i^*}{\gamma} + \frac{\lambda_i \lambda_i^* - \lambda_i^* \alpha_i}{\gamma^*} \\ &\quad - 2(ds + d^*s^*)(\zeta_i, \xi_i)\end{aligned}\quad (39)$$

where

$$\begin{aligned}A_\mathcal{L}\xi_i &= \lambda_i \xi_i, & A_\mathcal{L}^*\xi_i &= \lambda_i^* \xi_i, \\ A_\mathcal{L}\zeta_i &= \alpha_i \zeta_i, & A_\mathcal{L}^*\zeta_i &= \alpha_i^* \zeta_i.\end{aligned}\quad (40)$$

Proof. Putting $F = \xi_i$ and $E = \zeta_i$, $i = 1, 2, 3$ into relation (32) and using (28), we get

$$\begin{aligned}\frac{\bar{c}}{2}\{\mu_i(\xi)\nu_i(\zeta_i) - \mu_i(\zeta_i)\nu_i(\xi_i)\} &= \rho(\xi_i, A'_\zeta \zeta_i) - \rho(\zeta_i, A'_\zeta \xi_i) \\ &\quad + \rho^*(\xi_i, A'^*_\zeta \zeta_i) - \rho^*(\zeta_i, A'^*_\zeta \xi_i), \\ &\quad - \sigma(\xi_i, \zeta)\bar{\mathfrak{g}}(A_\mathcal{L}^* \zeta_i, \mathcal{L}) + \sigma(\zeta_i, \zeta)\bar{\mathfrak{g}}(A_\mathcal{L}^* \xi_i, \mathcal{L}) \\ &\quad - \sigma^*(\xi_i, \zeta)\bar{\mathfrak{g}}(A_\mathcal{L} \zeta_i, \mathcal{L}) + \sigma^*(\zeta_i, \zeta)\bar{\mathfrak{g}}(A_\mathcal{L} \xi_i, \mathcal{L}) \\ &\quad - 2ds(\zeta_i, \xi_i) - 2d^*s^*(\zeta_i, \xi_i)\end{aligned}$$

Using relations (22) in the above equation and taking into account that \mathcal{M} is screen conformal, we have

$$A_{\zeta}^* \zeta_i = \frac{1}{\gamma^*} A_{\mathcal{L}}^* \zeta_i = \frac{1}{\gamma^*} \alpha^* \zeta_i, \quad A'_{\zeta} \zeta_i = \frac{1}{\gamma} A_{\mathcal{L}} \zeta_i = \frac{1}{\gamma} \alpha \zeta_i$$

Since ξ_i, ζ_i are eigenvalues of $A_{\mathcal{L}}, A_{\mathcal{L}}^*$. From two last equations and the fact that $\bar{\mathfrak{g}}(\xi_i, \zeta_i) = 1, \bar{\mathfrak{g}}(\xi_i, \xi_i) = \bar{\mathfrak{g}}(\mathcal{L}, \mathcal{L}) = 0 = \bar{\mathfrak{g}}(\zeta_i, \zeta_i) = \bar{\mathfrak{g}}(\zeta, \zeta)$, we can state

$$\begin{aligned} \frac{\bar{c}}{2} = & \tau_i \frac{\lambda_i \lambda_i^* - \tau_i \lambda_i \bar{\mathfrak{g}}(A_{\mathcal{L}}^* \zeta_i, \xi_i)}{\gamma} + \frac{\lambda_i \lambda_i^* - \tau_i \lambda_i^* \bar{\mathfrak{g}}(A_{\mathcal{L}} \zeta_i, \xi_i)}{\gamma^*} \\ & - 2(ds + d^* s^*)(\zeta_i, \xi_i). \end{aligned} \quad (41)$$

So, we get the assertion. \square

Corollary 3.8. *Let $(\mathcal{M}, \mathfrak{g}, \nabla)$ be a screen locally conformal lightlike real hypersurface of an almost para-hyperhermitian statistical manifold $(\bar{\mathcal{M}}, \bar{\mathfrak{g}}, \bar{\nabla}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_i)_{i=1,2,3})$ of constant holomorphic sectional curvature \bar{c} . If the vector fields ζ_i, ξ_i $i = 1, 2, 3$ are eigenvalues of the shape operators $A_{\mathcal{L}}, A_{\mathcal{L}}^*$, such that*

$$\begin{aligned} A_{\mathcal{L}} \xi_i &= \lambda_i \xi_i, & A_{\mathcal{L}}^* \xi_i &= \lambda_i^* \xi_i \\ A_{\mathcal{L}} \zeta_i &= \alpha_i \zeta_i, & A_{\mathcal{L}}^* \zeta_i &= \alpha_i^* \zeta_i \end{aligned} \quad (42)$$

then we have

$$2(ds + d^* s^*)(\xi_i, \zeta_j) = 0. \quad (43)$$

Proof. Putting $F = \xi_i$ and $E = \zeta_j$ into relation (32) and using (20), we get the assertion. \square

Example 3.9. Let consider $\bar{\mathcal{M}} = \mathbb{R}_4^8$ equipped with para-hyperhermitian structure $\bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_1, \bar{\mathfrak{X}}_2, \bar{\mathfrak{X}}_3)$ as follows

$$\begin{aligned} \bar{\mathfrak{X}}_1(y_1, y_2, y_3, y_4, y_5, y_6, y_7, y_8) &= (-y_3, y_4, -y_1, y_2, -y_7, y_8, -y_5, y_6) \\ \bar{\mathfrak{X}}_2(y_1, y_2, y_3, y_4, y_5, y_6, y_7, y_8) &= (y_4, y_3, y_2, y_1, y_8, y_7, y_6, y_5) \\ \bar{\mathfrak{X}}_3(y_1, y_2, y_3, y_4, y_5, y_6, y_7, y_8) &= (-y_2, y_1, -y_4, y_3, -y_6, y_5, -y_8, y_7). \end{aligned}$$

with Cartesian coordinate $(y_1, y_2, y_3, y_4, y_5, y_6, y_7, y_8)$ with the metric \bar{g}

$$\bar{g} = -dy_1^2 - dy_2^2 - dy_3^2 - dy_4^2 + dy_5^2 + dy_6^2 + dy_7^2 + dy_8^2.$$

By taking $\frac{\partial}{\partial y_i} = w_i$, we define statistical connections $\bar{\nabla}, \bar{\nabla}^*$ on $\bar{\mathcal{M}}$ as follows

$$\bar{\nabla}_{w_i} w_i = w_i = -\bar{\nabla}_{w_i}^* w_i, \quad i = 1, \dots, 8$$

and other components are zero. Then, $\bar{\mathcal{M}}$ is an almost para-hyperhermitian statistical manifold.

Let \mathcal{M} be a hypersurface of $(\mathbb{R}_4^8, \bar{g}, \mathfrak{X} = (\mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_3))$ such that

$$\begin{aligned} y_1 &= t_1 + \cos \alpha t_5, & y_2 &= t_4 \\ y_3 &= -t_2, & y_4 &= t_3 + t_7 \\ y_5 &= \cos \alpha t_1 - \sin \alpha t_4 + t_5, & y_6 &= \sin \alpha t_1 + \cos \alpha t_4 \\ y_7 &= -\cos \alpha t_2 + \sin \alpha t_3 + t_6, & y_8 &= \sin \alpha t_2 + \cos \alpha t_3 \end{aligned}$$

where $\alpha \in \mathbb{R} - \{\pi + k\pi, k \in \mathbb{Z}\}$. Then, $T\mathcal{M}$ is spanned by

$$\begin{aligned} E_1 &= \partial y_1 + \cos \alpha \partial y_5 + \sin \alpha \partial y_6, & E_2 &= -\partial y_3 - \cos \alpha \partial y_7 + \sin \alpha \partial y_8, \\ E_3 &= \partial y_4 + \sin \alpha \partial y_7 + \cos \alpha \partial y_8, & E_4 &= \partial y_2 - \sin \alpha \partial y_5 + \cos \alpha \partial y_6, \\ E_5 &= \cos \alpha \partial y_1 + \partial y_5, & E_6 &= \partial y_7, \\ E_7 &= \partial y_4 \end{aligned}$$

and we can see that $\mathfrak{X}_1 E_1 = E_2$, $\mathfrak{X}_2 E_1 = E_2$ and $\mathfrak{X}_3 E_1 = E_3$. Considering $E' = \cot \alpha \partial y_2 + \csc \alpha \partial y_6$, we get $\mathcal{L} = \cot \alpha \partial y_2 - \frac{1}{2}(\partial y_1 + \cos \alpha \partial y_5) + (\csc \alpha - \frac{1}{2} \sin \alpha) \partial y_6$. Thus, we have

$$\begin{aligned} \xi_1 &= \cot \alpha \partial y_4 + \frac{1}{2} \partial y_3 + \frac{1}{2} \cos \alpha \partial y_7 + (\csc \alpha - \frac{1}{2} \sin \alpha) \partial y_8, \\ \xi_2 &= \cot \alpha \partial y_3 - \frac{1}{2} \partial y_4 - \frac{1}{2} \cos \alpha \partial y_8 + (\csc \alpha - \frac{1}{2} \sin \alpha) \partial y_7, \\ \xi_3 &= -\cot \alpha \partial y_1 - \frac{1}{2} \partial y_2 - \frac{1}{2} \cos \alpha \partial y_6 - (\csc \alpha - \frac{1}{2} \sin \alpha) \partial y_5 \end{aligned}$$

$$\begin{aligned} \zeta_1 &= -\partial y_3 - \cos \alpha \partial y_7 + \sin \alpha \partial y_8, \\ \zeta_2 &= \partial y_4 + \cos \alpha \partial y_8 + \sin \alpha \partial y_7, & \zeta_3 &= \partial y_2 + \cos \alpha \partial y_6 - \sin \alpha \partial y_5. \end{aligned}$$

Using the Gauss formulas for \mathcal{M} and $\mathfrak{s}(T\mathcal{M})$, we obtain

$$\begin{aligned}
\bar{\nabla}_{E_1}E_1 &= -\partial y_1 + \cos^2\alpha\partial y_5 + \sin^2\alpha\partial y_6, \\
\bar{\nabla}_{E_2}E_2 &= \partial y_3 + \cos^2\alpha\partial y_7 + \sin^2\alpha\partial y_8 \\
\bar{\nabla}_{E_3}E_3 &= \partial y_4 + \sin^2\alpha\partial y_7 + \cos^2\alpha\partial y_8, \\
\bar{\nabla}_{E_4}E_4 &= \partial y_2 + \sin^2\alpha\partial y_5 + \cos^2\alpha\partial y_6, \\
\bar{\nabla}_{E_5}E_5 &= \cos^2\alpha\partial y_1 + \partial y_5, \\
\bar{\nabla}_{E_6}E_6 &= \partial y_7, \\
\bar{\nabla}_{E_7}E_7 &= \partial y_4, \\
\bar{\nabla}_{\xi_1}\xi_1 &= \cot^2\alpha\partial y_4 + \frac{1}{4}\partial y_3 + \frac{1}{4}\cos^2\alpha\partial y_7 + (\csc\alpha - \frac{1}{2}\sin\alpha)\partial y_8 \\
\bar{\nabla}_{\xi_1}\zeta_1 &= \frac{1}{2}\cos^2\alpha\partial y_7 + \sin\alpha(\csc\alpha - \frac{1}{2}\sin\alpha)\partial y_8.
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
\sigma(E_1, E_1) &= \cos^3\alpha + \sin^3\alpha, & \sigma(E_2, E_2) &= 0, \\
\sigma(E_3, E_3) &= 0, & \sigma(E_4, E_4) &= \sin^2\alpha\cos\alpha + \cos^2\alpha\sin\alpha, \\
\sigma(E_5, E_5) &= -\cos^2\alpha + \cos\alpha, & \sigma(E_6, E_6) &= 0, \\
\sigma(E_7, E_7) &= 0.
\end{aligned}$$

We compute

$$\begin{aligned}
\nabla_{\xi_1}\xi_1 &= \cot^2\alpha\partial_4 + \frac{1}{4}\partial y_3 + \frac{1}{4}\cos^2\alpha\partial y_7 + (\csc\alpha - \frac{1}{2}\sin\alpha)\partial y_8(44) \\
\nabla_{\xi_1}\zeta_1 &= \frac{1}{2}\cos^2\alpha\partial y_7 + \sin\alpha(\csc\alpha - \frac{1}{2}\sin\alpha)\partial y_8 \\
\rho(\xi_1, \xi_1) &= \rho(\xi_1, \zeta_1) = 0.
\end{aligned}$$

Thus, \mathcal{M} is a real lightlike hypersurface of $(\mathbb{R}_4^8, \bar{\mathfrak{g}}, \bar{\mathfrak{X}} = (\bar{\mathfrak{X}}_1, \bar{\mathfrak{X}}_2, \bar{\mathfrak{X}}_3), \bar{\nabla})$.

Acknowledgments

The authors would like to express sincerely thanks to the referees for carefully reading the paper and giving helpful comments.

References

- [1] Ö. Aksu, E. Erkan, and M. Gülbahar, Screen invariant lightlike hypersurfaces of almost product-like statistical manifolds and locally product-like statistical manifolds, *Malaya J. Matematik*, 11(4) (2023) 363–377.
- [2] S. Amari, *Differential Geometric Methods in Statistics*, Springer, Berlin, 1985.
- [3] C. Atindogbe and K.L. Duggal, Conformal screen on lightlike hypersurfaces, *Int. J. Pure Appl. Math.* 11(4) (2004) 421–442.
- [4] O. Bahadir and M.M. Tripathi, Geometry of lightlike hypersurfaces of a statistical manifold, *WSEAS Transactions on Mathematics* 22 (2023) 466–474.
- [5] K.L. Duggal, Foliations of lightlike hypersurfaces and their physical interpretation, *Cent. Eur. J. Math.* 10 (2012) 1789–1800.
- [6] K.L. Duggal and C.L. Bejan, *Lightlike Submanifolds of Semi-Riemannian Manifolds and Applications*, Mathematics and its Applications, 364. Kluwer Academic Publication Group, Dordrecht, 1996.
- [7] K.L. Duggal and B. Sahin, *Differential Geometry of Lightlike Submanifolds*, Frontiers in Mathematics, Birkhauser, 2010.
- [8] K. L. Duggal and A. Bejancu, Lightlike submanifolds of semi-Riemannian manifolds, *Acta Appl. Math.* 38 (1995) 197—215.
- [9] H. Furuhashi and I. Hasegawa, *Submanifold Theory in Holomorphic Statistical Manifolds*, in: S. Dragomir, M. H. Shahid, F. R. Al-Solamy (Eds), *Geometry of Cauchy-Riemann Submanifolds*, Springer, Singapore, 2016.
- [10] S. Ianus, L. Ornea and G.E. Vilcu, Submanifolds in Manifolds with Metric Mixed 3-Structures, *Mediterr. J. Math.* 9(1) (2012) 105–128.

- [11] M.B. Kazemi Balgesir and S. Salahvarzi, On statistical submersions from 3-Sasakian statistical manifolds, *Differ. Geom. Appl.* 94 (2024), 102-124.
- [12] A.N. Siddiqui, B.Y. Chen and M.D. Siddiqi, Chen inequalities for statistical submersions between Statistical manifolds, *Int. J. Geom. Methods Models Phys.* 18(4) (2021) 2150049.
- [13] G.E. Vilcu, Para-hyperhermitian structures on tangent bundles, *Proc. Est. Acad. Sci.* 60(3) (2011) 165-173.
- [14] K. Yano and M. Kon, *Structures on Manifolds*, World Scientific, 1984.

Mohammad Bagher Kazemi Balgeshir

Assistant Professor of Mathematics
Department of Mathematics
University of Zanzan
Zanzan, Iran
E-mail: mbkazemi@znu.ac.ir

Sara Miri

Ph.D. in Mathematics
Department of Mathematics
Azarbaijan Shahid Madani University
Tabriz, Iran
E-mail: s.miri@azaruniv.ac.ir

Mohammad Ilmakchi

Associate Professor of Mathematics
Department of Mathematics
Azarbaijan Shahid Madani University
Tabriz, Iran
E-mail: ilmakchi@azaruniv.ac.ir