

Journal of Mathematical Extension
Journal Pre-proof
ISSN: 1735-8299
URL: <http://www.ijmex.com>
Original Research Paper

Existence Results for Some Fractional Stochastic Integro-differential Equations via Measures of Non-compactness

A. R. Yaghoobnia

Lahijan Branch, Islamic Azad University

M. Kazemi*

Ashtian Branch, Islamic Azad University

V. N. Mishra

Indira Gandhi National Tribal University

Abstract. Using fixed point theorems is one method used to prove the existence of solutions in many types of integral equations. This study focuses on applying a generalization of Petryshyn's fixed point theorem to solve a general form of fractional stochastic integro-differential equations in the Banach algebra $C([0, a])$. Besides stating and proving the relevant theorem, the reasons for the superiority of the new method compared some similar methods, were explained. In addition, to confirm the efficiency and check the validity of results, a part of the paper dedicated to solving some stochastic integral equations.

AMS Subject Classification: 47H09, 47H10

Keywords and Phrases: Existence of solution, measures of noncompactness, stochastic Integral equations, fractional calculus, Petryshyn's fixed point theorem

Received: April 2024; Accepted: October 2024

*Corresponding Author

1 Introduction

Integral equations (IEs) are an important tool for scientific expression of many phenomena and modeling of a wide range of scientific processes and have wide applications in various scientific fields such as mathematical physics, economics, biology, scattering theory, mechanics and population dynamics, [13, 19, 39, 40, 41, 47]. The importance of the existence of the solution in such studies cannot be overstated, as many times no analytical solution can be found for such problems.

So far, many researchers have been study in this field and have reflected the results of their research [6, 17]. Employing fixed point theorems (FPTs) to check the existence of solutions in different types of IEs is one of the most important methods used by scientists in this field, for more instance, consider [2, 22, 25, 26]. Some systems are such that to model them, familiarity with fractional integral equations (FIEs) is necessary. Also, in the solution's existence, we can refer to the works done in [8, 32, 48, 50], and other classes of these equations in [4, 34, 49], which all are based on FPTs. Some phenomena include random parameters that leads to encountering stochastic IEs [38, 46]. Such systems have more complexities and it is important to make sure they have solution. Methods based on FPTs are some studies that researchers have done to ensure the existence of solutions in different classes of such equations [12, 16, 27, 42]. There are equations that contain a combination of random parameters and derivative of fractional order. Such complex equations can be found in [18, 23].

The introduction and study on the existence of the solution of fractional functional IE in the Riemann-Liouville (RL) sense, using FPT in Banach algebra is given in [5, 9, 31]. In 2021, Samei *et al.* investigated the following singular fractional q -integro-differential equation involving Caputo fractional q -derivative, for $0 < s < 1$,

$${}^C D_q^\sigma y(s) = g\left(s, y(s), y'(s), {}^C D_q^\zeta y(s), \int_0^s y(r)\beta(r) dr\right), \quad (1)$$

under boundary conditions $y(0) = 0$ and $y(1) = {}^C D_q^\eta y(\tau)$, where $y \in C([0, 1])$, $\sigma \in [1, 2)$, $\zeta, \eta, \tau \in (0, 1)$, $\beta \in L^1([0, 1])$ is nonnegative with $\|\beta\|_1 = m$ and $g(s, y_1, y_2, y_3, y_4)$ is singular at some points of s [44].

Aydogan in [7] considered the following k -dimensional hybrid differential system

$$\left\{ \begin{array}{l} {}^C D_q^\sigma \left(m_1(s) \left(\frac{y_1(s)}{l_1(s, y_1(s))} \right)' - \tilde{m}_1(s) \tilde{h}_1(y_1(s)) \right) + q_1(s) y_1(s) \\ \quad = f_1(s) h_1(y_1(s)), \\ {}^C D_q^\sigma \left(m_2(s) \left(\frac{y_2(s)}{l_2(s, y_2(s))} \right)' - \tilde{m}_2(s) \tilde{h}_2(y_2(s)) \right) + q_2(s) y_2(s) \\ \quad = f_2(s) h_2(y_2(s)), \\ \dots \\ {}^C D_q^\sigma \left(m_k(s) \left(\frac{y_k(s)}{l_k(s, y_k(s))} \right)' - \tilde{m}_k(s) \tilde{h}_k(y_k(s)) \right) + q_k(s) y_k(s) \\ \quad = f_k(s) h_k(y_k(s)), \end{array} \right.$$

under the sigma boundary value conditions

$$\left. \left(\frac{y_i(s)}{l_i(s, y_i(s))} \right)' \right|_{s=0} = \frac{\tilde{m}_i(s)}{m_i(s)} \tilde{h}_i(y_i(s)) \Big|_{s=0}, \quad 1 \leq i \leq k,$$

and $\sum_{i=1}^k \zeta_i \left(\frac{y_i(a_i)}{l_i(a_i, y_i(a_i))} \right) = \rho_i \sum_{i=1}^k \frac{y_i(z_i)}{l_i(z_i, y_i(z_i))}$. Bhupeshwar *et al.* first, focused on examining the existence and uniqueness of solutions Ψ -Hilfer FDI

$${}^H D^{\sigma, \zeta; \Psi} y(s) = g(s, y(s)), \quad s \in [\mathfrak{s}_1, \mathfrak{s}_2] \subset \mathbb{R}^{>0}, \quad (2)$$

under conditions $y(\mathfrak{s}_1^+) = y'(\mathfrak{s}_1^+)$ and $y(u) = K I^{\sigma; \Psi} y(s)$, with $\eta = \sigma + \zeta(3 - \sigma)$, where $2 < \sigma < 3$, $K \in \mathbb{R}$, $I^{\sigma; \Psi} y(s)$ is the Ψ -RL FI of order σ , and in the second stage, provided two distinct existence results for Ψ -Hilfer FDI (2) via new conditions

$$\left\{ \begin{array}{l} y_\Psi^{(j)}(\mathfrak{s}_1) = 0, \quad j = 0, 1, 2, \dots, (n-2), \quad y_\Psi^{(j)} = \left(\frac{d}{\Psi'(s) ds} \right)^j y(s), \\ y_\Psi^{n-1}(u) = y_u \in \mathbb{R}, \end{array} \right.$$

where $g(\cdot, y(\cdot)) \in C([0, 1])$ [11].

In this study, we examine the existence of the solution of following

fractional stochastic integro-differential equations (FSIDEs),

$$\begin{aligned} {}^C D^\sigma (y(s) + g(s, y(s))) &= f(s, y(\alpha(s))) \\ &+ F\left(s, y(\beta(s)), \int_0^s k_1(s, t, y(\theta(t))) ds, \right. \\ &\left. \int_0^s k_2(s, t, y(\mu(t))) dW(t)\right), \end{aligned} \quad (3)$$

for $s \in I_a := [0, a]$, under the initial conditions

$$y^{(i)}(0) = x_i, \quad i = 0, 1, \dots, n-1, \quad (4)$$

where, $y \in C(I_a)$ as the analytical solution of (3) is unknown and all other functions are known stochastic processes defined on the some probability space (Ω, \mathcal{F}, P) , and $W(s)$ is the Brownian motion. Also, $y^{(i)}(0)$ is the i -th order of derivative of continuous function x at point 0 and x_i 's are constant. In addition $\alpha, \theta, \mu \in C(I_a)$, $f, g \in C(I_a \times \mathbb{R})$, and $k_1, k_2 \in C(I_a \times I_a \times \mathbb{R})$ are continuous functions. The development of the concept of measures of non-compactness (M.N.C) was first done by Kuratowski [30]. Later, other researchers used this concept in investigating the existence of different types of solutions for the IEs [3, 10, 15, 33]. This research examines the existence of a solution to FSIDE (3) by applying the concept of M.N.C and in this way, the FPT of Petryshyn is used.

2 Auxiliary Facts and Notations

In this section we review some definitions and theorems, by stating some auxiliary facts and notations. First, we provide some preliminary concepts of fractional calculus. Then some basic introductions about stochastic calculations, and in the next subsection about FPT of Petryshyn, which depend on the concept of M.N.C, brief explanations will be provided.

2.1 Fractional calculus

Definition 2.1 ([28]). The RL FI of order $\sigma > 0$ of a function ξ , is defined as

$$I^\sigma \xi(\tau) = \int_0^\tau \frac{(\tau-\mu)^{\sigma-1}}{\Gamma(\sigma)} \xi(\mu) d\mu, \quad \tau > 0.$$

Of course, to learn about the properties of the RL derivative, you can see [28]. In this article, the definition of Caputo derivative is considered, which can better model the phenomenon and is compatible with the initial conditions of the problems.

Definition 2.2 ([28]). The Caputo derivative of fractional order $\sigma \geq 0$ for a function $\xi(\tau)$ is defined by

$$({}^C D^\sigma \xi)(\tau) = \int_0^\tau \frac{(\tau-\mu)^{n-\sigma-1}}{\Gamma(n-\sigma)} \tau^{(n)}(\mu) d\mu,$$

for $n-1 < \sigma \leq n$, $n \in \mathbf{N}$.

Lemma 2.3 ([28]). Let $\sigma > 0$ and $n = [\sigma] + 1$. For two fractional operators defined above, the following properties yield

- (i) $(I^\sigma {}^C D^\sigma \xi)(\tau) = \xi(\tau) - \sum_{i=0}^{n-1} \frac{\xi^{(i)}(0)}{i!} \tau^i;$
- (ii) $({}^C D^\sigma I^\sigma \xi)(\tau) = \xi(\tau).$

2.2 Stochastic calculus

Systems types have been described and evolved using DEs and IEs, since their inception, based on their applications (i.e. economic, mechanical and social systems). These equations applied to model phenomena that in part deal with movement. Stochastic Differential Equations (SDEs) is a new branch of mathematics that defines the characteristics of random motion based on very broad mathematical foundations. Mathematical models involving measuring uncertainty, are key to the solution and play an important part in the branch of science and industry, which is why scientists use SDEs as needed in systems modeling. The SDEs are equations in which one, or more terms are random processes. Therefore, the solution of SDEs may also be of

the type of stochastic processes that despite the similarity to the methods of solving ordinary DEs, there are differences. We studied the basic concepts of this discussion using the concept of Brownian motion.

Definition 2.4. ([29]) Brownian motion $W(s)$ which is the following properties is a stochastic process.

- a) For $0 \leq s_1 < s_2 < \dots < s_n$, the increments

$$W(s_1), W(s_2) - W(s_1), \dots, W(s_n) - W(s_{n-1}),$$

are independent of the path;

- b) $W(s) - W(t)$ having mean and variance 0 and variance $s - t$, respectively, has a normal distribution, as a result $W(s)$ has normal distribution with mean and variance 0 and variance s ;
- c) The $W(s)$, for $s \geq 0$ is a continuous functions.

The definition in part (a), (b) and (c), assumes the start of movement from s . The condition $P(W(0) = 0) = 0$ standardizes Brownian motion where it start at 0.

Before explaining the next theorem which implies the existence of a SI, it is necessary to state the following definition.

Definition 2.5 ([29]). When for all s , $Y(s)$ be \tilde{F}_s -measurable, the process Y is called adapted to the filtration $\tilde{F} = (\tilde{F}_s)$.

Theorem 2.6 ([29]). *If Y be a process that satisfies the continuous adapted condition, then the $\int_0^T Y(s) dW(s)$ exists.*

If Brownian motion was derivable everywhere, its integral would not be a problem, but considering that it is not derivable anywhere, therefore the SI cannot be calculated by normal methods. The common method for calculating the SI is to use the integration by parts method, which converts the SI into a computable normal or simple integral. So that for the differentiable and bounded function ϕ , we have [36]:

$$\int_0^s \phi(t) dW(t) = \phi(s)W(s) - \int_0^s W(t)\phi'(t) dt, \quad 0 \leq s \leq 1, \quad (5)$$

which is an alternative method for calculating SIs.

2.3 Petryshyn's fixed-point theorem

Here, we employ the symbol E for *Real Banach space*, the symbol \bar{B}_r for *Closed ball with center 0 and radius r* , the symbol $\partial\bar{B}_r$ for *Sphere in E around 0 with radius $r > 0$* , and finally the symbol $C(I_a)$ for *Space of all continuous and real-valued functions on $I_a = [0, a]$* . We recall some definitions and theorems that are required for the sequel.

Definition 2.7 ([30]). Let $Y \subset E$ be a bounded set, then $\alpha(Y) = \inf \{ \rho > 0 : Y \text{ can be covered by a finite number of sets with diameter } \leq \rho \}$, is said to be the Kuratowski M.N.C.

Definition 2.8 ([21]). Let $Y \subset E$ be a bounded set, then the Hausdorff M.N.C is given by $\mu(Y) = \inf \{ \rho > 0 : Y \text{ has a finite } \rho\text{-net in } E \}$.

Theorem 2.9 ([21]). *Let $Y \subset E$ be a bounded set, then the M.N.C α and μ fulfill $\mu(Y) \leq \alpha(Y) \leq 2\mu(Y)$.*

The space $C(I_a)$ is a Banach space under the norm $\|y\| = \sup \{ |y(s)| : s \in I_a \}$, and we shall write the modulus of continuity of a function $y \in C(I_a)$ as

$$\omega(y, \rho) = \sup \{ |y(s) - y(t)| : |s - t| \leq \rho \}.$$

Since y is uniformly continuous on I_a , we have $\omega(y, \rho) \rightarrow 0$, as $\rho \rightarrow 0$.

Theorem 2.10 ([21]). *In Hausdorff M.N.C, for all bounded sets $Y \subset C(I_a)$*

$$\mu(Y) = \lim_{\rho \rightarrow 0} \left\{ \sup \omega(y, \rho) : y \in Y \right\}. \quad (6)$$

Definition 2.11. [35] Let $Q : E \rightarrow E$ be a continuous map. Q is said to be a k -set contraction if for all $Y \subset C(I_a)$ be bounded, $Q(Y)$ is bounded and $\alpha(QY) \leq k\alpha(Y)$, $0 < k < 1$. Moreover, Q is said to be condensing (densifying) map if $\alpha(QY) < \alpha(Y)$.

Note that, a k -set contraction with $0 < k < 1$ yields condensing (densifying) but not vice versa.

Theorem 2.12 ([37, 45]). *Suppose that $Q : \bar{B}_r \rightarrow E$ is a densifying mapping that satisfies the boundary condition,*

(P) *if $Q(Y) = kY$, for some Y in $\partial\bar{B}_r$ then $k \leq 1$.*

Then the set of fixed points of Q in \bar{B}_r is nonempty.

3 Main Results

In this section, we examine the solvability of the FSIDE (3). Because of the continuity of g and f , we apply the operator I^σ on sides of Eq. (3),

$$\begin{aligned} y(s) &= \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} s^i - g(s, y(s)) \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{f(t, y(\alpha(t)))}{(s-t)^{1-\sigma}} dt \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{F(t, y(\beta(t)), (H_1 y)(t), (H_2 y)(t))}{(s-t)^{1-\sigma}} dt, \end{aligned} \quad (7)$$

where

$$\begin{aligned} (H_1 y)(s) &= \int_0^s k_1(s, t, y(\theta(t))) dt, \\ (H_2 y)(s) &= \int_0^s k_2(s, t, y(\mu(t))) dW(t). \end{aligned}$$

The Eq. (3) is equivalent to the FSIE (7). This means every solution of Eq. (7) is also a solution of Eq. (3), and vice versa. Next, we consider the following conditions for Eq. (7):

H1) $g, f \in C(I_a \times \mathbb{R})$, $F \in C(I_a \times \mathbb{R} \times \mathbb{R} \times \mathbb{R})$, $k_1, k_2 \in C(I_a \times I_a \times \mathbb{R})$, and $\alpha, \beta, \theta, \mu : I_a \rightarrow I_a$ are continuous;

H2) There exist nonnegative constants k, c_1, c_2, c_3, c_4 , and $k < 1$ such that $|g(s, u) - g(s, \bar{u})| \leq k|u - \bar{u}|$, and

$$|F(s, u, v, w) - F(s, \bar{u}, \bar{v}, \bar{w})| \leq c_1|u - \bar{u}| + c_2|v - \bar{v}| + c_3|w - \bar{w}|;$$

H3) (Bounded condition) There exists nonnegative r_0 such that

$$\sup \left\{ L + A + \frac{M_1 a^\sigma}{\Gamma(1+\sigma)} + \frac{M_2 a^\sigma}{\Gamma(1+\sigma)} \right\} \leq r_0, \quad (8)$$

where

$$\begin{aligned} L &= \sup \left\{ \left| \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} s^i \right| : \forall s \in I_a \right\}, \\ A &= \sup \left\{ |g(s, u)| : \forall s \in I_a, u \in [-r_0, r_0] \right\}, \\ M_1 &= \sup \left\{ |f(s, u)| : \forall s \in I_a, u \in [-r_0, r_0] \right\}, \\ M_2 &= \sup \left\{ |F(s, u, v, w)| : \forall s \in I_a, u \in [-r_0, r_0], \right. \\ &\quad \left. |v| \leq aA_1, |w| \leq \lambda B \right\}, \end{aligned}$$

and

$$\begin{aligned} A_1 &= \sup \left\{ |k_1(s, t, u)| : \forall s, t \in I_a, u \in [-r_0, r_0] \right\}, \\ B &= \sup \left\{ |k_2(s, t, u)| : \forall s, t \in I_a, u \in [-r_0, r_0] \right\}, \\ \lambda &= \sup \left\{ |W(s)| : \forall s \in I_a \right\}. \end{aligned}$$

Theorem 3.1. *By conditions (H1)-(H3) on $E = C(I_a)$, FSIDE. (3) has at least one solution.*

Proof. We define the operator $Q : B_{r_0} \rightarrow C(I_a)$, as follows

$$\begin{aligned} (Qy)(s) &= \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} s^i - g(s, y(s)) \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{f(t, y(\alpha(t)))}{(s-t)^{1-\sigma}} dt \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{F(t, y(\beta(t)), (H_1 y)(t), (H_2 y)(t))}{(s-t)^{1-\sigma}} dt. \end{aligned}$$

We will demonstrate that the operator Q is continuous on the ball B_{r_0} . Take arbitrary $x, y \in B_{r_0}$ and $\varepsilon > 0$ such that $\|x - y\| \leq \varepsilon$, then for

$s \in I_a$, we get

$$\begin{aligned}
|(Qy)(s) - (Qx)(s)| &\leq |g(s, x(s)) - g(s, y(s))| \\
&+ \frac{1}{\Gamma(\sigma)} \int_0^s \frac{|f(t, y(\alpha(t))) - f(t, x(\alpha(t)))|}{(s-t)^{1-\sigma}} dt \\
&+ \frac{1}{\Gamma(\sigma)} \int_0^s \frac{1}{(s-t)^{1-\sigma}} \left[\left| F(t, y(\beta(t)), (H_1y)(t), (H_2y)(t)) \right. \right. \\
&\quad \left. \left. - F(t, x(\beta(t)), (H_1x)(t), (H_2x)(t)) \right| \right] dt \\
&\leq k \|y - x\| + \frac{s^\sigma}{\Gamma(1+\sigma)} \omega(f, \omega(\alpha, \varepsilon)) \\
&+ \frac{1}{\Gamma(\sigma)} \int_0^s \frac{1}{(s-t)^{1-\sigma}} \left[\left| F(t, y(\beta(t)), (H_1y)(t), (H_2y)(t)) \right. \right. \\
&\quad \left. \left. - F(t, x(\beta(t)), (H_1y)(t), (H_2y)(t)) \right| \right] dt \\
&+ \frac{1}{\Gamma(\sigma)} \int_0^s \frac{1}{(s-t)^{1-\sigma}} \left[\left| F(t, x(\beta(t)), (H_1y)(t), (H_2y)(t)) \right. \right. \\
&\quad \left. \left. - F(t, x(\beta(t)), (H_1x)(t), (H_2y)(t)) \right| \right] dt \\
&+ \frac{1}{\Gamma(\sigma)} \int_0^s \frac{1}{(s-t)^{1-\sigma}} \left[\left| F(t, x(\beta(t)), (H_1x)(t), (H_2y)(t)) \right. \right. \\
&\quad \left. \left. - F(t, x(\beta(t)), (H_1x)(t), (H_2x)(t)) \right| \right] dt \\
&\leq k \|y - x\| + \frac{s^\sigma}{\Gamma(1+\sigma)} \omega(f, \omega(\alpha, \varepsilon)) + \frac{c_1 s^\sigma}{\Gamma(1+\sigma)} \|y - x\| \\
&\quad + \frac{c_2 a s^\sigma}{\Gamma(1+\sigma)} \omega(k_1, \varepsilon) + \frac{c_3 \lambda s^\sigma}{\Gamma(1+\sigma)} \omega(k_2, \varepsilon),
\end{aligned}$$

where for $\sigma > 0$, we define

$$\begin{aligned}
\omega(f, \varepsilon) &= \sup \left\{ |f(t, y) - f(t, x)| : t \in I_a, \right. \\
&\quad \left. y, x \in [-r_0, r_0], \|y - x\| \leq \varepsilon \right\}, \\
\omega(k_i, \varepsilon) &= \sup \left\{ |k_i(s, t, y) - k_i(s, t, x)| : s, t \in I_a, \right. \\
&\quad \left. y, x \in [-r_0, r_0], \|y - x\| \leq \varepsilon \right\}, \quad i = 0, 1.
\end{aligned}$$

Since the functions $f = f(t, y)$ and $k = k(s, t, y)$ are uniformly continuous on $I_a \times \mathbb{R}$ and $I_a \times I_a \times \mathbb{R}$, we indicate that $\omega(f, \omega(\alpha, \varepsilon)) \rightarrow 0$,

$\omega(k_1, \varepsilon) \rightarrow 0$ and $\omega(k_2, \varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. Consequently, the operator Q is continuous on B_{r_0} . In the following, we prove the operator Q fulfills densifying condition in view of μ . To do this, we take arbitrary $\varepsilon > 0$ and assumed that $x \in Y \subset C(I_a)$ is a bounded set. Here for $s_1, s_2 \in I_a$ such that $s_1 \leq s_2$ while $s_2 - s_1 \leq \varepsilon$, gives

$$\begin{aligned}
|(Qy)(s_2) - (Qy)(s_1)| &= \left| \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} s_2^i \right. \\
&\quad - g(s_2, y(s_2)) + \frac{1}{\Gamma(\sigma)} \int_0^{s_2} \frac{f(t, y(\alpha(t)))}{(s_2-t)^{1-\sigma}} dt \\
&\quad + \frac{1}{\Gamma(\sigma)} \int_0^{s_2} \frac{F(t, y(\beta(t)), (H_1y)(t), (H_2y)(t))}{(s_2-t)^{1-\sigma}} dt \\
&\quad - \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} s_1^i \\
&\quad + g(s_1, y(s_1)) - \frac{1}{\Gamma(\sigma)} \int_0^{s_1} \frac{f(t, y(\alpha(t)))}{(s_1-t)^{1-\sigma}} dt \\
&\quad \left. - \frac{1}{\Gamma(\sigma)} \int_0^{s_1} \frac{F(t, y(\beta(t)), (H_1y)(t), (H_2y)(t))}{(s_1-t)^{1-\sigma}} dt \right| \\
&\leq \left| \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} (s_2^i - s_1^i) \right| + |g(s_1, y(s_1)) \\
&\quad - g(s_2, y(s_2))| + \frac{1}{\Gamma(\sigma)} \left| \int_0^{s_1} \frac{f(t, y(\alpha(t)))}{(s_2-t)^{1-\sigma}} dt \right. \\
&\quad \left. + \int_{s_1}^{s_2} \frac{f(t, y(\alpha(t)))}{(s_2-t)^{1-\sigma}} dt + \int_0^{s_1} \frac{f(t, y(\alpha(t)))}{(s_1-t)^{1-\sigma}} dt \right| \\
&\quad + \frac{1}{\Gamma(\sigma)} \left| \int_0^{s_1} \frac{F(t, y(\beta(t)), (H_1y)(t), (H_2y)(t))}{(s_2-t)^{1-\sigma}} dt \right. \\
&\quad \left. + \int_{s_1}^{s_2} \frac{F(t, y(\beta(t)), (H_1y)(t), (H_2y)(t))}{(s_2-t)^{1-\sigma}} dt \right. \\
&\quad \left. + \int_0^{s_1} \frac{F(t, y(\beta(t)), (H_1y)(t), (H_2y)(t))}{(s_1-t)^{1-\sigma}} dt \right| \\
&\leq |g(s_1, y(s_1)) - g(s_1, y(s_2))| + |g(s_1, y(s_2)) \\
&\quad - g(s_2, y(s_2))| + \frac{1}{\Gamma(\sigma)} \int_0^{s_1} \left| \frac{f(t, y(\alpha(t)))}{(s_2-t)^{1-\sigma}} \right.
\end{aligned}$$

$$\begin{aligned}
& -\frac{f(t,y(\alpha(t)))}{(s_1-t)^{1-\sigma}} \Big| dt + \frac{1}{\Gamma(\sigma)} \int_{s_1}^{s_2} \left| \frac{f(t,y(\alpha(t)))}{(s_2-t)^{1-\sigma}} \right| dt \\
& + \frac{1}{\Gamma(\sigma)} \int_0^{s_1} \left| \frac{F(t,y(\beta(t)),(H_1y)(t),(H_2y)(t))}{(s_2-t)^{1-\sigma}} \right. \\
& \quad \left. - \frac{F(t,y(\beta(t)),(H_1y)(t),(H_2y)(t))}{(s_1-t)^{1-\sigma}} \right| dt \\
& + \frac{1}{\Gamma(\sigma)} \int_{s_1}^{s_2} \left| \frac{F(t,y(\beta(t)),(H_1y)(t),(H_2y)(t))}{(s_2-t)^{1-\sigma}} \right| dt.
\end{aligned}$$

For simplicity we use the following notation:

$$\omega_g(I_a, \varepsilon) = \sup \left\{ |g(s, y) - g(\bar{s}, y)| : |s - \bar{s}| \leq y \in [-r_0, r_0] \right\},$$

and using the above relation we get

$$\begin{aligned}
|(Qy)(s) - (Qx)(s)| & \leq k\omega(y, \varepsilon) + \omega_g(I_a, \varepsilon) \\
& + \frac{M_1}{\Gamma(1+\sigma)} \{s_1^\sigma - s_2^\sigma + (s_2 - s_1)^\sigma\} + \frac{M_1}{\Gamma(1+\sigma)} (s_2 - s_1)^\sigma \\
& + \frac{M_2}{\Gamma(1+\sigma)} \{s_1^\sigma - s_2^\sigma + (s_2 - s_1)^\sigma\} + \frac{M_2}{\Gamma(1+\sigma)} (s_2 - s_1)^\sigma \\
& \leq k\omega(x, \varepsilon) + \omega_g(I_a, \varepsilon) + \frac{3\varepsilon^\sigma M_1}{\Gamma(1+\sigma)} + \frac{3\varepsilon^\sigma M_2}{\Gamma(1+\sigma)}.
\end{aligned}$$

Taking limit as $\varepsilon \rightarrow 0$, we obtain $\omega(Qy, \varepsilon) \leq k\omega(y, \varepsilon)$, for $y \in Y$. Therefore, $\mu(QY) \leq k\mu(Y)$. Now, we get Q is a condensing mapping with constant $k < 1$. It remains to verify condition (P) of Theorem 2.12. Suppose $y \in \partial \bar{B}_{r_0}$. If $Qy = ky$ then we have $kr_0 = k\|y\| = \|Qy\|$ and by condition (H3), we concluded that

$$\begin{aligned}
|Qy(s)| & = \left| \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} s^i - g(s, y(s)) \right. \\
& \quad \left. + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{f(t,y(\alpha(t)))}{(s-t)^{1-\sigma}} dt \right. \\
& \quad \left. + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{F(t,y(\beta(t)),(H_1y)(t),(H_2y)(t))}{(s-t)^{1-\sigma}} dt \right| \leq r_0,
\end{aligned}$$

hence $\|Qy\| \leq r_0$, which gives $k \leq 1$. \square

The following corollary which is the main results of Dadsetadi *et al.* [14], would be obtained from Theorem 3.1.

Corollary 3.2 ([14]). *Suppose*

M1) $g \in C(I_a \times \mathcal{R})$, $f \in C(I_a \times \mathcal{R})$, $F \in C(I_a \times \mathcal{R}^2)$, $k \in C(I_a^2 \times \mathcal{R})$
and, $\mu : I_a \rightarrow I_a$ are continuous;

M2) *There exist non negative constants k_1, k_2, c_1, c_2 , and c_3 , so that $k_1 < 1$,*

$$|g(\vartheta, \omega_1) - g(\vartheta, \varpi_1)| \leq k_1 |\omega_1 - \varpi_1|,$$

and

$$|F(\vartheta, \omega_1, \omega_2) - F(\vartheta, \varpi_1, \varpi_2)| \leq c_1 |\omega_1 - \varpi_1| + c_2 |\omega_2 - \varpi_2|;$$

M3) $\exists \delta_0 \geq 0$ such that

$$\sup \left\{ L + A + \frac{M_1 a^\sigma}{\Gamma(1+\sigma)} + \frac{M_2 a^\sigma}{\Gamma(1+\sigma)} \right\} \leq \delta_0,$$

where,

$$L = \sup \left\{ \left| \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta^i \right| : \forall \vartheta \in I_a \right\};$$

$$A = \sup \left\{ |g(\vartheta, \omega_1)| : \forall \vartheta \in I_a, \omega_1 \in [-\delta_0, \delta_0] \right\};$$

$$M_1 = \sup \left\{ |f(\vartheta, \omega_1)| : \forall \vartheta \in I_a, \omega_1 \in [-\delta_0, \delta_0] \right\};$$

$$M_2 = \sup \left\{ |F(\vartheta, \omega_1, \omega_2)| : \forall \vartheta \in I_a, \omega_1 \in [-\delta_0, \delta_0], |\omega_2| \leq aB \right\};$$

$$B = \sup \left\{ |k(\vartheta, \ell, \omega_1)| : \forall \vartheta, \ell \in I_a, \omega_1 \in [-\delta_0, \delta_0] \right\}.$$

Then

$$\begin{aligned} {}^C D^\sigma (y(\vartheta) + g(\vartheta, y(\vartheta))) &= f(\vartheta, y(\vartheta)) \\ &+ F\left(\vartheta, y(\vartheta), \int_0^\vartheta k(\vartheta, \ell) H(y(\mu(\ell))) d\ell\right), \quad \vartheta \in I_a, \end{aligned} \quad (9)$$

with the initial conditions

$$y^{(i)}(0) = y_i, \quad i = 0, 1, \dots, n-1, \quad (10)$$

has at least a solution in I_a .

Proof. It is clear that Eq. (9) is a particular case of Eq. (3). Here $\alpha(\vartheta) = \beta(\vartheta) = \varsigma(\vartheta) = \vartheta$, $k(\vartheta, \ell, y(\mu(\ell))) = k(\vartheta, \ell)H(y(\mu(\ell)))$. By employing RL fractional integrating and Lemma 2.3, Eq. (9) changes into

$$y(s) = \sum_{i=0}^{n-1} \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} \vartheta^i - g(\vartheta, y(\vartheta)) \\ + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{f(\ell, y(\ell))}{(\vartheta - \ell)^{1-\sigma}} d\ell + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{F(\ell, y(\ell), (Hy)(\ell))}{(\vartheta - \ell)^{1-\sigma}} d\ell.$$

The proof is connected to Theorem 3.1, so we can drop these parts. \square

Remark 3.3. The above Corollary is the main result of [14], which was proved here using FPT of Petryshyn simpler and with fewer conditions, and this is the advantage of using Petryshyn's theorem.

Corollary 3.4. *If $g(s, y(s)) = k_1 \equiv 0$, $\alpha(s) = \mu(s) = s$ and $F(s, u, v, w) = w$, then Eq. (3) has the following form, which was studied in [24],*

$${}^C D_s^\sigma y(s) = f(s, y(s)) + \int_0^s k_2(s, y(t)) dW(t), \quad (11)$$

with $\sigma \in (0, 1)$ and the initial condition $y(0) = y_0$, where

$${}^C D_s^\sigma \psi(z) = \frac{1}{\Gamma(1-\sigma)} \int_0^s \frac{\psi'(z)}{(s-z)^\sigma} dz, \quad s \in [0, a].$$

It is well-known that Eq. (11) is equivalent to the following FSIE with a weakly singular kernel of the form

$$y(s) = y_0 + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{f(\zeta, y(\zeta))}{(s-\zeta)^{1-\sigma}} d\zeta \\ + \frac{1}{\Gamma(\sigma)} \int_0^s \int_0^\zeta \frac{k_2(\zeta, y(t))}{(s-\zeta)^{1-\sigma}} dW(t) d\zeta.$$

4 Examples

Here, we provide two examples to confirm the efficiency and check the validity of the main results.

Example 4.1. Consider following \mathbb{FSDDE} on $C(I_a)$, $a = 1$,

$$\begin{aligned} {}^C D^\sigma \left(y(s) + \frac{\cos(s)}{\sqrt{64+s^2}} y(s) \right) &= \frac{e^{-4s}}{10+s^2} \sin \left(\frac{1}{1+s} \right) + \frac{y(1-s)\sqrt{s}}{7} \\ &+ \frac{3s^2 y(s^2)}{5+5s^2} + \frac{e^{-s}}{6+5s} \int_0^s \frac{1}{1+s^3} \left[1 + \int_0^t \frac{\ln(1+|y(\xi)|)}{\sqrt{4+\xi+t}} d\xi \right] dt \\ &+ \frac{1}{5} \int_0^s \frac{e^{-3st}}{2+s^2+\ln(1+t)} \sin(y(t)) dW(t), \end{aligned} \quad (12)$$

under the initial conditions $y^{(i)}(0) = y_i$, $i = 0, 1$ for $\sigma = \frac{5}{4}, \frac{3}{2}, \frac{7}{4}$. Eq. (12) is a particular form of Eq. (3) such that $n = 2$,

$$\begin{aligned} g(s, y(s)) &= \frac{\cos(s)}{\sqrt{64+s^2}} y(s), \\ f(s, y(\alpha(s))) &= \frac{e^{-4s}}{10+s^2} \sin \left(\frac{1}{1+s} \right) + \frac{y(1-s)\sqrt{s}}{7}, \\ F(s, u, v, w) &= \frac{3s^2 u}{5+5s^2} + \frac{e^{-s}}{6+5s} v + \frac{1}{5} w, \end{aligned}$$

and

$$\begin{aligned} v &= \int_0^s \frac{1}{1+s^3} \left[1 + \int_0^t \frac{\ln(1+|y(\xi)|)}{\sqrt{4+\xi+t}} d\xi \right] dt, \\ w &= \int_0^s \frac{e^{-3st}}{2+s^2+\ln(1+t)} \sin(y(t)) dW(t). \end{aligned}$$

It can be seen that $|g(s, u) - g(s, \bar{u})| \leq \frac{1}{8}|u - \bar{u}|$ and

$$|F(s, u, v, w) - F(s, \bar{u}, \bar{v}, \bar{w})| \leq \frac{3}{5}|u - \bar{u}| + \frac{1}{6}|v - \bar{v}| + \frac{1}{5}|w - \bar{w}|.$$

Here $k = \frac{1}{8} < 1$, $c_1 = \frac{3}{5}$, $c_2 = \frac{1}{6}$, $c_3 = \frac{1}{5}$. So, the conditions (H1) and (H2) hold. Moreover, for $\|y\| \leq r_0$, $r_0 > 0$ and $y_0 = 0$, $y_1 = 1$, we have

$$\begin{aligned} |y(s)| &= \left| \sum_{i=0}^1 \frac{y^{(i)}(0) + g^{(i)}(0, y_0)}{i!} s^i - g(s, y(s)) \right. \\ &+ \frac{1}{\Gamma(\sigma)} \int_0^s \frac{f(t, y(\alpha(t)))}{(s-t)^{\sigma-1}} dt \\ &+ \left. \frac{1}{\Gamma(\sigma)} \int_0^s \frac{F(t, y(\beta(t)), y(\theta(t)), (Hy)(t))}{(s-t)^{\sigma-1}} dt \right| \\ &\leq \frac{9}{8} + \frac{1}{8} r_0 + \frac{1}{\Gamma(\sigma+1)} \left(\frac{1}{10} + \frac{r_0}{7} \right) \\ &+ \frac{1}{\Gamma(\sigma+1)} \left[\frac{3r_0}{5} + \frac{1}{6} \left(\frac{r_0}{2} + 1 \right) + \frac{\lambda}{10} \right], \quad \forall s \in I_a. \end{aligned}$$

Therefore (H3) holds if according to the Eq. (8),

$$\begin{aligned}
L + A + \frac{M_1 a^\sigma}{\Gamma(1+\sigma)} + \frac{M_2 a^\sigma}{\Gamma(1+\sigma)} \\
\leq \frac{9}{8} + \frac{1}{8} r_0 + \frac{1}{\Gamma(\sigma+1)} \left(\frac{1}{10} + \frac{r_0}{7} \right) + \frac{1}{\Gamma(\sigma+1)} \left[\frac{3r_0}{5} \right. \\
\left. + \frac{1}{6} \left(\frac{r_0}{2} + 1 \right) + \frac{\lambda}{10} \right] \simeq \left\{ \begin{array}{l} 1.790, \quad \sigma = \frac{5}{4}, \\ 1.701, \quad \sigma = \frac{3}{2}, \\ 1.613, \quad \sigma = \frac{7}{4}, \end{array} \right\} \leq r_0. \quad (13)
\end{aligned}$$

This shows that

$$r_0 \simeq \left\{ \begin{array}{l} 1.215 + 0.088\lambda, \quad \sigma = \frac{5}{4}, \\ 1.072 + 0.075\lambda, \quad \sigma = \frac{3}{2}, \\ 0.929 + 0.062\lambda, \quad \sigma = \frac{7}{4}, \end{array} \right. \quad (14)$$

is a solution of the above inequality. Table 1 shows the numerical results of Eqs.(13) and (14). Furthermore, Figs. 1a and 1b show M_1 and M_2 , respectively. One can see the 2D plot of suitable r_0 in Fig. 1c, well.

The result is followed from Theorem 3.1. Therefor, assumptions (H1)-(H3) be fulfilled and Theorem 3.1 indicates the solution of (12) in $C(I_a)$.

Example 4.2. Consider following FSIIDE,

$$\begin{aligned}
{}^C D^\sigma \left(y(s) + \frac{2+\ln(1+|y(s)|)}{(2s+3)^2} \right) &= \frac{1}{5} e^{-s} + \frac{3 \cos(s)y(s^3)}{4+3s} \\
&+ \frac{1}{9} \sin \left(\sqrt{\frac{\pi}{2}} y(\sqrt{s}) \right) + \frac{s^2}{2(1+s^2)} \int_0^s \sqrt{\frac{se^{-3t}}{1+s}} \left[\frac{1}{5} \right. \\
&+ \left. \int_0^t \xi \left(\frac{|y(\xi)|}{1+|y(\xi)|} + y(\xi) \right) d\xi \right] dt \\
&+ \frac{e^{-s}}{3+s^2} \int_0^s \frac{st \cos(sy(\sqrt{t}))}{2+t^2+5s} dW(t), \quad (15)
\end{aligned}$$

for $y \in C(I_a)$, $a = 1$ and for $\sigma = \frac{1}{5}, \frac{1}{3}, \frac{1}{2}$, via condition $y(0) = y_0 = 0$.

Table 1: Numerical results of A , M_1 , M_2 and suitable r_0 of FSIDE (12) for three different values of σ in Example 4.1.

s	A	M_1	M_2	$r_0 \geq \dots$	r_0
$\sigma = \frac{5}{4}$					
0.00	0.125	0.000	0.000	1.250	$1.215 + 0.088\lambda$
0.10	0.124	0.005	0.017	1.271	$1.215 + 0.088\lambda$
0.20	0.122	0.011	0.040	1.299	$1.215 + 0.088\lambda$
0.30	0.119	0.019	0.068	1.332	$1.215 + 0.088\lambda$
0.40	0.115	0.029	0.103	1.372	$1.215 + 0.088\lambda$
0.50	0.109	0.041	0.145	1.420	$1.215 + 0.088\lambda$
0.60	0.103	0.054	0.196	1.478	$1.215 + 0.088\lambda$
0.70	0.095	0.069	0.254	1.544	$1.215 + 0.088\lambda$
0.80	0.087	0.087	0.320	1.618	$1.215 + 0.088\lambda$
0.90	0.077	0.106	0.392	1.700	$1.215 + 0.088\lambda$
1.00	0.067	0.127	0.471	1.790	$1.215 + 0.088\lambda$
$\sigma = \frac{3}{2}$					
0.00	0.125	0.000	0.000	1.250	$1.072 + 0.075\lambda$
0.10	0.124	0.002	0.008	1.260	$1.072 + 0.075\lambda$
0.20	0.122	0.007	0.023	1.277	$1.072 + 0.075\lambda$
0.30	0.119	0.012	0.043	1.300	$1.072 + 0.075\lambda$
0.40	0.115	0.020	0.070	1.329	$1.072 + 0.075\lambda$
0.50	0.109	0.029	0.104	1.368	$1.072 + 0.075\lambda$
0.60	0.103	0.040	0.147	1.415	$1.072 + 0.075\lambda$
0.70	0.095	0.054	0.198	1.472	$1.072 + 0.075\lambda$
0.80	0.087	0.070	0.258	1.539	$1.072 + 0.075\lambda$
0.90	0.077	0.088	0.326	1.616	$1.072 + 0.075\lambda$
1.00	0.067	0.108	0.401	1.701	$1.072 + 0.075\lambda$
$\sigma = \frac{7}{4}$					
0.00	0.125	0.000	0.000	1.250	$0.929 + 0.062\lambda$
0.10	0.124	0.001	0.004	1.254	$0.929 + 0.062\lambda$
0.20	0.122	0.004	0.013	1.264	$0.929 + 0.062\lambda$
0.30	0.119	0.007	0.026	1.278	$0.929 + 0.062\lambda$
0.40	0.115	0.013	0.046	1.299	$0.929 + 0.062\lambda$
0.50	0.109	0.020	0.072	1.327	$0.929 + 0.062\lambda$
0.60	0.103	0.029	0.107	1.364	$0.929 + 0.062\lambda$
0.70	0.095	0.041	0.150	1.411	$0.929 + 0.062\lambda$
0.80	0.087	0.055	0.202	1.468	$0.929 + 0.062\lambda$
0.90	0.077	0.071	0.262	1.535	$0.929 + 0.062\lambda$
1.00	0.067	0.089	0.332	1.613	$0.929 + 0.062\lambda$

Here $n = 1$,

$$g(s, y(s)) = \frac{2 + \ln(1 + |y(s)|)}{(2s+3)^2},$$




Figure 1: Graphical representation of M_1 , M_2 and suitable r_0 of of FSIDE (12) for three different values of σ in Example 4.1.

$$f(s, y(\alpha(s))) = \frac{1}{5}e^{-s} + \frac{3 \cos(s)y(s^3)}{4+3s},$$

$$F(s, u, v, w) = \frac{1}{9} \sin\left(\sqrt{\frac{\pi}{2}}u\right) + \frac{s^2}{2(1+s^2)}v + \frac{e^{-s}}{3+s^2}w,$$

and

$$v = \int_0^s \sqrt{\frac{se^{-3t}}{1+s}} \left(\frac{1}{5} + \int_0^t \xi \left[\frac{|y(\xi)|}{1+|y(\xi)|} + y(\xi) \right] d\xi \right) dt,$$

$$w = \int_0^s \frac{st \cos(sy(\sqrt{t}))}{2+t^2+5s} dW(t).$$

It can be seen that we have $|g(s, u) - g(s, \bar{u})| \leq \frac{1}{9}|u - \bar{u}|$ and

$$|F(s, u, v, w) - F(s, \bar{u}, \bar{v}, \bar{w})| \leq \frac{\sqrt{\pi}}{18}|u - \bar{u}| + \frac{1}{2}|v - \bar{v}| + \frac{1}{3}|w - \bar{w}|.$$

So, we can choose $k = \frac{1}{9} < 1$, $c_1 = \frac{\sqrt{\pi}}{18}$, $c_2 = \frac{1}{2}$, $c_3 = \frac{1}{3}$. It follows that the conditions (H1) and (H2) hold. Moreover, for $\|y\| \leq r_0$, $r_0 > 0$, we have

$$\begin{aligned} |y(s)| &= \left| y(0) + g(0, y_0) - g(s, y(s)) + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{f(t, y(\alpha(t)))}{(s-t)^{1-\sigma}} dt \right. \\ &\quad \left. + \frac{1}{\Gamma(\sigma)} \int_0^s \frac{F(t, y(\beta(t)), y(\theta(t)), (Hy)(t))}{(s-t)^{1-\sigma}} dt \right| \\ &\leq \frac{2}{9} + \frac{1}{\Gamma(\sigma+1)} \left(\frac{1}{5} + \frac{r_0}{4} \right) \\ &\quad + \frac{1}{\Gamma(\sigma+1)} \left[\frac{1}{9} + \frac{1}{4} \left(1 + r_0 + \frac{1}{5} \right) + \frac{1}{6} \lambda \right], \quad \forall s \in I_a. \end{aligned}$$

Therefore (H3) holds if according to the Eq. (8),

$$\begin{aligned} L + A + \frac{M_1 a^\sigma}{\Gamma(1+\sigma)} + \frac{M_2 a^\sigma}{\Gamma(1+\sigma)} \\ \leq \frac{2}{9} + \frac{1}{\Gamma(\sigma+1)} \left(\frac{1}{5} + \frac{r_0}{4} \right) + \frac{1}{\Gamma(\sigma+1)} \left[\frac{1}{9} + \frac{1}{4} \left(1 + r_0 + \frac{1}{5} \right) + \frac{1}{6} \lambda \right] \\ \simeq \begin{cases} 1.128, & \sigma = \frac{1}{5}, \\ 1.151, & \sigma = \frac{1}{3}, \\ 1.158, & \sigma = \frac{1}{2}, \end{cases} \leq r_0, \end{aligned} \quad (16)$$

$|y(s)| \leq r_0$. This shows that

$$r_0 = \frac{2\sqrt{\pi} + 11 + 3\lambda}{9(\sqrt{\pi} - 1)} \simeq \begin{cases} 0.432 + 0.182\lambda, & \sigma = \frac{1}{5}, \\ 0.466 + 0.187\lambda, & \sigma = \frac{1}{3}, \\ 0.476 + 0.188\lambda, & \sigma = \frac{1}{2}. \end{cases} \quad (17)$$

Table 1 shows the numerical results of Eqs.(16) and (17). Furthermore, Figs. 2a and 2b show M_1 and M_2 , respectively. One can see the 2D plot of suitable r_0 in Fig. 2c, well. The result is followed from Theorem 3.1.

Table 2: Numerical results of A , M_1 , M_2 and suitable r_0 of FSIIDE (15) for three different values of σ in Example 4.2.

s	A	M_1	M_2	$r_0 \geq \dots$	r_0
$\sigma = \frac{1}{5}$					
0.00	0.222	0.000	0.000	0.444	$0.432 + 0.182\lambda$
0.10	0.195	0.601	0.286	1.305	$0.432 + 0.182\lambda$
0.20	0.173	0.634	0.315	1.345	$0.432 + 0.182\lambda$
0.30	0.154	0.628	0.336	1.340	$0.432 + 0.182\lambda$
0.40	0.139	0.603	0.356	1.320	$0.432 + 0.182\lambda$
0.50	0.125	0.569	0.377	1.293	$0.432 + 0.182\lambda$
0.60	0.113	0.528	0.400	1.263	$0.432 + 0.182\lambda$
0.70	0.103	0.482	0.424	1.231	$0.432 + 0.182\lambda$
0.80	0.095	0.434	0.448	1.198	$0.432 + 0.182\lambda$
0.90	0.087	0.384	0.471	1.163	$0.432 + 0.182\lambda$
1.00	0.080	0.332	0.493	1.128	$0.432 + 0.182\lambda$
$\sigma = \frac{1}{3}$					
0.00	0.222	0.000	0.000	0.444	$0.466 + 0.187\lambda$
0.10	0.195	0.455	0.217	1.089	$0.466 + 0.187\lambda$
0.20	0.173	0.526	0.262	1.183	$0.466 + 0.187\lambda$
0.30	0.154	0.550	0.294	1.220	$0.466 + 0.187\lambda$
0.40	0.139	0.549	0.324	1.233	$0.466 + 0.187\lambda$
0.50	0.125	0.533	0.354	1.234	$0.466 + 0.187\lambda$
0.60	0.113	0.507	0.384	1.227	$0.466 + 0.187\lambda$
0.70	0.103	0.473	0.415	1.214	$0.466 + 0.187\lambda$
0.80	0.095	0.433	0.447	1.196	$0.466 + 0.187\lambda$
0.90	0.087	0.389	0.477	1.175	$0.466 + 0.187\lambda$
1.00	0.080	0.342	0.507	1.151	$0.466 + 0.187\lambda$
$\sigma = \frac{1}{2}$					
0.00	0.222	0.000	0.000	0.444	$0.476 + 0.188\lambda$
0.10	0.195	0.312	0.149	0.878	$0.476 + 0.188\lambda$
0.20	0.173	0.405	0.202	1.002	$0.476 + 0.188\lambda$
0.30	0.154	0.453	0.242	1.072	$0.476 + 0.188\lambda$
0.40	0.139	0.475	0.280	1.116	$0.476 + 0.188\lambda$
0.50	0.125	0.479	0.317	1.143	$0.476 + 0.188\lambda$
0.60	0.113	0.469	0.356	1.160	$0.476 + 0.188\lambda$
0.70	0.103	0.449	0.394	1.169	$0.476 + 0.188\lambda$
0.80	0.095	0.420	0.434	1.171	$0.476 + 0.188\lambda$
0.90	0.087	0.385	0.473	1.167	$0.476 + 0.188\lambda$
1.00	0.080	0.344	0.511	1.158	$0.476 + 0.188\lambda$




Figure 2: Graphical representation of M_1 , M_2 and suitable r_0 of of FSIDE (15) for three different values of σ in Example 4.2.

5 Conclusion and Perspective

In this work, Theorem 2.12 and the M.N.C idea were used to analyze the of solutions some nonlinear functional FSIDE in the Banach algebra $C(I_a)$. The superiority of Theorem 2.12 compared to other similar FPTs, such as Darbo and Schauder, is that here the condition that involved operator maps a closed convex subset onto itself is not needed. Thus by applying weaker conditions, the method is extended and includes a

larger range. In fact, Some valuable articles such as [1, 11, 20, 43], can be generalized or be used with the results of this research.

References

- [1] N. Adjimi, A. Boutiara, M. E. Samei, S. Etemad and S. Rezapour, On solutions of a hybrid generalized Caputo-type problem via the measure of non-compactness in the generalized version of Darbo's theorem, *Journal of Inequalities and Applications*, 2023 (2023), 34.
- [2] R. P. Agarwal, N. Hussain and M. A. Taoudi, Fixed point theorems in ordered Banach spaces and applications to nonlinear integral equations, *Abstract and Applied Analysis*, Article ID 245872 (2012), 15 pages.
- [3] A. Aghajani, J. Banaś, and Y. Jalilian, Existence of solutions for a class of nonlinear Volterra singular integral equations, *Computers & Mathematics with Applications*, 62(3) (2011), 1215–1227.
- [4] S. N. Al-Rawi, On the Solution of Certain Fractional Integral Equations, *Kirkuk Journal of Science*, 1(2) (2006), 125–136.
- [5] M. Ammi, E. Kinani and D. F. M. Torres, Existence and uniqueness of solution to a functional integro-differential fractional equation, *Electronic Journal of Differential Equations*, 2012(103) (2012), 1–9.
- [6] K. E. Atkinson, *The Numerical Solution of Integral Equations of the Second Kind*, Cambridge University Press, New York (1997).
- [7] S. M. Aydogan, On a K -dimensional system of hybrid fractional differential equations with multi-point boundary conditions, *Journal of Mathematical Extension*, 15(10) SI-NTFCA (2021), 1–18.
- [8] S. M. Aydogan, A. Hussain and F. M. Saker, On a nonlinear fractional-Order model of COVID-19 under AB-fractional derivative, *Journal of Mathematical Extension*, 15(11) SI-NTFCA (2021), 1–31.

- [9] K. Balachandran, S. Kiruthika and J. J. Trujillo, Existence of solutions of nonlinear fractional pantograph equations, *Acta Mathematica Scientia*, 33(3) 2013, 712–720.
- [10] J. Banaś, Measures of noncompactness in the study of solutions of nonlinear differential and integral equations, *Central European Journal of Mathematics*, 10(6) (2012), 2003–2011.
- [11] Bhupeshwar, D. K. Patel, M. Gabeleh and M. E. Samei, On existence of solutions for Ψ -Hilfer type fractional BVP of generalized higher order, *Computational and Applied Mathematics*, 43 (2024), 155.
- [12] S. Carl and S. Heikkilä, *Fixed Point Theory in Ordered Sets and Applications: From Differential and Integral Equations to Game Theory*, Springer Science & Business Media, India (2010).
- [13] D. Colton, and R. Kress, *Integral Equation Methods in Scattering Theory*, SIAM, Society for Industrial and Applied Mathematics, Philadelphia, USA (2013).
- [14] S. Dadsetadi, K. Nouri and L. Torkzadeh, Solvability of some nonlinear integro-differential equation of fractional order via measure of noncompactness, *The Pure and Applied Mathematics, Korean Society of Mathematical Education*, 27(1) (2020), 13–24.
- [15] M. A. Darwish and S. K. Ntouyas, On a quadratic fractional Hammerstein-Volterra integral equation with linear modification of the argument, *Nonlinear Analysis: Theory, Methods & Applications*, 74(11) (2011), 3510–3517.
- [16] A. Deep, S. Abbas, B., Singh, M. R. Alharthi and K. S. Nisar, Solvability of functional stochastic integral equations via Darbo's fixed point theorem. *Alexandria Engineering Journal*, 60(6) (2021), 5631–5636.
- [17] L. M. Delves and J. L. Mohamed, *Computational Methods for Integral Equations*, Cambridge University Press, UK (1988).

- [18] A. M. A. El-Sayed, On the stochastic fractional calculus operators, *Journal of Fractional Calculus and Applications*, 6(1) (2015), 101–109.
- [19] L. Eskola, *Geophysical Interpretation using Integral Equations*, Springer Science & Business Media, Netherlands (2012).
- [20] M. Fatehi, S. Rezapour, M. E. Samei, Investigation of the solution for the k -dimensional device of differential inclusion of Laplacein fraction with sequential derivatives and boundary conditions of integral and derivative, *Journal of Mathematical Extension*, 17(11) (2023), (6)1–33.
- [21] L. S. Gol'denšteĭn and A. S. Markus, On the measure of non-compactness of bounded sets and of linear operators, *Studies in Algebra and Math. Anal. (Russian)*, Izdat. “Karta Moldovenjaske”, Kishinev (1965), 45–54.
- [22] S. Gülyaz and I. M. Erhan, Existence of solutions of integral equations via fixed point theorems, *Journal of Inequalities and Applications*, 2014(1) (2014), 1–15.
- [23] F. M. Hafiz, The fractional calculus for some stochastic processes, *Stochastic Analysis and Applications*, 22(2) (2004), 507–523.
- [24] M. Kamrani, Numerical solution of stochastic fractional differential equations, *Numerical Algorithms*, 68 (2015), 81–93.
- [25] E. Karapinar, A. Fulga, N. Shahzad and A. F. Roldán López de Hierro, Solving integral equations by means of fixed point theory, *Journal of Function Spaces*, 2022 (2022).
- [26] M. Kazemi and R. Ezzati, Existence of solutions for some nonlinear Volterra integral equations via Petryshyn fixed point theorem, *International Journal of Nonlinear Analysis and Applications*, 9(1) (2018), 1–12.
- [27] M. Kazemi and A. R. Yaghoobnia, Application of fixed point theorem to solvability of functional stochastic integral equations, *Applied Mathematics and Computation*, 417 (2022), 126759.

- [28] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, 204, New York (2006).
- [29] F. C. Klebaner, *Introduction to Stochastic Calculus with Applications*, World Scientific Publishing Company, New York (2012).
- [30] K. Kuratowski, Sur les espaces completes, *Fundamenta Mathematicae*, 15 (1934), 301–335.
- [31] N. Laksaci, A. Boudaoui, K. Abodayeh, W. Shatanawi and T. A. M. Taqi, Existence and uniqueness results of coupled fractional-order differential systems involving Riemann-Liouville derivative in the space $W_{a^+}^{\gamma_1,1}(a,b) \times W_{a^+}^{\gamma_2,1}(a,b)$ with Perov's Fixed Point Theorem, *Fractal and Fractional*, 5(4) (2021), 217.
- [32] Ü. Lepik, Solving fractional integral equations by the Haar wavelet method, *Applied Mathematics and Computation*, 214(2) (2009), 468–478.
- [33] K. Maleknejad, K. Nouri and R. Mollapourasl, Investigation on the existence of solutions for some nonlinear functional-integral equations, *Nonlinear Anal.*, 71(12) (2009), e1575–e1578.
- [34] N. Mandal and A. Chakrabarti, Solution of a system of generalized Abel integral equations using fractional calculus, *Applied Mathematics Letters*, 9(5) (1996), 1–4.
- [35] R. D. Nussbaum, *The fixed-point index and fixed point theorem for k -set contractions*, ProQuest LLC, Ann Arbor, MI, Thesis (Ph.D.)—The University of Chicago (1969).
- [36] B. Øksendal, *Stochastic Differential Equations*, Springer, pp:65–84, London (2003).
- [37] W. V. Petryshyn, Structure of the fixed points sets of k -set-contractions, *Archive for Rational Mechanics and Analysis*, 40 (1970/1971), 312–328.

- [38] P. E. Protter, On the existence, uniqueness, convergence and explosions of solutions of systems of stochastic integral equations, *The Annals of Probability*, 5(2) (1977), 243–261.
- [39] J. Prüss, *Evolutionary Integral Equations and Applications*, Birkhäuser Basel, Switzerland (2013).
- [40] M. Rahman, *Integral Equations and their Applications*, WIT Press publishes, Ashurst Lodge, Ashurst, Southampton, SO40 7AA, UK (2007).
- [41] A. P. Raiche, An integral equation approach to three-dimensional modeling, *Geophysical Journal International*, 36(2) (1974), 363–376.
- [42] R. A. Rashwan and D.M. Albaqeri, A common random fixed point theorem and application to random integral equations, *International Journal of Applied Mathematics Research*, 3(1) (2014), pages 71.
- [43] M. E. Samei, Employing Kuratowski measure of non-compactness for positive solutions of system of singular fractional q -differential equations with numerical effects, *Filomat*, 34(9) (2020), 1–19.
- [44] M. E. Samei, H. Zanganeh and S. M. Aydogan, Investigation of a class of the singular fractional integro-differential quantum equations with multi-step methods, *Journal of Mathematical Extension*, 15(15) SI-NTFCA (2021), 1–54.
- [45] S. P. Singh, B. Watson and P. Srivastava, “*Fixed point theory and best approximation: The KKM-map principle*”, Springer Science & Business Media, 424 (2013).
- [46] C. P. Tsokos and W. J. Padgett, *Random Integral Equations with Applications to Stochastic Systems*, Springer (2006).
- [47] A. M. Wazwaz, *Linear and Nonlinear Integral Equations: Methods and Applications*, Higher Education Press, Springer-Verlag Berlin Heidelberg (2011).

- [48] W. Wei, X. Li, and X. Li, New stability results for fractional integral equation, *Computers & Mathematics with Applications*, 67(10) (2012) 3468–3475.
- [49] N. Xu, and W. Liu, Iterative solutions for a coupled system of fractional differential–integral equations with two-point boundary conditions, *Applied Mathematics and Computation*, 244 (2014), 903–911.
- [50] M. A. Zaky, An accurate spectral collocation method for nonlinear systems of fractional differential equations and related integral equations with nonsmooth solutions, *Applied Numerical Mathematics*, 154 (2020), 205–222.

Alireza Yaghoobnia

Assistant Professor of Mathematics
Department of Mathematics
Lahijan Branch, Islamic Azad University
Lahijan, Iran
E-mail: yaghoobnia@liau.ac.ir

Manochehr Kazemi

Associate Professor of Mathematics
Department of Mathematics
Ashtian Branch, Islamic Azad University
Ashtian, Iran
E-mail: univer_ka@yahoo.com

Vishnu Narayan Mishra

Professor of Mathematics
Department of Mathematics, Faculty of Science
Indira Gandhi National Tribal University
Lalpur, Amarkantak 484 887, Madhya Pradesh, India
E-mail: vishnunarayanmishra@gmail.com, vnm@igntu.ac.in