

Journal of Mathematical Extension
Journal Pre-proof
ISSN: 1735-8299
URL: <http://www.ijmex.com>
Original Research Paper

Essential Norm of Generalized Integral Type Operators From Mixed-Norm to Zygmund-Type Spaces

Y. Estaremi*

Golestan University

M. Hassanlou

Urmia University of Technology

M. S. AL Ghafri

University of Technology and Applied Sciences

Abstract. In this paper, we consider the generalized integration operator from mixed-norm space into Zygmund-type and Bloch-type spaces and find an estimation for the essential norm of this operator.

AMS Subject Classification: 47B33; 47B38; 30H05.

Keywords and Phrases: Essential norm, generalized integration operator, mixed-norm space Zygmund-type space, weighted Bloch space.

1 Introduction and Preliminaries

We denote by \mathbb{D} the open unit disc in the complex plane \mathbb{C} and $H(\mathbb{D})$ the space of analytic functions on \mathbb{D} . Every analytic function (self-map) $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ induces a linear operator via composition called composition operator C_φ , $C_\varphi f = f \circ \varphi$. A complete history of these operators on the spaces of analytic functions can be found in [3]. Throughout the recent

Received: July 2024; Accepted: December 2024

*Corresponding Author

decades there has been a significant interest on investigating the operator properties of composition operator and generalization of it while they act on some spaces of analytic or measurable functions. The above operator can be generalized by

$$(C_{\varphi}^g f)(z) = \int_0^z f'(\varphi(\xi))g(\xi) d\xi, \quad f \in H(\mathbb{D}), z \in \mathbb{D},$$

where $g \in H(\mathbb{D})$, see [5]. Also, if we replace n -th derivative with 1-th derivative, then we have

$$(C_{\varphi,g}^n f)(z) = \int_0^z f^{(n)}(\varphi(\xi))g(\xi) d\xi, \quad f \in H(\mathbb{D}), z \in \mathbb{D},$$

where $n \in \mathbb{N}_0$. If $n = 0$, then $C_{\varphi,g}^0$ is the Volterra composition operator. Also for $\varphi(z) = z$ and g' we have Volterra type operator T_g ,

$$T_g f(z) = \int_0^z f(w)g'(w)dw.$$

$C_{\varphi,g}^n$ is called generalized integration operator, see for example [4, 12] and references therein. A natural question dealing with this kind of operators or other classes of operators is that, under which conditions they are bounded, compact, have closed range and \dots . Another aspect is computing or estimating norm, essential norm, numerical range, \dots of the operator. All characterizations are in terms of symbols induce the operators.

The aim of this paper is to estimate the essential norm of generalized integration operators from mixed-norm spaces, which include some well-known spaces of analytic functions, into weighted Zygmund space. As a result, compactness criteria for the operator, has been proved in [4], is obtained.

A positive continuous function ϕ on $[0, 1)$ is called normal if there are two constants $b > a > 0$ such that

- (i) $\frac{\phi(r)}{(1-r)^a}$ is non-increasing in $[0, 1)$ and $\frac{\phi(r)}{(1-r)^a} \downarrow 0$
- (ii) $\frac{\phi(r)}{(1-r)^b}$ is non-decreasing in $[0, 1)$ and $\frac{\phi(r)}{(1-r)^b} \uparrow \infty$

as $r \rightarrow 1^-$. For $0 < p, q < \infty$ and ϕ normal, the mixed-norm space $H(p, q, \phi)$ consists of all analytic functions in \mathbb{D} for which

$$\|f\|_{p,q,\phi}^p = \int_0^1 M_q^p(f, r) \frac{\phi^p(r)}{1-r} dr < \infty$$

where

$$M_q(f, r) = \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^q d\theta \right)^{1/q}.$$

If $1 \leq p < \infty$, then $H(p, q, \phi)$ becomes a Banach space equipped with the norm $\|\cdot\|_{p,q,\phi}$. In the case $0 < p < 1$, $\|\cdot\|_{p,q,\phi}$ is a quasi-norm on $H(p, q, \phi)$ and this space is a Fréchet space but is not a Banach space. For $\alpha > -1$, $\phi(r) = (1-r)^{(\alpha+1)/p}$ is a normal function. If $p = q$, then $H(p, p, (1-r)^{(\alpha+1)/p})$ is weighted Bergman space A_α^p which is defined by

$$A_\alpha^p = \{f \in H(\mathbb{D}) : \|f\|_{A_\alpha^p}^p = \int_{\mathbb{D}} |f(z)|^p (1-|z|^2)^\alpha dA(z) < \infty\},$$

where dA is the normalized area measure on \mathbb{D} .

A positive continuous function μ on \mathbb{D} is called weight. For a weight μ , the weighted Zygmund space \mathcal{Z}_μ , Zygmund-type space, is the space of $f \in H(\mathbb{D})$ such that

$$\sup_{z \in \mathbb{D}} \mu(|z|) |f''(z)| < \infty.$$

The weighted Zygmund space \mathcal{Z}_μ is a Banach space with respect to the norm:

$$\|f\|_{\mathcal{Z}_\mu} = |f(0)| + |f'(0)| + \sup_{z \in \mathbb{D}} \mu(|z|) |f''(z)|.$$

The weighted Bloch space \mathcal{B}_μ defined by

$$\mathcal{B}_\mu = \{f \in H(\mathbb{D}) : \|f\|_{\mathcal{B}_\mu} = |f(0)| + \sup_{z \in \mathbb{D}} \mu(|z|) |f'(z)| < \infty\}.$$

Note that if $\mu(z) = (1-z^2)$, then we obtain classical Zygmund space \mathcal{Z} and Bloch space \mathcal{B} . For a complete information about classical Bloch space, see [13].

Hu in [6] has characterized those holomorphic symbols on the unit ball for which the induced extended cesàro operator is bounded (compact) on mixed norm space. The boundedness of cesàro operators on mixed norm space is studied in [7]. Boundedness and compactness of the generalized integration operator from a mixed norm space into a weighted Zygmund space investigated by Guo and Liu in [4]. More results on integral type operators acting on mixed norm, Zygmund and Bloch spaces can be found in [1, 5, 8, 9, 10, 11, 12].

Guo and Liu in [4] have characterized compact generalized integration operators $C_{\varphi,g}^n$ between mixed norm and weighted Zygmund spaces. Indeed, they have proved that $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is compact if and only if

$$A = \lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n+1}} = 0$$

$$B = \lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|g'(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n}} = 0$$

In this paper, motivated by the above results, we find an approximation of the essential norm of the generalized integration operators $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ and prove that

$$\|C_{\varphi,g}^n\|_e \approx \max\{A, B\},$$

it means that there exists constants C, D such that $CB \leq A \leq DB$ or equivalently $CA \leq B \leq DA$.

Recall that for any operator T between two Banach spaces X and Y , the essential norm of T is denoted by $\|T\|_{e, X \rightarrow Y}$ and is defined as follows

$$\|T\|_{e, X \rightarrow Y} = \inf\{\|T - S\| : S \text{ is a compact operator from } X \text{ to } Y\}.$$

The operator is compact if and only if $\|T\|_{e, X \rightarrow Y} = 0$.

2 Essential Norm of $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ or \mathcal{B}_μ

In this section we give an estimation for the essential norm of the operator $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ or \mathcal{B}_μ . First we recall a fundamental lemma for later use in the paper.

Lemma 2.1. [10] *Assume that $p, q \in (0, \infty)$, ϕ is normal and $f \in H(p, q, \phi)$. Then for each $n \in \mathbb{N}_0$, there is a positive constant C independent of f such that*

$$|f^{(n)}(z)| \leq C \frac{\|f\|_{p,q,\phi}}{\phi(|z|)(1-|z|^2)^{1/q+n}}, \quad z \in \mathbb{D}.$$

The following lemma can be obtained by a standard argument, see [2].

Lemma 2.2. *Let φ be a analytic self-map of \mathbb{D} . Then $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is compact if and only if $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded and for any bounded sequence $\{f_k\}$ in $H(p, q, \phi)$ which converges uniformly to zero on compact subsets of \mathbb{D} as $k \rightarrow \infty$, we have $\|C_{\varphi,g}^n f_k\|_{\mathcal{Z}_\mu} \rightarrow 0$ as $k \rightarrow \infty$.*

By the above observations, in the next proposition, we get that every bounded generalized integration operator $C_{\varphi,g}^n$ with $\|\varphi\|_\infty = \sup_{z \in \mathbb{D}} |\varphi(z)| < 1$, is compact.

Proposition 2.3. *Let $p, q \in (0, \infty)$, ϕ is normal, $g \in H(\mathbb{D})$ and φ be a analytic self-map of \mathbb{D} . If $\|\varphi\|_\infty < 1$ and the operator $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded then it is compact*

Proof. Since $C_{\varphi,g}^n$ is bounded, then there exists a positive constant C such that for

$$\|C_{\varphi,g}^n f\|_{\mathcal{Z}_\mu} \leq C \|f\|_{p,q,\phi}, \quad (1)$$

for all $f \in H(p, q, \phi)$. Let $f_1(z) = \frac{z^n}{n!}$. Then we have

$$\sup_{z \in \mathbb{D}} \mu(|z|) |g'(z)| = \|C_{\varphi,g}^n f_1\|_{\mathcal{Z}_\mu} \leq C \|f_1\|_{p,q,\phi} < \infty. \quad (2)$$

Similarly for $f_2(z) = \frac{z^{n+1}}{(n+1)!}$, by using triangle inequality and the fact that $|\varphi(z)| < 1$, we get that

$$\sup_{z \in \mathbb{D}} \mu(|z|) |\varphi'(z)g(z)| < \infty. \quad (3)$$

Now assume that $\{f_k\}_{k \in \mathbb{N}}$ is a bounded sequence in $H(p, q, \phi)$ such that converges to 0, uniformly on compact subsets of \mathbb{D} , as $k \rightarrow \infty$. This

implies that $\{f'_k\}_{k=1}^\infty$ and $\{f''_k\}_{k=1}^\infty$ converge uniformly to 0, on compact subsets of \mathbb{D} , as $k \rightarrow \infty$. By (2) and (3) we obtain

$$\begin{aligned}
\|C_{\varphi,g}^n f_k\|_{\mathcal{Z}_\mu} &= |(C_{\varphi,g}^n f_k)'(0)| + \sup_{z \in \mathbb{D}} \mu(z) |(C_{\varphi,g}^n f_k)''(z)| \\
&\leq |f_k^{(n)}(\varphi(0))g(0)| + \sup_{z \in \mathbb{D}} \mu(|z|) |(C_{\varphi,g}^n f_k)''(z)| \\
&\leq |f_k^{(n)}(\varphi(0))g(0)| + \sup_{z \in \mathbb{D}} \mu(|z|) |f_k^{(n+1)}(\varphi(z))\varphi'(z)g(z)| \\
&\quad + \sup_{z \in \mathbb{D}} \mu(|z|) |f_k^{(n)}(\varphi(z))g'(z)| \\
&= |f_k^{(n)}(\varphi(0))g(0)| + \sup_{|\varphi(z)| \leq \rho} \mu(|z|) |f_k^{(n+1)}(\varphi(z))\varphi'(z)g(z)| \\
&\quad + \sup_{|\varphi(z)| \leq \rho} \mu(|z|) |f_k^{(n)}(\varphi(z))g'(z)|,
\end{aligned}$$

which tends to 0, as $k \rightarrow \infty$. Here $\rho = \|\varphi\|_\infty$, $\rho \in (0, 1)$. Hence by Lemma 2.2, $C_{\varphi,g}^n$ is compact. \square

Now, in the next theorem we estimate the essential norm of $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$. We use $\|C_{\varphi,g}^n\|_e$ for the essential norm of $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ instead of $\|C_{\varphi,g}^n\|_{e, H(p, q, \phi) \rightarrow \mathcal{Z}_\mu}$.

Theorem 2.4. *Let $p, q \in (0, \infty)$, ϕ is normal and φ be a analytic self-map of \mathbb{D} . If the operator $C_{\varphi,g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded then*

$$\|C_{\varphi,g}^n\|_e \approx \max \left\{ \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |\varphi'(z)g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n+1}}, \right. \\
\left. \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |g'(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n}} \right\}.$$

Proof. If $\|\varphi\|_\infty < 1$, then by the Proposition 2.3, the operator $C_{\varphi,g}^n$ is compact and therefore $\|C_{\varphi,g}^n\|_e = 0$. Also by Theorem 2.7 of [4], we have

$$\limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |\varphi'(z)g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n+1}} = 0$$

and

$$\limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |g'(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n}} = 0.$$

Now assume that $\|\varphi\|_\infty = 1$. Let $\{z_j\}_{j \in \mathbb{N}}$ be a sequence in \mathbb{D} such that $|\varphi(z_j)| \rightarrow 1$, as $j \rightarrow \infty$. Since ϕ is a normal function, then we have a positive constant b from the definition. Put $\alpha = 1/q + b + 1$ and define the sequence $\{f_k\}$ as follows

$$f_k(z) = \frac{(1 - |\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \left(\frac{1}{(1 - \overline{\varphi(z_k)}z)^\alpha} - \frac{\alpha(1 - |\varphi(z_k)|^2)}{(\alpha + n)(1 - \overline{\varphi(z_k)}z)^{\alpha+1}} \right).$$

Direct calculations shows that

$$\begin{aligned} f_k^{(n)}(z) &= \frac{(1 - |\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \frac{\alpha(\alpha + 1) \cdots (\alpha + n - 1) \overline{\varphi(z_k)}^n}{(1 - \overline{\varphi(z_k)}z)^{\alpha+n}} \\ &\quad - \frac{(1 - |\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \\ &\quad \times \frac{\alpha(\alpha + 1) \cdots (\alpha + n - 1)(\alpha + n)(1 - |\varphi(z_n)|^2) \overline{\varphi(z_n)}^n}{(\alpha + n)(1 - \overline{\varphi(z_k)}z)^{\alpha+n+1}} \\ f_k^{(n+1)}(z) &= \frac{(1 - |\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \frac{\alpha(\alpha + 1) \cdots (\alpha + n) \overline{\varphi(z_k)}^{n+1}}{(1 - \overline{\varphi(z_k)}z)^{\alpha+n}} \\ &\quad - \frac{(1 - |\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \\ &\quad \times \frac{\alpha(\alpha + 1) \cdots (\alpha + n - 1)(\alpha + n)(1 - |\varphi(z_k)|^2) \overline{\varphi(z_k)}^n}{(\alpha + n)(1 - \overline{\varphi(z_k)}z)^{\alpha+n+1}}. \end{aligned}$$

It is not difficult to show that $\{f_k\}$ is a bounded sequence in $H(p, q, \phi)$, i.e., there exists a positive constant C such that

$$\sup_k \|f_k\|_{p,q,\phi} \leq C. \quad (4)$$

Also, it is convergent to zero uniformly on compact subsets of \mathbb{D} . Moreover, we have $f_k^{(n)}(\varphi(z_k)) = 0$ and

$$f_k^{(n+1)}(\varphi(z_k)) = - \frac{\alpha(\alpha + 1)(\alpha + 2) \cdots (\alpha + n - 1) \overline{\varphi(z_k)}^{n+1}}{\phi(|\varphi(z_k)|)(1 - |\varphi(z_k)|^2)^{1/q+n+1}}.$$

Therefore for any compact operator $T : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ we have

$$\lim_{n \rightarrow \infty} \|T f_k\|_{\mathcal{Z}_\mu} = 0.$$

By these observations we get that

$$\begin{aligned} \|C_{\varphi,g}^n - T\| &\succeq \limsup_{k \rightarrow \infty} \|C_{\varphi,g}^n f_k - T f_k\|_{\mathcal{Z}_\mu} \\ &\succeq \limsup_{k \rightarrow \infty} \|C_{\varphi,g}^n f_k\|_{\mathcal{Z}_\mu} - \limsup_{k \rightarrow \infty} \|T f_k\|_{\mathcal{Z}_\mu} \\ &\succeq \limsup_{k \rightarrow \infty} \frac{\mu(|z_k|)|\varphi'(z_k)g(z_k)||\varphi(z_k)|^{n+1}}{\phi(|\varphi(z_k)|)(1-|\varphi(z_k)|^2)^{1/q+n+1}}. \end{aligned}$$

And so

$$\|C_{\varphi,g}^n\|_e = \inf_T \|C_{\varphi,g}^n - T\| \succeq \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)g(z)|}{\phi(|\varphi(z)|)(1-|\varphi(z)|^2)^{1/q+n+1}}. \quad (5)$$

Now we define the sequence $\{h_k\}$ as follows

$$h_k(z) = \frac{(1-|\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \left(\frac{\alpha+n+1}{(1-\overline{\varphi(z_k)}z)^\alpha} - \frac{\alpha(1-|\varphi(z_k)|^2)}{(1-\overline{\varphi(z_k)}z)^{\alpha+1}} \right).$$

It is easy to see that

$$\begin{aligned} h_k^{(n)}(z) &= \frac{(1-|\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \frac{\alpha(\alpha+1) \cdots (\alpha+n-1)(\alpha+n+1)\overline{\varphi(z_k)}^n}{(1-\overline{\varphi(z_k)}z)^{\alpha+n}} \\ &\quad - \frac{(1-|\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \\ &\quad \times \frac{\alpha(\alpha+1) \cdots (\alpha+n)(1-|\varphi(z_k)|^2)\overline{\varphi(z_k)}^n}{(\alpha+n)(1-\overline{\varphi(z_k)}z)^{\alpha+n+1}} \\ h_k^{(n+1)}(z) &= \frac{(1-|\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \frac{\alpha(\alpha+1) \cdots (\alpha+n+1)\overline{\varphi(z_k)}^{n+1}}{(1-\overline{\varphi(z_k)}z)^{\alpha+n+1}} \\ &\quad - \frac{(1-|\varphi(z_k)|^2)^{b+1}}{\phi(|\varphi(z_k)|)} \\ &\quad \times \frac{\alpha(\alpha+1) \cdots (\alpha+n+1)(1-|\varphi(z_k)|^2)\overline{\varphi(z_k)}^{n+1}}{(1-\overline{\varphi(z_k)}z)^{\alpha+n+2}}. \end{aligned}$$

Then $\{h_k\}$ is a bounded sequence in $H(p, q, \phi)$ converging to zero uniformly on compact subsets of \mathbb{D} and

$$h_k^{(n)}(\varphi(z_k)) = \frac{\alpha(\alpha+1)(\alpha+2) \cdots (\alpha+n-1)\overline{\varphi(z_k)}^n}{\phi(|\varphi(z_k)|)(1-|\varphi(z_k)|^2)^{1/q+n}}, \quad h_k^{(n+1)}(\varphi(z_k)) = 0.$$

Hence

$$\lim_{n \rightarrow \infty} \|Th_k\|_{\mathcal{Z}_\mu} = 0,$$

for any compact operator $T : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$. By definition of the operator norm we have

$$\begin{aligned} \|C_{\varphi, g}^n - T\| &\succeq \limsup_{k \rightarrow \infty} \|C_{\varphi, g}^n h_k - Th_k\|_{\mathcal{Z}_\mu} \\ &\succeq \limsup_{k \rightarrow \infty} \|C_{\varphi, g}^n h_k\|_{\mathcal{Z}_\mu} - \limsup_{k \rightarrow \infty} \|Th_k\|_{\mathcal{Z}_\mu} \\ &\succeq \limsup_{k \rightarrow \infty} \frac{\mu(|z_k|)|g(z_k)||\varphi(z_k)|^n}{\phi(|\varphi(z_k)|)(1 - |\varphi(z_k)|^2)^{1/q+n+}}. \end{aligned}$$

Therefore, using the definition of the essential norm we obtain

$$\|C_{\varphi, g}^n\|_e = \inf_T \|C_{\varphi, g}^n - T\| \succeq \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n}}. \quad (6)$$

From (5) and (6), the lower estimate of the essential norm is achieved.

Suppose that $\{r_j\}$ be a sequence in $(0, 1)$ and $r_j \rightarrow 1$ as $j \rightarrow \infty$. Since $C_{\varphi, g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded, then it is clear that $C_{r_j \varphi, g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is also bounded. Also, since $\|r_j \varphi\|_\infty < 1$, then by Proposition 2.3, we get that $C_{\varphi, g}^n : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is a compact operator for any j . Thus

$$\|C_{\varphi, g}^n\|_e \leq \|C_{\varphi, g}^n - C_{r_j \varphi, g}^n\| = \sup_{\|f\|_{p, q, \phi} \leq 1} \|(C_{\varphi, g}^n - C_{r_j \varphi, g}^n)f\|_{\mathcal{Z}_\mu}, \quad j \in \mathbb{N}.$$

Now let $f \in H(p, q, \phi)$ and $\|f\|_{p,q,\phi} \leq 1$. Then

$$\begin{aligned}
& \| (C_{\varphi,g}^n - C_{r_j\varphi,g}^n) f \|_{\mathcal{Z}_\mu} = |((C_{\varphi,g}^n - C_{r_j\varphi,g}^n) f)(0)| \\
& \quad + |((C_{\varphi,g}^n - C_{r_j\varphi,g}^n) f)'(0)| + \sup_{z \in \mathbb{D}} \mu(|z|) |((C_{\varphi,g}^n - C_{r_j\varphi,g}^n) f)''(z)| \\
& \leq |(f^{(n)}(\varphi(0)) - f^{(n)}(r_j\varphi(0)))g(0)| \\
& \quad + \sup_{z \in \mathbb{D}} \mu(|z|) |\varphi'(z)g(z)(f^{(n+1)}(\varphi(z)) - r_j f^{(n+1)}(r_j\varphi(z)))| \\
& \quad + \sup_{z \in \mathbb{D}} \mu(|z|) |g'(z)(f^{(n)}(\varphi(z)) - f^{(n)}(r_j\varphi(z)))| \\
& = |(f^{(n)}(\varphi(0)) - f^{(n)}(r_j\varphi(0)))g(0)| \\
& \quad + \sup_{|\varphi(z)| \leq \delta} \mu(|z|) |\varphi'(z)g(z)(f^{(n+1)}(\varphi(z)) - r_j f^{(n+1)}(r_j\varphi(z)))| \\
& \quad + \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |\varphi'(z)g(z)(f^{(n+1)}(\varphi(z)) - r_j f^{(n+1)}(r_j\varphi(z)))| \\
& \quad + \sup_{|\varphi(z)| \leq \delta} \mu(|z|) |g'(z)(f^{(n)}(\varphi(z)) - f^{(n)}(r_j\varphi(z)))| \\
& \quad + \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |g'(z)(f^{(n)}(\varphi(z)) - f^{(n)}(r_j\varphi(z)))|, \tag{7}
\end{aligned}$$

where $\delta \in (0, 1)$, is fixed. Since $f_{r_j} \rightarrow f$ uniformly on compact subsets of \mathbb{D} , where $f_r(z) = f(rz)$, then

$$|(f^{(n)}(\varphi(0)) - f^{(n)}(r_j\varphi(0)))g(0)| \rightarrow 0$$

as $j \rightarrow \infty$. Also

$$\limsup_{j \rightarrow \infty} \sup_{|\varphi(z)| \leq \delta} \mu(|z|) |\varphi'(z)g(z)(f^{(n+1)}(\varphi(z)) - r_j f^{(n+1)}(r_j\varphi(z)))| = 0$$

$$\limsup_{j \rightarrow \infty} \sup_{|\varphi(z)| \leq \delta} \mu(|z|) |g'(z)(f^{(n)}(\varphi(z)) - f^{(n)}(r_j\varphi(z)))| = 0.$$

Here we use (2) and (3) and the fact that $f_{r_j}^{(n+1)} \rightarrow f^{(n+1)}$ and $f_{r_j}^{(n)} \rightarrow f^{(n)}$ uniformly on compact subsets of \mathbb{D} . Also by (7) and Lemma

2.1 we have

$$\begin{aligned}
 & \| (C_{\varphi, g}^n - C_{r_j \varphi, g}^n) f \|_{\mathcal{Z}_\mu} \\
 & \leq \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |\varphi'(z) g(z) (f^{(n+1)}(\varphi(z)) - r_j f^{(n+1)}(r_j \varphi(z)))| \\
 & \quad + \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |g'(z) (f^{(n)}(\varphi(z)) - f^{(n)}(r_j \varphi(z)))| \\
 & \leq \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |\varphi'(z) g(z) f^{(n+1)}(\varphi(z))| \\
 & \quad + \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |\varphi'(z) g(z) r_j f^{(n+1)}(r_j \varphi(z))| \\
 & \quad + \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |g'(z) f^{(n)}(\varphi(z))| \\
 & \quad + \sup_{\delta < |\varphi(z)| < 1} \mu(|z|) |g'(z) f^{(n)}(r_j \varphi(z))| \\
 & \leq 2C \sup_{\delta < |\varphi(z)| < 1} \frac{\mu(|z|) |\varphi'(z) g(z)|}{\phi(|\varphi(z)|) (1 - |\varphi(z)|^2)^{1/q+n+1}} \\
 & \quad + 2C \sup_{\delta < |\varphi(z)| < 1} \frac{\mu(|z|) |g'(z)|}{\phi(|\varphi(z)|) (1 - |\varphi(z)|^2)^{1/q+n}}. \tag{8}
 \end{aligned}$$

So

$$\begin{aligned}
 \|C_{\varphi, g}^n\|_e & \leq \sup_{\|f\|_{p, q, \phi} \leq 1} \| (C_{\varphi, g}^n - C_{r_j \varphi, g}^n) f \|_{\mathcal{Z}_\mu} \\
 & \leq 2C \sup_{\delta < |\varphi(z)| < 1} \frac{\mu(|z|) |\varphi'(z) g(z)|}{\phi(|\varphi(z)|) (1 - |\varphi(z)|^2)^{1/q+n+1}} \\
 & \quad + 2C \sup_{\delta < |\varphi(z)| < 1} \frac{\mu(|z|) |g'(z)|}{\phi(|\varphi(z)|) (1 - |\varphi(z)|^2)^{1/q+n}}.
 \end{aligned}$$

Therefore as $\delta \rightarrow 1$ we obtain

$$\begin{aligned}
 \|C_{\varphi, g}^n\|_e & \leq \max \left\{ \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |\varphi'(z) g(z)|}{\phi(|\varphi(z)|) (1 - |\varphi(z)|^2)^{1/q+n+1}}, \right. \\
 & \quad \left. \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|) |g'(z)|}{\phi(|\varphi(z)|) (1 - |\varphi(z)|^2)^{1/q+n}} \right\}.
 \end{aligned}$$

This completes the proof. \square

If we set $n = 1$ in the Theorem 2.4, then we have the following corollary.

Corollary 2.5. *Let $p, q \in (0, \infty)$, ϕ be normal and φ be a analytic self-map of \mathbb{D} . If the operator $C_\varphi^g : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded, then*

$$\|C_\varphi^g\|_e \approx \max\left\{ \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+2}}, \right. \\ \left. \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|g'(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+1}} \right\},$$

and as a result, $C_\varphi^g : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is compact if and only if

$$\limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+2}} = 0, \\ \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|g'(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+1}} = 0.$$

When $n = 1$ and $g(z) = \varphi'(z)$, then $C_{\varphi, \varphi'}^1 = C_\varphi$, the well-known composition operator, and we have the next corollary.

Corollary 2.6. *Let $p, q \in (0, \infty)$, ϕ is normal and φ be a analytic self-map of \mathbb{D} . If the operator $C_\varphi : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded then*

$$\|C_\varphi\|_e \approx \max\left\{ \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'^2(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+2}}, \right. \\ \left. \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+1}} \right\},$$

and as a result, $C_\varphi : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is compact if and only if

$$\limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'^2(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+2}} = 0, \\ \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|\varphi'(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+1}} = 0.$$

With different choices on ϕ and g and the other parameters of the operator and space, we can obtain some other results.

Theorem 2.7. *Let $p, q \in (0, \infty)$, ϕ is normal and φ be a analytic self-map of \mathbb{D} . If the operator $C_{\varphi, g}^n : H(p, q, \phi) \rightarrow \mathcal{B}_\mu$ is bounded then*

$$\|C_{\varphi, g}^n\|_e \approx \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n}}.$$

The proof is similar to the proof of Theorem 2.4.

Corollary 2.8. *Let $p, q \in (0, \infty)$, ϕ is normal and φ be a analytic self-map of \mathbb{D} . Then the operator $C_{\varphi, g}^n : H(p, q, \phi) \rightarrow \mathcal{B}_\mu$ is compact if and only if*

$$\limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|g(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{1/q+n}} = 0.$$

References

- [1] O. Blasco, Cesàro-type operators on Hardy spaces, *J. Math. Anal. Appl.*, 529 (2023), 127017.
- [2] B. Choe, H. Koo and W. Smith, Composition operators on small spaces, *Integral Equ. Oper. Theory*, 56(3) (2006), 357-380.
- [3] C. C. Cowen and B. D. Maccluer, *Composition Operators on Spaces of Analytic Functions*, CRC Press, Boca Raton, 1995.
- [4] J. Guo and Y. Liu, Generalized integration operators from mixed-norm to Zygmund-type spaces, *Bull. Malay. Math. Soc.*, 39 (2016), 1043-1057.
- [5] S. Li and S. Stević, Generalized composition operators on Zygmund spaces and Bloch type spaces, *J. Math. Anal. Appl.*, 338 (2008) 1282-1295.
- [6] Z. Hu, Extended cesàro operators on mixed norm spaces, *Proc. Am. Math. Soc.*, 131(7) (2002), 2171-2179.

- [7] J. H. Shi and G. P. Ren, Boundedness of the Cesàro operator on mixed norm spaces, *Proc. Amer. Math. Soc.*, 126 (1998), 3553-3560.
- [8] Y. Liu and Y. Yu, Riemann-Stieltjes operator from mixed norm spaces to Zygmund-type spaces on the unit ball, *Taiwan. J. Math.*, 17(5) (2013), 1751-1764.
- [9] Y. Liu and Y. Yu, Weighted Differentiation Composition Operators from Mixed-Norm to Zygmund Spaces, *Numer. Funct. Anal. Optim.*, 31(8) (2010), 936-954.
- [10] S. Stević, Generalized composition operators between mixed-norm and some weighted spaces, *Numer. Funct. Anal. Optim.*, 29(7-8) (2008), 959-978.
- [11] S. Stević, Products of integral-type operators and composition operators from a mixed norm space to Bloch-type spaces, *Sib. Math. J.*, 50(4) (2009), 726-736.
- [12] X. Zhu, An integral-type operator from H^∞ to Zygmund-type spaces, *Bull. Malays. Math. Sci. Soc.*, 35 (2012) 679-686.
- [13] K. Zhu, Bloch type spaces of analytic functions, *Rocky Mountain J. Math.*, 23 (1993), 1143-1177.

Yousef Estaremi

Associate Professor of Mathematics
Department of Mathematics
Faculty of Sciences, Golestan University
Gorgan, Iran
E-mail: y.estaremi@gu.ac.ir

Mostafa Hassanlou

Associate Professor of Mathematics
Department of Mathematics
Engineering Faculty of Khoy, Urmia University of Technology
Urmia, Iran
E-mail: m.hassanlou@uut.ac.ir

Mohammad said Al Ghafri

Assistant Professor of Mathematics
Department of Mathematics
University of Technology and Applied Sciences
Rustaq, Sultanate of Oman
E-mail: mohammed.alghafri@utas.edu.om