

Journal of Mathematical Extension
Vol. 19, No. 2, (2025) (3) 1-19
ISSN: 1735-8299
URL: <http://doi.org/10.30495/JME.2025.3161>
Original Research Paper

Mean Value Functions and Comparative Convexity

M.H. Hooshmand

Shi.C., Islamic Azad University

Abstract. This paper formally defines mean value and pointwise MV-functions associated with a given real function and extends certain aspects of the mean value theorem and its properties. Additionally, we introduce and study an induced concept called comparative convexity (and concavity). As an application of this study, we establish some uniqueness conditions for related functional equations and completely solve several functional inequalities.

AMS Subject Classification: 26E60; 26A51; 39B22; 39B62; 39B72

Keywords and Phrases: Mean value theorem, convex function, functional equation and inequality, limit summability of functions

1 INTRODUCTION

In this paper, we examine certain classes of useful and applicable functions derived from the mean value theorem (MVT). Several articles have explored related functional equations (e.g., see [1, 2, 3, 5, 7, 10]). However, to the best of our knowledge, these works primarily focus on mean value-type theorems, their generalizations, and associated functional equations, without explicitly defining and studying the related functions themselves. In this study, we not only formally define and analyze

Received: September 2024; Accepted: March 2025

such functions but also introduce an induced concept called comparative convexity (and concavity).

2 MEAN VALUE FUNCTIONS of A REAL FUNCTION

Now, let's define MV-functions as follows. During this paper we consider I as a (bounded or unbounded) interval.

Definition 2.1. Let $f : I \rightarrow \mathbb{R}$ be a given function. We call $g : I^\circ \rightarrow \mathbb{R}$ a "Mean Value function of f " (MV-function) if for every two distinct elements $x, y \in I$ there exists a $c (= c_{x,y})$ between them such that $f(y) - f(x) = g(c)(y - x)$.

If f is differentiable on I° and continuous on I , then $g = f'$ is a MV-function of f . The next example shows possibility of existence of other MV-functions even if f is (infinitely) differentiable every where.

Example 2.2. The Dirichlet function $D = \chi_{\mathbb{Q}}$ (that is the indicator of rational numbers) is a MV-function for the identity and all constant functions. Indeed, if f is a constant (resp. the identity) function, then every function g such that $g^{-1}(0)$ (resp. $g^{-1}(1)$) is a dense subset of \mathbb{R} (e.g., $g = \chi_{\mathbb{Q}}$ or $g = 1 - \chi_{\mathbb{Q}}$) is a MV-function of f .

Now, two natural questions arise which are: how large is the class of MV-functions of f ?, and under what conditions f' is the only MV-function?

Theorem 2.3. *Every function $f : I \rightarrow \mathbb{R}$ has infinitely many MV-functions, and cardinality of the class of such functions is 2^c (the biggest possible cardinality, see the next remark).*

But, if f is differentiable and f' (resp. f'^{-1}) is continuous, then f' is the unique continuous (inverse-continuous) MV-function of f .

Proof. Let $g : I \rightarrow \mathbb{R}$ be a function such that preimage of every point (i.e., $g^{-1}(\{t\})$; $t \in \mathbb{R}$) is dense in \bar{I} . For each distinct $x, y \in \mathbb{R}$, we can pick an arbitrary point $t \in g^{-1}\left(\frac{f(y)-f(x)}{y-x}\right) \cap (\min\{x, y\}, \max\{x, y\})$ and put $c = c_{x,y} := t$ in the definition. Now, we show that the cardinality

of such functions g is $2^{\mathfrak{c}}$. A function g has the property if and only if it is everywhere surjective (i.e., it transforms every open interval into the whole real line). For if g is everywhere surjective, then given $x \in \mathbb{R}$ and $(x - \varepsilon, x + \varepsilon) \subseteq I$, we have $f((x - \varepsilon, x + \varepsilon)) = \mathbb{R}$, thus $f^{-1}(t) \in (x - \varepsilon, x + \varepsilon)$ for all $t \in \mathbb{R}$. Since this must hold for every $\varepsilon > 0$ such that $(x - \varepsilon, x + \varepsilon) \subseteq I$, we get the property. Conversely, if the preimage $g^{-1}(t)$ is dense in \bar{I} , for every t , then it must intersect every open set. In particular, it must intersect every set of the form $(x - \varepsilon, x + \varepsilon)$ in I , hence g is everywhere surjective. Therefore, Theorem 4.3 of [4] completes the first part of proof.

For the second part, let f be differentiable and g be a MV-function of f . If g is continuous, then one can easily see that $g = f'$. But, if g is invertible and g^{-1} is continuous, then put $h = g^{-1}$ and consider $x < y$ (without loss of generality). Thus $x < h(\frac{f(y)-f(x)}{y-x}) < y$. By fixing x and $y \rightarrow x$ we get

$$\lim_{y \rightarrow x^+} h\left(\frac{f(y) - f(x)}{y - x}\right) = x.$$

This implies $hf'_+(x) = x$, since h is continuous. Hence $h = f'^{-1}$ and so $g = f'$. \square

Remark 2.4. It is important to know that the family of MV-functions of every function $f : I \rightarrow \mathbb{R}$ is $2^{\mathfrak{c}}$ -lineable as the aspect of pathological properties of functions (see [4, 6]). Also, one can construct infinitely many such functions $g : \mathbb{R} \rightarrow \mathbb{R}$ in the above theorem (that the preimage of every point is a dense subset) explicitly. A famous example of such g is Conway's base 13 function (see [8]). Another method to construct such functions is possible as follows. Consider the additive quotient group \mathbb{R}/\mathbb{Q} with the cardinal number \mathfrak{c} . Thus there is a bijection $\varphi : \mathbb{R}/\mathbb{Q} \rightarrow \mathbb{R}$. Defining $g(x) := \varphi([x]_{\mathbb{R}/\mathbb{Q}})$ gives us a function with the desired property.

As we know behaviors of the derivative of a function determines several properties of that function. Thus, it is very important to know that when the role of a MV-function g is similar to the derivative function.

Lemma 2.5. (*Basic relations between a real function and its MV-functions*)

Let g be a MV-function of $f : I \rightarrow \mathbb{R}$.

(a) If $g \geq 0$ (on I°), then f is increasing. But the converse is not true

(although it is true when $g = f'$).

(b) If g is continuous, then f is differentiable (but the converse is not true) and $g = f'$ on I° .

Hence, every function has at most one continuous MV-function, and the followings are equivalent:

- f is continuously differentiable;

- f has a continuous MV-function;

- f' is the only continuous MV-function of f .

(c) If g is increasing, then f is convex. But the converse is not true (however, if g is continuous then the converse is also valid).

(d) If g is increasing and $0 \in I$, then there exist constants a, b such that

$$a + bx \leq f(x) \leq a + xg(x) ; \quad x \in I^\circ.$$

(e) If g is invertible, then $g^{-1}(\frac{f(y)-f(x)}{y-x})$ is a strict mean function on I (see the next explanation).

Proof. (a) The first part is obvious, and for the second part let $f(t) = t$, and $g(t)$ be 1 on \mathbb{Q} and -1 on \mathbb{Q}^c .

(b) The first statement is easy to prove, and for disproving the converse consider every differentiable function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $g = f'$ is not continuous. Theorem 2.3 proves the rest of sentences in this part.

(c) Let $x < y < z$ be arbitrary elements of I and $c_{x,y}, c_{y,z}$ are as mentioned in Definition 2.1. Then, $x < c_{x,y} < y < c_{y,z} < z$ and $g(c_{x,y}) = \frac{f(y)-f(x)}{y-x}$, $g(c_{y,z}) = \frac{f(z)-f(y)}{z-y}$. These relations imply that f is convex if g is increasing (indeed the inequalities (6) hold). The converse is not true by Example 3.10.

(d) Putting $a = f(0)$ and $b = g(0)$, one can get the result easily from (c).

(e) This is clear by considering the definition. \square

Remark 2.6. Up to now, we have determined the continuous MV-functions of a given function, and in Theorem 3.9, we do for increasing (monotone) MV-functions. Hence, we leave the following important problem and questions:

Problem I. Characterize (or give some necessary and sufficient conditions for the existence of) MV-functions with the intermediate value (Darboux) property, piecewise continuity, left or right continuity, etc.

Question II. Let f be a real function with one-sided derivatives and at most finitely many non-differentiable points. Then, are there some MV-functions of f with at most finitely many non-continuous points? What happens if such points are at most countable?

Question III. Let f be a non-continuously differentiable function. Is it true that f' is the only MV-function with the intermediate value property? If no, what conditions are needed to be true?

Putting $I\check{\times}I := I \times I \setminus \{(x, x) : x \in I\}$, a function $\eta : I\check{\times}I \rightarrow I^\circ$ is called “strict mean function on I ” if $\min\{x, y\} < \eta(x, y) < \max\{x, y\}$, for all $(x, y) \in I\check{\times}I$. Replacing $I\check{\times}I$ by $I \times I$ and $<$ by \leq , η is called “mean function on I ”.

Let g be a MV-function of $f : I \rightarrow \mathbb{R}$. Then, we can construct functions $\eta : I\check{\times}I \rightarrow I^\circ$ such that $\eta(x, y) = c = c_{x,y}$, for a value c in the definition (choice functions). Therefore, η is a strict mean function on I and it can be considered as a symmetric function (i.e. $\eta(x, y) = \eta(y, x)$), and also can be naturally extended on whole $I \times I$, by defining $\eta(x, x) = x$ (to be a mean function).

On the other hand, we have the following equivalent conditions:

$$\min\{x, y\} < \eta(x, y) < \max\{x, y\} \Leftrightarrow 0 < \frac{\eta(x, y) - x}{y - x} < 1 \quad (1)$$

$$\Leftrightarrow 0 < \frac{\eta(x, y) - y}{x - y} < 1 \Leftrightarrow \eta(x, y) = \lambda(x, y)x + (1 - \lambda(x, y))y$$

$$\Leftrightarrow \eta(x, y) = \mu(x, y)y + (1 - \mu(x, y))x,$$

where $0 < \lambda(x, y) < 1$ and $0 < \mu(x, y) < 1$ (of course, $\lambda(x, y) + \mu(x, y) = 1$). Thus, for every strict mean function η there exist functions $\lambda, \mu : I \times I \rightarrow (0, 1)$ such that

$$\eta(x, y) = y + (x - y)\lambda(x, y) = x + (y - x)\mu(x, y),$$

for all $(x, y) \in I \times I$, and vice versa. This fact gives us a simple way to construct strict mean and symmetric strict mean functions.

Proposition 2.7. *For a given function $f : I \rightarrow \mathbb{R}$, the following statements are equivalent:*

- (a) g is a MV-function of f .
 (b) There exists a (symmetric) strict mean function η on I such that

$$f(y) - f(x) = (y - x)g(\eta(x, y)) ; (x, y) \in I \times I.$$

- (c) There exists a (symmetric) function $\lambda : I \times I \rightarrow (0, 1)$ such that

$$f(y) - f(x) = (y - x)g(\lambda(x, y)x + (1 - \lambda(x, y))y) ; (x, y) \in I \times I$$

- (d) There exists a (symmetric) mean function η on I such that

$$f(y) - f(x) = (y - x)g(\eta(x, y)) ; (x, y) \in I \times I.$$

- (e) There exists a (symmetric) function $\lambda : I \times I \rightarrow (0, 1)$ such that

$$f(y) - f(x) = (y - x)g(x + (y - x)\lambda(x, y)) ; x, y \in I, x < y.$$

- (f) There exists a (symmetric) function $\lambda : I \times I \rightarrow (0, 1)$ such that

$$f(y) - f(x) = (y - x)g(\min\{x, y\} + |y - x|\lambda(x, y)) ; (x, y) \in I \times I$$

or equivalently

$$f(y) - f(x) = (y - x)g\left(\frac{x + y}{2} + (\lambda(x, y) - \frac{1}{2})|y - x|\right) ; (x, y) \in I \times I.$$

2.1 Pointwise MV-functions

It may be helpful to study the next easier case of MV-functions.

Definition 2.8. Let $f : I \rightarrow \mathbb{R}$ be a given function and fix an element $x_0 \in I$. We call $g : I^\circ \rightarrow \mathbb{R}$ a “Mean Value function of f at x_0 ” (x_0 -MV-function) if one of the equivalent conditions (a)-(f) (in Proposition 2.7) holds, for every two distinct elements $x, y \in I$ with $x = x_0$.

It is obvious that g is a MV-function of f if and only if it is a x_0 -MV-function for all $x_0 \in I$.

Lemma 2.9. All functions of the form

$$g(t) = g_{\mu, x_0}(t) := \frac{\mu(t)}{t - x_0} \left(f\left(x_0 + \frac{t - x_0}{\mu(t)}\right) - f(x_0) \right),$$

(for all t such that $\frac{t-x_0}{\mu(t)} \in I - x_0$) are x_0 -MV-functions of f , where $\mu : I \rightarrow (0, 1)$ satisfies the functional equation

$$\mu(t\mu(t) - x_0\mu(t) + x_0) = \mu(t) ; t \in I.$$

In particular, putting $\mu(t) = \mu$ where $\mu \in (0, 1)$, we obtain infinitely many such functions, and specially $g_{\frac{1}{2}, x_0}(t) = \frac{f(2t-x_0)-f(x_0)}{2t-2x_0}$, if $\mu = \frac{1}{2}$.

Proof. For every $t \in I \setminus \{x_0\}$ put $z = z_{t, x_0} = \mu(t)t + (1 - \mu(t))x_0$, then

$$\frac{z - x_0}{\mu(t)} = t - x_0 \in I - x_0,$$

so $g(z)$ is defined and we have $g(z) = \frac{f(t)-f(x_0)}{t-x_0}$ \square

A question arises that when a x_0 -MV-function is the same f' . For the case $x_0 = 0$ and $\mu(t) = \mu$ it means that

$$f'(t) = \frac{\mu}{t} \left(f\left(\frac{t}{\mu}\right) - f(0) \right) ; 0 \neq t \in \mu I,$$

and we arrive at the differential equation

$$t f'(t) - \mu f\left(\frac{t}{\mu}\right) + \mu f(0) = 0 ; t \in \mu I.$$

The related integral equation is

$$\int^x \frac{f\left(\frac{1}{\mu}t\right)}{t} dt = \frac{1}{\mu} f(x) + f(0) \log |x| ; x \neq 0.$$

that is interesting and may be useful for the topic. It is obvious that all functions $f(t) = kt$ enjoy the properties.

3 COMPARATIVE CONVEX AND CONCAVE FUNCTIONS

We defined the MV-functions via replacing f' by an arbitrary function g satisfying the equation (and condition). On the other hand a function f is convex on I if and only if $f(x) \geq f(y) + f'(y)(x - y)$, for all $x, y \in I$. Therefore, we are led to the following definition.

Definition 3.1. Let $g : I \rightarrow \mathbb{R}$ be a given function. We call $f : I \rightarrow \mathbb{R}$ “ g -compared convex” if $f(x) \geq f(y) + g(y)(x - y)$, for all $x, y \in I$. Here g is called “difference quotient bound” of f (see the inequality (6)). Analogously, “strictly g -compared convex” and “ g -compared concave” functions are defined.

As a geometric interpretation, f is g -compared convex if and only if for every point $x_0 \in I$ there is a line of support of f with the slope $g(x_0)$. If f is a differentiable convex function, then it is f' -compared convex. More generally, every convex function is f'_{\pm} -compared convex. It is clear that f is g -compared concave if and only if $-f$ is $-g$ -compared convex. Also, f is 0-compared convex if and only if f is constant.

Convention. For comfort, in this paper, we use “ g -convex” instead of “ g -compared convex”.

Example 3.2. The absolute value function is sgn-convex (where sgn denotes the sign function). Indeed $|\cdot|$ is g -convex if and only if there exists $-1 \leq \alpha \leq 1$ such that

$$g(x) = \begin{cases} \text{sgn}(x) & \text{if } x \neq 0 \\ \alpha & \text{if } x = 0 \end{cases}$$

Conversely, f is sgn-convex if and only if $f = |\cdot| + c$ for some constant c (also see Theorem 3.8).

Proposition 3.3. A function $f : I \rightarrow \mathbb{R}$ is g -convex if and only if one of the following equivalent conditions hold:

$$g(x)(y - x) \leq f(y) - f(x) \leq g(y)(y - x) ; \quad x, y \in I. \quad (2)$$

$$f(\lambda x + (1 - \lambda)y) \leq f(y) + \lambda g(x)(x - y) ; \quad x, y \in I, \lambda \in [0, 1]. \quad (3)$$

$$g(x) \leq \frac{f(y) - f(x)}{y - x} \leq g(y) ; \quad x, y \in I, x < y. \quad (4)$$

$$f(\lambda x + (1 - \lambda)y) \leq f(x) + (1 - \lambda)g(y)(y - x) ; \quad x, y \in I, \lambda \in [0, 1]. \quad (5)$$

Proof. If f is g -convex, then by changing the roles of x, y in its inequality, and combining the two inequalities we arrive at (2). Now, let $x < y$ in I and consider $0 < \lambda < 1$. If (2) holds, then

$$g(x) \leq \frac{f(\lambda x + (1 - \lambda)y) - f(x)}{(1 - \lambda)(y - x)} \leq g(\lambda x + (1 - \lambda)y)$$

$$\leq \frac{f(y) - f(\lambda x + (1 - \lambda)y)}{\lambda(y - x)} \leq g(y)$$

Thus, we obtain two inequalities

$$f(\lambda x + (1 - \lambda)y) \leq f(y) + \lambda g(x)(x - y)$$

and

$$f(\lambda x + (1 - \lambda)y) \leq f(x) + (1 - \lambda)g(y)(y - x),$$

for $x < y$ and $0 < \lambda < 1$. By changing the roles of x , y and replacing $1 - \lambda$ by λ in the second inequality, we derive the first inequality for $y < x$ and all $0 < \lambda < 1$. Also, (3) holds for $\lambda = 0, 1$ obviously. Hence, we arrive at (3) for all $x, y \in I$ and $\lambda \in [0, 1]$.

Now, if (3) holds, then putting $\lambda = 1$ we conclude that f is g -convex and so (2) is valid, thus we deduce (4). Notice that (4) \Rightarrow (5) is similar to (2) \Rightarrow (3), and (5) implies g -convexity of f by putting $\lambda = 0$. Therefore, the proof is complete. \square

Remark 3.4. It is easy to show that if I is open or f is continuous, then we can reduce the condition $\lambda \in [0, 1]$ to $\lambda \in (0, 1)$. Now, the question is that whether it is valid without the extra conditions, i.e., does $f(\lambda x + (1 - \lambda)y) \leq f(x) + (1 - \lambda)g(y)(y - x)$ for $0 < \lambda < 1$ (on I) imply that f is g -convex?

An important result of Proposition 3.3 is

$$g(x) \leq \frac{f(y) - f(x)}{y - x} \leq g(y) \leq \frac{f(z) - f(y)}{z - y} \leq g(z), \quad (6)$$

for all $x < y < z$ in I . Hence, we arrive at the following important corollary.

Corollary 3.5. *A function f is convex if and only if it is g -convex for some g . Also, if f is g -convex, the f is convex and g is increasing.*

Now, we are ready to prove the essential relations between f and g for g -convex functions f . It together with Proposition 3.3 enables us to characterize them in several ways.

Theorem 3.6. *If $f : I \rightarrow \mathbb{R}$ is g -convex, then*

$$f'_-(x) = g(x^-) \leq g(x) \leq g(x^+) = f'_+(x) ; x \in I^\circ.$$

Hence, f is differentiable (on I°) if and only if g is continuous, and if this is the case, then $g = f'$ on I° .

Proof. The inequality (6) together with increasingness of g implies that $f'_-(x) \leq g(x^-) \leq g(x) \leq g(x^+) \leq f'_+(x)$. But we claim that $f'_+(x) = g(x^+)$ and $f'_-(x) = g(x^-)$. For fixed $x < y$ in I° consider the function

$$h(t) = f(t) - \frac{f(y) - f(x)}{y - x}(t - x)$$

which has the left derivative

$$h'_-(t) = f'_-(t) - \frac{f(y) - f(x)}{y - x}.$$

We have $h(x) = h(y)$ so that h attains its minimum on the interval $[x, y]$ at some point t_0 with $x < t_0 \leq y$. Then $h'_-(t_0) \leq 0$ so that

$$\frac{f(y) - f(x)}{y - x} = f'_-(t_0) - h'_-(t_0) \geq f'_-(t_0) \geq g(t_0^-) \geq g(x^+).$$

Taking the limit $y \rightarrow x^+$ we conclude that $f'_+(x) \geq g(x^+)$. This proves the first equality; the second one can be proved in the same way. \square
The first corollary states a necessary and sufficient condition on (f, g) for g -convexity of f .

Corollary 3.7. *A function $f : I \rightarrow \mathbb{R}$ is g -convex if and only if it is convex and $f'_- \leq g \leq f'_+$ on I° . Hence, every differentiable convex function has a unique difference quotient bound function (that is f').*

Now, we are ready to state some characterizations of g -convex and their difference quotient bound functions.

Theorem 3.8. *Assume that the interval I is open.*

(a) *(A characterization of g -convex functions). Let $g : I \rightarrow \mathbb{R}$ be a given*

increasing function. Then, $f : I \rightarrow \mathbb{R}$ is g -convex if and only if there is a constant $c \in I$ such that for all $x \in I$,

$$f(x) - f(c) = \int_c^x g(t)dt. \quad (7)$$

(b) (A characterization of difference quotient bounds of f).

Let $f : I \rightarrow \mathbb{R}$ be a given convex function. Then, $g : I \rightarrow \mathbb{R}$ is a difference quotient bound of f if and only if there exists $J \subseteq I$ such that f is differentiable on J , $I \setminus J$ is at most countable (thus $I \setminus J = \{c_n\}_{n=1}^N$ or \emptyset , for some $N \in \mathbb{N} \cup \{\aleph_0\}$) and

$$g(x) = \begin{cases} f'(x) & \text{if } x \in J \\ d_n & \text{if } x = c_n \in I \setminus J \end{cases}$$

for some $d_n \in [f'_-(c_n), f'_+(c_n)]$.

Proof. The above corollary implies that (f is convex and so) $g = f'$ on I except possibly in at most countable points (i.e., there exists $J \subseteq I$ such that f is differentiable on J , $I \setminus J$ is at most countable and thus $I \setminus J = \{c_n\}_{n=1}^N$ or \emptyset , for some $N \in \mathbb{N}$). Also, we have $f'_- \leq g \leq f'_+$ on I . Therefore, there exists $c \in I$ such that

$$f(x) - f(c) = \int_c^x f'_+(t)dt = \int_c^x f'_-(t)dt = \int_c^x g(t)dt.$$

Now, one can complete the proof easily. \square

3.1 Relations with MV-functions

The following theorem says that if a function has an increasing MV-function g , then it is g convex, and characterizes such functions g . Hence, it can be considered as a generalization of a famous fact about real convex functions.

Theorem 3.9. *If $f : I \rightarrow \mathbb{R}$ has an increasing MV-function g , then it is g -convex (but the converse is not true), hence*

$$f'_-(x) = g(x^-) \leq g(x) \leq g(x^+) = f'_+(x); \quad x \in I^\circ,$$

and the inequalities (2), (3), (4), (5), and (6) hold.

Proof. This is a direct result of Lemma 2.5(c), Theorem 3.6 and the next example. \square

Now, we show that there exists a convex function without any increasing MV-function.

Example 3.10. The real function $f(x) = |x|$ does not have any increasing (monoton) MV-function (although it is convex). For if g is an increasing MV-function of f , then g is the same as mentioned in Example 3.2, and cannot satisfy one of the equivalent conditions in Proposition 2.7 (because the equation $|y| - |x| = g(\eta(x, y))(y - x)$ is impossible for all $x < 0 < y$). Note that, here, f'_\pm are increasing but not MV-functions of f .

Considering the above example, we arrive at the following problem.

Problem IV. Give an example of a function f with more than one increasing MV-function (f must be convex and non-differentiable at some points).

4 MEAN VALUE AND COMPARATIVE CONVEX FUNCTIONAL EQUATIONS AND INEQUALITIES

Now, we focus on functional equations and inequalities induced by the topics MV-functions and compared convexity.

4.1 MV-functional equations and inequalities

The equivalent conditions mentioned in Propositions 2.7 and 3.3 give us several equivalent functional equations and inequalities. Now, we will discuss them.

Let I be an interval, $f : I \rightarrow \mathbb{R}$ and $g : I^\circ \rightarrow \mathbb{R}$ real functions, and $\eta : I \times I \rightarrow I^\circ$ a strict mean function, $\lambda : I \times I \rightarrow (0, 1)$ and $h : D_h \rightarrow \mathbb{R}$ (where D_h contains the image of the function $\frac{f(y)-f(x)}{y-x}$ for all $(x, y) \in I \check{\times} I$). Note that some of these functions are unknown and others known functions. Then, we call (8) (resp. (9)) MV-functional equation (resp.

MV-functional inequality).

$$\frac{f(y) - f(x)}{y - x} = g(\eta(x, y)) ; (x, y) \in I \check{\times} I \quad (8)$$

$$0 < \frac{1}{y - x} (h(\frac{f(y) - f(x)}{y - x}) - x) < 1 ; (x, y) \in I \check{\times} I \quad (9)$$

It is interesting to know that the MV-equation and inequality have close relations if $h = g^{-1}$. In this case, every solution (f, g) of (8) satisfies (9). But the converse is true if g is monotone and enjoys the Darboux property. For if it is so, then (9) implies that $\frac{f(y) - f(x)}{y - x}$ (for $(x, y) \in I \check{\times} I$) lies between x and y , and so g is a monotonically continuous MV-function of f , hence (f, g) satisfies (8) (for some strict mean function η). Note that (1) provides several equivalent inequalities to (9) that one of them is

$$\min\{x, y\} < h(\frac{f(y) - f(x)}{y - x}) < \max\{x, y\} ; (x, y) \in I \check{\times} I$$

For studying MV-equations, we divided them to the following types:

(a) f is a known function.

The MV-equation has infinitely many solutions (g, η) . For if $\eta : I \check{\times} I \rightarrow I$ is an arbitrary invertible strict mean, then for every $t \in I^\circ$ there exists a unique $(x_t, y_t) \in I \check{\times} I$ such that $t = \eta(x_t, y_t)$. Now, defining $g = g_\eta$ (on I°) by $g(t) := \frac{f(y_t) - f(x_t)}{y_t - x_t}$ we observe that (g, η) satisfies the equation. Since the family of such functions η is infinite, we are done. Also, the MV-inequality has infinitely many solutions h , by Theorem 2.3 and the above explanation. Therefore, a uniqueness theorem should be useful for this case.

Lemma 4.1. *(A uniqueness conditions for the MV-equation and inequality).*

If f is differentiable and f' is invertible, then $h = f'^{-1}$ is a solution of the MV-inequality. Moreover, if f'^{-1} is continuous (e.g., if f' is strictly increasing and continuous) then $h = f'^{-1}$ (resp. $g = f'$) is the unique solution of the MV-inequality (resp. MV-equation) with the properties.

Proof. The first part of the claim is clear by Lemma 2.5(e) and (1). Now, let h be an arbitrary solution of the inequality and $x < y$ (without

lose of generality). Then, $x < h(\frac{f(y)-f(x)}{y-x}) < y$. By fixing x and $y \rightarrow x$ we get

$$\lim_{y \rightarrow x^+} h\left(\frac{f(y) - f(x)}{y - x}\right) = x.$$

This implies $hf'_+(x) = x$, since h is continuous. Thus the assumption yields $h = f'^{-1}$. Therefore, $g = h^{-1} = f'$ is also a unique solution of the MV-equation. \square

(b) g (resp. h) is a known function. For this case, we have also infinitely many solutions (f, η) (resp. f) for (8) (resp. (9)), e.g., put $\eta(x, y) = c$ and $f(t) = g(c)t$, for all constants $c \in I$. However, the next theorem gives the general solution of them under some conditions on g and h .

Theorem 4.2. *If h (resp. g) is strictly monoton and continuous, then all solutions of the MV-inequality (resp. MV-equation) are of the form*

$$f(t) = \int^t h^{-1}(s)ds + c,$$

(resp.

$$f(t) = \int^t g(s)ds + c,$$

$$\eta(x, y) = g^{-1}\left(\frac{f(y) - f(x)}{y - x}\right),$$

of course, $\min\{x, y\} < \eta(x, y) < \max\{x, y\}$), for all real constants c .

Proof. If $f(t) = \int^t h^{-1}(s)ds + c$ and $x < y$ then there exists a $x < z < y$ such that

$$\frac{f(y) - f(x)}{y - x} = \frac{1}{y - x} \int_x^y h^{-1}(s)ds = h^{-1}(z),$$

and so, the inequality (9) is equivalent to $0 < \frac{z-x}{y-x} < 1$ that is clearly true. For uniqueness, if h is strictly monotonic and continuous, then

$$h^{-1}(x) < \frac{f(y) - f(x)}{y - x} < h^{-1}(y),$$

whenever $x < y$. Now, fixing x and $y \rightarrow x^+$ we get $h^{-1}(x) = f'_+(x)$, and in a similar way $h^{-1}(y) = f'_-(y)$. Therefore, $f' = h^{-1}$ on the related interval, and the first part (the MV-inequality) is proved. Proof for the MV-equation is similar. \square

(c) η is a known function.

In this case, the MV-equation has infinitely many solutions, obviously. As a special interesting equation, we have

$$\frac{f(y) - f(x)}{y - x} = g(\lambda x + \mu y),$$

where $\lambda, \mu \geq 0$ and $\lambda + \mu = 1$ ($\eta(x, y) = \lambda x + \mu y$). One can see its general solution in [10].

(d) Two of the functions f , g and η are known.

The MV-equation does not have a solution necessarily. For if $\eta(x, y) = (1 - \lambda)x + \lambda y$ where $0 < \lambda < 1$ is a constant, then f must be a polynomial of order at most two (see, for example, [10]). Also, if $g \equiv 0$ and f is not one-to-one, then they do not satisfy the equation.

4.2 Compared convex and concave functional inequalities

Now, we consider the functional inequalities associated to the g -convexity (and concavity). The obtained results of the topic enable us to solve several such functional inequalities. Hence, we first state their general solutions as a theorem, and then completely solve several functional inequalities and related systems, as some applications of the study. The following theorem is a direct result of Theorem 3.8 and Proposition 3.3.

Theorem 4.3. *The (g -comparative convex) functional inequality*

$$f(x) \geq f(y) + g(y)(x - y) \ ; \ x, y \in I.$$

has a solution if and only if f is convex, g is increasing and $f'_+(x) = g(x^+)$, $f'_-(x) = g(x^-)$ on I° . Moreover,

(a) If g is known and I open, then the general solution is

$$f(x) - f(c) = \int_c^x g(t)dt = \int_c^x g^\pm(t)dt \ ; \ x \in I,$$

where c is an arbitrary fixed point in I .

(b) If f is given, then the general solution is

$$g(x) = \lambda(x)f'_-(x) + (1 - \lambda(x))f'_+(x) ; x \in I^\circ,$$

for all functions $\lambda : I^\circ \rightarrow [0, 1]$.

Another general solution is of the form

$$g(x) = \begin{cases} f'(x) & \text{if } x \in I_f \\ d_n & \text{if } x = c_n \in I \setminus I_f \end{cases}$$

where I is open, $I_f := \{x \in I : f \text{ is differentiable at } x\}$, $I \setminus I_f = \{c_n\}_{n=1}^N$ or \emptyset , $d_n \in [f'_-(c_n), f'_+(c_n)]$.

5 SOME APPLICATIONS of THE COMPARATIVE CONVEXITY AND CONCAVITY

General solutions of the following functional inequalities and related systems are obtained from the topic of g -convexity and g -concavity. In continuation, the interval I is assumed open.

Example 5.1. General solutions of the functional inequality (self-convexity that is equivalent to f -convexity of f):

$$f(x) \geq f(y)(1 + x - y) ; x, y \in I, \quad (10)$$

is of the form $f(t) = \lambda e^t$ with $\lambda \geq 0$. Thus, general solution of

$$\frac{f(x)}{f(y)} \geq 1 + x - y ; x, y \in \mathbb{R}$$

is λe^t , for all $\lambda \neq 0$.

Because f must be continuous (since it is convex and I open) so $f' = f$ and it is increasing (by Theorem 3.6 and Corollary 3.5). Thus $f(t) = \lambda e^t$ (for a constant $\lambda \geq 0$ and all $t \in \mathbb{R}$). Therefore, the natural exponential function is the only real function satisfying (10) (or the second inequality) with the initial condition $f(0) = 1$.

A similar inequality and its general solution can be stated for the self-concavity. As a generalization of (10), let ϕ be a given continuous function on I , k a real constant and consider the functional inequality

$$f(x) \geq f(y) + (kf + \phi)(y)(x - y); \quad x, y \in I. \quad (11)$$

If it has a solution, then f satisfies the linear differential equation $f' - kf = \phi$ on I . Thus, f can be determined through (11) and the condition that $kf + \phi$ must be increasing. For example, putting $\phi = 0$ on I , the general solution of $f(x) \geq f(y) + kf(y)(x - y)$ is of the form $f(t) = \lambda e^{kt}$ with $\lambda, k \geq 0$.

Example 5.2. General solution of the system of functional inequalities (symmetric convexity, that says f is g -convex and g is f -convex):

$$\begin{cases} f(x) \geq f(y) + g(y)(x - y) \\ g(x) \geq g(y) + f(y)(x - y) \end{cases}; \quad x \in I \quad (12)$$

is

$$f(t) = \lambda_1 e^t + \lambda_2 e^{-t}, \quad g(t) = \mu_1 e^t + \mu_2 e^{-t},$$

for all $t \in I$ and $\lambda_1, \mu_1, \lambda_2, \mu_2$ such that $\lambda_1 \theta_I \geq |\lambda_2|$ and $\mu_1 \theta_I \geq |\mu_2|$, where $\theta_I = \inf_{t \in I} e^{2t}$.

Because both f and g (where (f, g) is a solution of (12)) must be continuous and so $f' = g, g' = f$. The conditions on the coefficients come from increasingness of them. Note that if $I = \mathbb{R}$, then the general solution is simplified to the form $f(t) = \lambda e^t, g(t) = \mu e^t$ for $\lambda, \mu \geq 0$ ($t \in \mathbb{R}$).

Example 5.3. General solution of the system of functional inequalities (convex-concavity, that says f is g -convex and h -concave):

$$\begin{cases} f(x) \geq f(y) + g(y)(x - y) \\ f(x) \leq f(y) + h(y)(x - y) \end{cases}; \quad x \in I \quad (13)$$

is $f(t) = at + b, g(t) = h(t) = a$ ($t \in I$), for all constants a, b .

Because (13) implies that $f' = g = h$, g must be increasing and h decreasing.

Acknowledgements

The author would like to thank the people who answered or sent their feedback and comments to some of my questions (related to this paper) in the math.stackexchange website.

References

- [1] J. Aczel, A mean value property of the derivative of quadratic polynomials - without mean values and derivatives, *Math. Mag.*, 58(1)(1985), 42-45.
- [2] J. Aczel and M. Kuczma, On two mean value properties and functional associated with them, *Aequat. Math.*, 38 (1989), 216–235.
- [3] J. Aczel and M. Kuczma, On two related types of functional equations describing mean value properties, *Zeszyty Nauk. Politech. Slask. Mat.-Fiz.*, 64 (1990), 27-35.
- [4] R.M. Aron, V.I. Gurariy, J.B. Seoane-Sepulveda, Lineability and spaceability of sets of functions on \mathbb{R} , *Proc. Amer. Math. Soc.*, 133 (2005), 795-803.
- [5] Z.M. Balogh, O.O. Ibrogimov and B.S. Mityagin, Functional equations and the Cauchy mean value theorem, *Aequat. Math.*, 90(2016), 683-697.
- [6] A. Farokhinia, , Lineability of space of quasi-everywhere surjective functions, *J. Maht. Ext.*, 6(3)(2012), 45-51.
- [7] J. Matkowski, Mean value property and associated functional equations, *Aequat. Math.*, 48(1999), 46-59.
- [8] D. G. Radcliffem, A Function that is Surjective on Every Interval, *Am. Math. Mon.*, 123(1)(2016), 88-89.
- [9] A.W. Roberts, D.E. Varberg, *Convex Functions*, Academic Press, 1973.
- [10] P. Sahoo and T. Riedel, *Mean value theorems and functional equations*, World Scientific, 1998.

Mohammad Hadi Hooshmand

Professor of Mathematics

Department of Mathematics, Shi.C., Islamic Azad University

Shiraz, Iran

E-mail: MH.Hooshmand@iau.ac.ir