

Solution and Hyperstability of Orthogonally Triple Lie Hom-derivations

J. Izadi*

Payame Noor University

M. E. Samei

Bu-Ali Sina University

Abstract. The goal of this paper introduces a generalized concept which is called the orthogonally Jensen s -functional equation, on triple Lie algebras while preserving orthogonality and presents its additive properties with preserving orthogonality. Further, the orthogonally triple Lie hom-derivation associated with Jensen s -functional equation on orthogonally triple Lie algebras, are described. Ultimately, based on fixed point method with the orthogonally conditions, we approach to delve into stability of Hyers-Ulam sense and hyperstability of this s -functional equation and the triple Lie hom-derivation with preserving orthogonality on orthogonally triple Lie algebras.

AMS Subject Classification: MSC 17A40; 39B52; 17B40

Keywords and Phrases: Hyperstability, orthogonally fixed point method, orthogonally triple Lie hom-derivations, Hyers-Ulam stability,

1 Introduction and Auxiliary Concepts

It seems that the pioneering use of ternary operations (TOs) with cubic matrices introduced by Cayley as an outstanding mathematician, in the 19th century. In fact, a TO is an n -ary operation with $n = 3$, such that for any given three elements of a set \mathcal{B} , mixes them to form a

Received: December 2024; Accepted: July 2025

single element of \mathcal{B} . Generally, the non-trivial TO is formulated by the following composition rule for $s, x, w \in \mathfrak{B}$:

$$(s, x, w)_{ijk} = \sum_{\ell, m, n} s_{ni\ell} x_{\ell jm} w_{mkn}, \quad i, j, k = 1, 2, \dots, \aleph, \aleph \in \mathbb{N}.$$

Following this innovation concept, n -ary algebraic structures have been developed and have been applied across a broad spectrum of disciplines, particularly those incorporating TOs [14, 27]. These include but are not limited to data processing, the realm of quantum mechanics, the intricate domain of mathematical physics, and notably in Nambu mechanics as evidenced by the references [25, 37, 41].

In 2006, Amyari and Moslehian introduced groundbreaking concepts related to ternary algebras [2]. A ternary algebra \mathfrak{B} , equipped with a TO, $(s_1, s_2, s_3) \mapsto [s_1, s_2, s_3]$ mapping from \mathfrak{B}^3 to \mathfrak{B} , constitutes a complex space where the operation is linear in the first and third arguments, and conjugate linear in the second argument, both respect to \mathbb{C} [2]. It is associative in that for each $s_i \in \mathfrak{B}$, the equality

$$[[s_1, s_2, s_3], s_4, s_5] = [s_1, [s_2, s_3, s_4], s_5] = [s_1, s_2, [s_3, s_4, s_5]],$$

holds, according to established results in [7]. Further, it adheres to the norm conditions,

$$\|[s_1, s_2, s_3]\| \leq \|s_1\| \cdot \|s_2\| \cdot \|s_3\|, \quad \|[s, s, s]\| = \|s\|^3,$$

and such a ternary algebra \mathfrak{B} forms a B-space is referred to as a ternary Banach algebra.

On the other hand, Lie algebras are a fundamental key in scope of mathematics that are named after the Norwegian mathematician Sophus Lie (1842-1899). In 1893, the work of Scheffers compiled much of the foundational knowledge of Lie algebras based on Lie's own lectures in Leipzig [32]. A Banach algebra can be defined as a Lie algebra, with the Lie product of the form:

$$[s_1, s_2] := \frac{1}{2}(s_1 s_2 - s_2 s_1), \quad s_i \in \mathcal{B}.$$

Extending this, for a triple Lie algebra, the product is defined by [29],

$$[[s_1, s_2], s_3] := \frac{1}{2}([s_1, s_2] s_3 - s_3 [s_1, s_2]), \quad s_i \in \mathcal{B}.$$

Lie theory's application extends beyond mathematics to include fields such as Physics, Engineering, Mathematical Finance, and Economics due to its problem-solving capabilities [21]. An application of Lie groups and algebras can be seen in Lie-Poisson dynamics, which are essential to formulate basic physical equations, that vividly demonstrate Lie-Poisson dynamics of Euler's equations for rigid body dynamics [44]. These equations make use of a Lie-Poisson bracket that is derived from the Lie algebra of infinitesimal rotations which highlights how the principles of Lie algebra and Poisson brackets are applied for analyzing the dynamic behavior of rigid bodies, showcasing the interplay between mathematical structures and physical systems [8, 24, 35]. In this regard, Lo and Lui demonstrate the use of Lie algebra techniques in evaluating financial derivatives, especially those involving multiple assets in [33, 34].

In 1940, Ulam introduced the concept of stability in functional equations (FEs) which raised the question of whether functions that are approximately solutions to a FE are also close to being true solutions of that equation [48]. This inquiry sparked a new area of study focused on investigating the stability properties of FEs, delving into the nuances in the input of a FE that impact the proximity of the output to the true solution. In 1941, Hyers became the first to provide a positive response in Theorem 1.1, to Ulam's query within the context of B-spaces for additive functions [20].

Theorem 1.1 ([20]). *Assume that \mathcal{Y} and \mathcal{B} be a normed and a Banach spaces respectively, and $\mathbb{k} : \mathcal{Y} \rightarrow \mathcal{B}$ be a mapping such that for a given $\varepsilon > 0$, the inequality*

$$\|\mathbb{k}(s + \grave{s}) - \mathbb{k}(s) - \mathbb{k}(\grave{s})\| \leq \varepsilon, \quad \forall s, \grave{s} \in \mathcal{Y},$$

holds. Then there exists a unique additive mapping $\mathfrak{u} : \mathcal{Y} \rightarrow \mathcal{B}$, with $\|\mathbb{k}(s) - \mathfrak{u}(s)\| \leq \varepsilon$ for each $s \in \mathcal{Y}$.

Subsequently, in 1978, Rassias expanded Hyers' work by introducing the generalized stability of Hyers-Ulam (HU) and a novel stability concept utilizing a control function for additive mappings, in Theorem 1.2.

Theorem 1.2 ([45]). *Let \mathcal{Y}, \mathcal{B} be real normed spaces with \mathcal{B} complete, and $\mathbb{k} : \mathcal{Y} \rightarrow \mathcal{B}$ be a mapping such that for each $s_{\circ} \in \mathcal{Y}$, the mapping*

$\mathbb{h}(s) = \mathbb{k}(s s_0)$ is continuous on \mathbb{R} . If for a given $\varepsilon \geq 0$, the inequality,

$$\|\mathbb{k}(s + \dot{s}) - \mathbb{k}(s) - \mathbb{k}(\dot{s})\| \leq \varepsilon(\|s\|^\xi + \|\dot{s}\|^\xi), \quad 0 \leq \xi < 1,$$

holds for $s, \dot{s} \in \mathcal{Y}$, then there exists a unique linear mapping $\mathfrak{u} : \mathcal{Y} \rightarrow \mathcal{B}$ such that

$$\|\mathbb{k}(s) - \mathfrak{u}(s)\| \leq \frac{\varepsilon\|s\|^\xi}{1-2^\xi-1}, \quad \forall s \in \mathcal{Y}.$$

Later, in 1994, Găvruta modified control function of Rassias to $\varphi(s, \dot{s})$ and demonstrated its stability in the next Theorem 1.3.

Theorem 1.3 ([17]). Let $(\mathcal{G}, +)$ be an abelian group, $(\mathcal{B}, \|\cdot\|)$ be a B-space, and consider a mapping $\varphi : \mathcal{G} \times \mathcal{G} \rightarrow \mathbb{R}^{\geq 0}$ with

$$\hat{\varphi}(s, \dot{s}) := \sum_{k=0}^{\infty} \frac{1}{2^k} \varphi(2^k s, 2^k \dot{s}) < \infty, \quad s, \dot{s} \in \mathcal{G}.$$

If $\mathbb{k} : \mathcal{G} \rightarrow \mathcal{B}$ be a mapping such that for each $s, \dot{s} \in \mathcal{G}$,

$$\|\mathbb{k}(s + \dot{s}) - \mathbb{k}(s) - \mathbb{k}(\dot{s})\| \leq \varphi(s, \dot{s}),$$

then there exists a unique mapping $\mathfrak{u} : \mathcal{G} \rightarrow \mathcal{B}$ such that for $s, \dot{s} \in \mathcal{G}$, we have $\mathfrak{u}(s + \dot{s}) = \mathfrak{u}(s) + \mathfrak{u}(\dot{s})$, and

$$\|\mathbb{k}(s) - \mathfrak{u}(s)\| \leq \frac{1}{2} \hat{\varphi}(s, s).$$

The investigation into the stability of FEs, fixed point (F.P) theory, linear operators, Fibonacci numbers, and optimization theory across different spaces, has been pursued by numerous scholars [1, 13, 39]. The study of stability in the realm of FEs on B-spaces, Obloza [40], in 1993, initiated the examination of stability for linear FEs. This opened up avenues for extensive research into the stability of fractional differential equations (FDEs), with several researchers extending UH stability to this domain [11, 19, 28, 36]. Aruchamy *et al.* in [3], studied the existence and stability results for the fractional nonlinear reaction-diffusion equation involving the Caputo fractional derivative with respect to the time variable of order $\alpha \in (0, 1)$, of the form:

$${}^C D_\xi^\alpha \mathfrak{u}(s, \xi) = \mathbb{D} \frac{\partial^2 \mathfrak{u}(s, \xi)}{\partial s^2} + \int_0^\xi a(\xi - r) \frac{\partial^2 \mathfrak{u}(s, r)}{\partial s^2} dr + g(\xi, \mathfrak{u}(s, \xi)),$$

for $s \in \Omega := (0, L)$, $\xi \geq 0$, with the positive kernel function $a(\xi - r)$ and

$$\begin{cases} u(s, 0) = u_0(s), & s \in \Omega, \\ u(s, \xi) = 0, & s \in \partial\Omega, \xi \geq 0. \end{cases}$$

Definition 1.4 ([15]). A binary relation $\perp \subseteq \mathfrak{B} \times \mathfrak{B}$, for the set $\mathfrak{B} \neq \emptyset$ has property orthogonally set (O -set) whenever there exists $s_o \in \mathfrak{B}$ such that, for each $s \in \mathfrak{B}$ either $s \perp s_o$ or $s_o \perp s$, and this O -set is denoted by (\mathfrak{B}, \perp) .

We will investigate some examples of O -sets based on the expressed concepts.

Example 1.5. Let $\mathfrak{B} = \mathbb{Z}$. Define $s \perp \dot{s}$ if and only if $\dot{s} \stackrel{s}{\equiv} 1$. Then (\mathfrak{B}, \perp) is an O -set. Further, $1 \perp s$ for every s .

Example 1.6. Let $\mathfrak{B} = \mathbb{R}^{\geq 0}$. Define $s \perp \dot{s}$ if $s \dot{s} \leq \min \{s, \dot{s}\}$. To get the desired result, it is enough to put $s = 0$.

Example 1.7. In the inner product space \mathfrak{B} with the inner product $\langle \cdot, \cdot \rangle$, we define $s \perp \dot{s}$ if and only if $\langle s, \dot{s} \rangle = 0$. One can easily check that $0 \perp s$ for each $s \in \mathfrak{B}$ and (\mathfrak{B}, \perp) is an O -set.

The O -set (\mathfrak{B}, \perp) is said to have property orthogonally generalized metric space (O -metric space), if (\mathfrak{B}, d) be a generalized metric space and denoted by (\mathfrak{B}, \perp, d) . We say a sequence $\{s_n\}_{n \in \mathbb{N}}$ in O -set (\mathfrak{B}, \perp) be an orthogonally sequence (O -sequence) if either $s_n \perp s_{n+1}$ or $s_{n+1} \perp s_n$ for $n \in \mathbb{N}$. Additionally, an O -sequence $\{s_n\}_{n \in \mathbb{N}} \subseteq \mathfrak{B}$ is called orthogonally Cauchy (O -Cauchy) when $\lim_{\ell, n \rightarrow \infty} d(s_\ell, s_n) = 0$. By concluding that, we say (\mathfrak{B}, \perp, d) is O -complete whenever every O -Cauchy sequence converges in \mathfrak{B} . In O -metric space (\mathfrak{B}, \perp, d) , a mapping $\mathbb{k} : \mathfrak{B} \rightarrow \mathfrak{B}$ is said to be orthogonally continuous (\perp -continuous) in $s \in \mathfrak{B}$ if, for each O -sequence $\{s_n\}_{n \in \mathbb{N}} \subseteq \mathfrak{B}$ with $s_n \rightarrow s$, we have, $\mathbb{k}(s_n) \rightarrow \mathbb{k}(s)$. Further, a mapping \mathbb{k} is called orthogonal contraction (\perp -contraction) via Lipschitz constant $0 \leq \gamma < 1$ when, for $s, \dot{s} \in \mathfrak{B}$,

$$d(\mathbb{k}(s), \mathbb{k}(\dot{s})) \leq \gamma d(s, \dot{s}), \quad s \perp \dot{s}.$$

We say \mathbb{k} is orthogonally preserving (\perp -preserving) if each $s \perp \dot{s}$ implies $\mathbb{k}(s) \perp \mathbb{k}(\dot{s})$.

Example 1.8. Let $\mathfrak{B} = [0, 1) \subset \mathbb{R}$ and d be the standard metric on \mathfrak{B} . Define the mapping $\mathbb{k}(s) = \sqrt{s}$ and $s \perp \dot{s}$ if $s \dot{s} \leq s$. Then, \mathfrak{B} is a \perp -contraction.

Example 1.9. Let $\mathfrak{B} = [0, 10)$ and d be the standard metric on \mathfrak{B} . Define the mapping $\mathbb{k}(s) = s^2$ and $s \perp \dot{s}$ if $s \leq \dot{s} \leq \frac{1}{2}$ and $s \dot{s} \leq s$. Easily, one can see \mathfrak{B} is a \perp -contraction. However, \mathfrak{B} isn't a contraction because, for $s = 0.4$ and $\dot{s} = 0.9$, we have,

$$d(\mathbb{k}(s), \mathbb{k}(\dot{s})) = d\left(\frac{81}{100}, \frac{4}{25}\right) = 0.65 > \gamma d\left(\frac{9}{10}, \frac{4}{10}\right).$$

This notion has been extensively explored in literatures [18, 22, 30]. Chandok *et al.* proposed some weaker orthogonal mathematical equation type of contraction mappings in the setting of metric spaces endowed with an orthogonal relation, as well as certain sufficient criteria for the existence of F.Ps for this class of mapping and used their results to investigate the solution and stability of Hyers-Ulam- Rassias - Wright sense, for the following Volterra type integral equation:

$$\mathfrak{u}(s) = \int_0^s g(s, r, \mathfrak{u}(r)) + F(s), \quad s \in \Omega,$$

under some Lipschitz conditions [10]. Nazam *et al.* by introducing (Ψ, Φ) -orthogonal interpolative contractions, investigated different conditions on the functions $\Psi, \Phi : (0, \infty) \rightarrow \mathbb{R}$ to prove the existence of F.Ps of set-valued (Ψ, Φ) -orthogonal interpolative contractions such that, for $s, \dot{s} \in \mathcal{B}$,

$$\Psi(\mathbb{k}(s, \dot{s})H(\mathcal{T}s, \mathcal{T}\dot{s})) \leq \Phi\left((d(s, \mathcal{T}s))^\gamma (d(\dot{s}, \mathcal{T}\dot{s}))^{1-\gamma}\right), \quad (1)$$

where \mathbb{k} is a strictly \perp -admissible mapping, $0 < \gamma < 1$, $s \notin \mathcal{T}(s)$, $\dot{s} \notin \mathcal{T}(\dot{s})$,

$$H(B_1, B_2) = \max \left\{ \sup_{p \in B_1} d(p, B_2), \sup_{a \in B_2} d(a, B_1) \right\} > 0,$$

B_1, B_2 in the set of all non-empty bounded and closed subsets of \mathcal{B} , $\mathcal{P}_{cb}(\mathcal{B})$, and to solve a \mathbb{FDE} involving Caputo-Fabrizio derivative in O -complete, of the form:

$$\begin{cases} {}^{\text{CF}}D_s^\alpha \mathfrak{u}(s) = g(s, \mathfrak{u}(s)), & s \in \Omega, \\ \mathfrak{u}(0) = 0, \quad I^\alpha \mathfrak{u}(1) = \mathfrak{u}'(0), \end{cases}$$

where $\mathfrak{u} \in C(\Omega)$ and $g : (\Omega \times \mathbb{R}) \rightarrow \mathbb{R}$ with a few conditions [38]. In 2018, Bahraini *et al.* in [6], demonstrated the innovative use of the O -set concept to extend a generalization of Diaz-Margolis's F.P theorem [12]. In 2019, Park *et al.* introduced hom-derivations on Banach algebras [42]. Then, Jahedi and Keshavarz proved ternary additive-quadratic hom-derivations [23].

In this research, motivated by the mentioned works, we first introduce orthogonally Jensen s -functional (\mathcal{J}_{sf}), $s \neq 0, \pm 1$ is a real number and orthogonally triple Lie hom-derivations on orthogonally triple Lie algebras, respectively. Then, we explore the FE of the form,

$$\begin{aligned} 3\mathcal{J}_{sf}\left(\frac{s_1+s_2+s_3}{3}\right) - \mathcal{J}_{sf}(s_1) - \mathcal{J}_{sf}(s_2) - \mathcal{J}_{sf}(s_3) \\ = s \left[\mathcal{J}_{sf}(s_1 + s_2 + s_3) + \mathcal{J}_{sf}(s_1) \right. \\ \left. - \mathcal{J}_{sf}(s_1 + s_2) - \mathcal{J}_{sf}(s_1 + s_3) \right], \end{aligned} \quad (2)$$

for $s_i \in \mathfrak{B}$ ($i = 1, 2, 3$) with $s_1 \perp s_2$, $s_1 \perp s_3$, $s_2 \perp s_3$, as a category of orthogonally additive mappings. A \mathbb{C} -linear orthogonally mappings (\perp -mappings) $\mathbb{k}_h, \mathbb{k}_d : \mathfrak{B} \rightarrow \mathfrak{B}$ are called the orthogonally triple Lie homomorphism and derivation, respectively, whenever

$$\mathbb{k}_h([s_1, s_2], s_3) = [\mathbb{k}_h(s_1), \mathbb{k}_h(s_2)], \mathbb{k}_h(s_3),$$

and

$$\begin{aligned} \mathbb{k}_d([s_1, s_2], s_3) &= [[\mathbb{k}_d(s_1), s_2], s_3] + [s_1, \mathbb{k}_d(s_2)], s_3 \\ &+ [s_1, s_2], \mathbb{k}_d(s_3). \end{aligned}$$

Definition 1.10. Consider an orthogonally triple Lie homomorphism $\mathbb{k}_h : \mathfrak{B} \rightarrow \mathfrak{B}$. The \mathbb{C} -linear \perp -mapping $\mathbb{k}_{hd} : \mathfrak{B} \rightarrow \mathfrak{B}$ is called an orthogonally triple Lie hom-derivations, when

$$\begin{aligned} \mathbb{k}_{hd}([s_1, s_2], s_3) &= [[\mathbb{k}_{hd}(s_1), \mathbb{k}_h(s_2)], \mathbb{k}_h(s_3)] \\ &+ [[\mathbb{k}_h(s_1), \mathbb{k}_{hd}(s_2)], \mathbb{k}_h(s_3)] \\ &+ [[\mathbb{k}_h(s_1), \mathbb{k}_h(s_2)], \mathbb{k}_{hd}(s_3)], \end{aligned}$$

for each $s_i \in \mathfrak{B}$, $i = 1, 2, 3$.

Further, we establish the stability of HU sense for the orthogonally Jensen s -FĒ and orthogonally triple Lie hom-derivations among orthogonally triple Lie algebras through the application of the next key Theorem 1.11.

Theorem 1.11 ([6]). *In O -complete metric space (\mathfrak{B}, d, \perp) , consider a \perp -preserving, \perp -continuous and \perp -contraction $\mathcal{J}_{sf} : \mathfrak{B} \rightarrow \mathfrak{B}$ with Lipschitz constant $0 < \gamma < 1$. Assume that for fixed $s_o \in \mathfrak{B}$ we have either $s_o \perp s$ or $s \perp s_circ$, for all $s \in \mathfrak{B}$ and consider the “ O -sequence of successive approximations with initial element s_o ”:*

$$s_o, \mathcal{J}_{sf}(s_o), \mathcal{J}_{sf}^2(s_o), \dots, \mathcal{J}_{sf}^n(s_o), \dots$$

Then, either

$$d(\mathcal{J}_{sf}^n(s_o), \mathcal{J}_{sf}^{n+1}(s_o)) = \infty, \quad \forall n \geq 0,$$

or there is $n_0 \in \mathbb{N}$ such that, $d(\mathcal{J}_{sf}^n(s_o), \mathcal{J}_{sf}^{n+1}(s_o)) < \infty$ for every $n > n_0$. If the second alternative holds, then

- i) The O -sequence of $\{\mathcal{J}_{sf}^n(s_o)\}$ is convergent to a F.P, s^* of \mathcal{J}_{sf} .
- ii) s^* is the unique F.P of \mathcal{J}_{sf} in

$$\mathfrak{B}^* = \left\{ s \in \mathfrak{B} : d(\mathcal{J}_{sf}^n(s_o), s) < \infty \right\}.$$

- iii) If $s^* \in \mathfrak{B}$, then $d(s, s^*) \leq \frac{1}{1-\gamma} d(s, \mathcal{J}_{sf}(s))$.

2 Main Results

In this section, let $s \neq 0, \pm 1$, $\gamma \in \mathbb{T}^1 := \{\gamma \in \mathbb{C} : |\gamma| = 1\}$ and \mathfrak{B} be a orthogonally triple Lie algebra. For simplicity of work, we define

$$V_{\perp} := \left\{ s_1, s_2, s_3 \in \mathfrak{B} : s_1 \perp s_2, s_1 \perp s_3, s_2 \perp s_3 \right\}.$$

In the sequel, we investigate the orthogonally Jensen s -FĒ which is an additive mapping and stability of HU for the triple Lie hom-derivations with preserving orthogonality. In Subsection 2.2, our focus shifts to examining the hyperstability of FĒ (2) and the triple Lie hom-derivations with preserving orthogonality such that, this investigation is conducted through Găvruta’s control function, which serves as a pivotal tool in our analysis.

2.1 Stability of the Triple Lie Hom-Derivations

Before delving into the proof of the main theorem in this section, it is imperative to establish the additivity of the \perp -mapping \mathcal{J}_{sf} .

Proposition 2.1. *If the proposed FE (2) holds true for each $s_i \in V_\perp$, $i = 1, 2, 3$, then the \perp -mapping $\mathcal{J}_{sf} : \mathfrak{B} \rightarrow \mathfrak{B}$ can be characterized as additive with \perp -preserving.*

Proof. Suppose that \mathcal{J}_{sf} fulfills condition FE (2). Setting $s_i = 0$, $i = 1, 2, 3$, in FE (2), we get $\mathcal{J}_{sf}(0) = 0$. Substituting $s_2 = s_3 = 0$ into (2), we get,

$$3\mathcal{J}_{sf}(s_1) = \mathcal{J}_{sf}(3s_1). \quad (3)$$

By using (3) and putting $s_1 = s_2 = s_3$ in FE (2), we have

$$2\mathcal{J}_{sf}(s_1) = \mathcal{J}_{sf}(2s_1).$$

Again putting $s_2 = -s_1$ and $s_3 = 0$, we get $\mathcal{J}_{sf}(-s_1) = -\mathcal{J}_{sf}(s_1)$. Finally, replace $s_3 = 0$ in FE (2) and employ (3), we have

$$\mathcal{J}_{sf}(s_1 + s_2) = \mathcal{J}_{sf}(s_1) + \mathcal{J}_{sf}(s_2).$$

Thus $\mathcal{J}_{sf} : \mathfrak{B} \rightarrow \mathfrak{B}$ is an \perp -mapping with additive property. \square

In the following proposition, we investigate \mathcal{J}_{sf} is an \mathbb{C} -linear \perp -mapping.

Proposition 2.2. *If a \perp -mapping $\mathcal{J}_{sf} : \mathfrak{B} \rightarrow \mathfrak{B}$ satisfies*

$$\begin{aligned} 3\mathcal{J}_{sf}\left(\frac{\gamma[s_1+s_2+s_3]}{3}\right) - \gamma\mathcal{J}_{sf}(s_1) - \gamma\mathcal{J}_{sf}(s_2) - \mathbb{k}_d(\mathcal{J}_{sf}(s_3)) \\ = s\left[\mathcal{J}_{sf}(\gamma[s_1+s_2+s_3]) + \gamma\mathcal{J}_{sf}(s_1) \right. \\ \left. - \gamma\mathcal{J}_{sf}(s_1+s_2) - \gamma\mathcal{J}_{sf}(s_1+s_3)\right], \end{aligned} \quad (4)$$

for every $s_i \in V_\perp$, $i = 1, 2, 3$ and $\gamma \in \mathbb{T}^1$, then the \perp -mapping $\mathcal{J}_{sf} : \mathfrak{B} \rightarrow \mathfrak{B}$ is an orthogonally \mathbb{C} -linear.

Proof. By Proposition 2.1 \mathcal{J}_{sf} is an orthogonally additive. Considering $s_2 = s_3 = 0$ in FE (4), we get $\mathcal{J}_{sf}(\gamma s_1) = \gamma\mathcal{J}_{sf}(s_1)$. Following the same

logic as in the proof of [43, Theorem 2.1], the mapping \mathcal{J}_{sf} is established to be \mathbb{C} -linear \perp -mapping. \square

Theorem 2.3 discusses regarding stability of orthogonally Jensen s -FE with Găvruta's control function on orthogonally triple Lie algebras approach orthogonally F.P theorem. Consider $\mathbb{k}_d, \psi : \mathfrak{B}^3 \rightarrow \mathbb{R}^{\geq 0}$ such that for some $0 < \lambda < 1$,

$$\mathbb{k}_d\left(\frac{s_1}{3}, \frac{s_2}{3}, \frac{s_3}{3}\right) \leq \frac{\lambda}{3} \mathbb{k}_d(s_1, s_2, s_3), \quad (5)$$

and

$$\psi\left(\frac{s_1}{3}, \frac{s_2}{3}, \frac{s_3}{3}\right) \leq \frac{\lambda}{3^3} \psi(s_1, s_2, s_3), \quad (6)$$

for each $s_i \in V_\perp$, $i = 1, 2, 3$. When setting $s_i = 0$, $i = 1, 2, 3$, we find that $\delta(0, 0, 0) = 0$ and $\psi(0, 0, 0) = 0$. From inequalities (5) and (6), it follows that,

$$\lim_{j \rightarrow \infty} 3^j \mathbb{k}_d\left(\frac{s_1}{3^j}, \frac{s_2}{3^j}, \frac{s_3}{3^j}\right) = 0, \quad \lim_{j \rightarrow \infty} 3^{3j} \psi\left(\frac{s_1}{3^j}, \frac{s_2}{3^j}, \frac{s_3}{3^j}\right) = 0. \quad (7)$$

In the upcoming theorem, we establish the stability of UH for FE (2) by employing the F.P theorem with \perp -preserving.

Theorem 2.3. *Let function $\mathbb{k}_d : \mathfrak{B}^3 \rightarrow \mathbb{R}^{\geq 0}$ satisfies in inequality (5) and $\mathcal{J}_{sf} : \mathfrak{B} \rightarrow \mathfrak{B}$ be a mapping such that,*

$$\begin{aligned} & \left\| 3 \mathcal{J}_{sf}\left(\frac{\gamma[s_1 + s_2 + s_3]}{3}\right) - \gamma \mathcal{J}_{sf}(s_1) - \gamma \mathcal{J}_{sf}(s_2) - \gamma \mathcal{J}_{sf}(s_3) \right. \\ & \quad - s \left[\mathcal{J}_{sf}(\gamma[s_1 + s_2 + s_3]) + \gamma \mathcal{J}_{sf}(s_1) \right. \\ & \quad \left. \left. - \gamma \mathcal{J}_{sf}[s_1 + s_2] - \gamma \mathcal{J}_{sf}[s_1 + s_3] \right] \right\| \\ & \leq \mathbb{k}_d(s_1, s_2, s_3), \end{aligned} \quad (8)$$

for every $s_i \in V_\perp$, $i = 1, 2, 3$ and $\gamma \in \mathbb{T}^1$. Then under these conditions, there exists a unique \mathbb{C} -linear \perp -mapping $\mathcal{T} : \mathfrak{B} \rightarrow \mathfrak{B}$ with \perp -preserving, for which

$$\|\mathcal{J}_{sf}(s_1) - \mathcal{T}(s_1)\| \leq \frac{1}{1-\lambda} \mathbb{k}_d(s_1, 0, 0). \quad (9)$$

Proof. Letting $\gamma = 1$ and $s_2 = s_3 = 0$ in (8), we get

$$\left\| 3 \mathcal{J}_{sf}\left(\frac{s_1}{3}\right) - \mathcal{J}_{sf}(s_1) \right\| \leq \mathbb{k}_d(s_1, 0, 0). \quad (10)$$

Let Υ be the set of all mappings $\mathcal{J}_{sf} : \mathfrak{B} \rightarrow \mathfrak{B}$ such that, $\mathcal{J}_{sf}(0) = 0$ and

$$\mathcal{J}_{sf}(s_1) \perp 3\mathcal{J}_{sf}\left(\frac{s_1}{3}\right) \quad \text{or} \quad 3\mathcal{J}_{sf}\left(\frac{s_1}{3}\right) \perp \mathcal{J}_{sf}(s_1).$$

Now, for every $\mathcal{J}_{sf}, \acute{\mathcal{J}}_{sf} \in \Upsilon$, we consider the metric is expressed by,

$$\begin{aligned} d(\mathcal{J}_{sf}, \acute{\mathcal{J}}_{sf}) &= \inf \left\{ \lambda \in (0, \infty) : \|\mathcal{J}_{sf}(s_1) - \acute{\mathcal{J}}_{sf}(s_1)\| \right. \\ &\quad \left. \leq \lambda \mathbb{k}_d(s_1, 0, 0), \forall s_1 \in V_\perp \right\}, \end{aligned}$$

and define the orthogonality relation \perp within Υ for each $\mathcal{J}_{sf}, \acute{\mathcal{J}}_{sf} \in \Upsilon$ in the manner described below,

$$\acute{\mathcal{J}}_{sf} \perp \mathcal{J}_{sf} \Leftrightarrow \acute{\mathcal{J}}_{sf}(s_1) \perp \mathcal{J}_{sf}(s_1) \quad \text{or} \quad \mathcal{J}_{sf}(s_1) \perp \acute{\mathcal{J}}_{sf}(s_1),$$

for each $s_1 \in V_\perp$. Demonstrating that (Υ, d) constitutes an O -metric space is straightforward. Now, we consider the \mathbb{C} -linear \perp -mapping,

$$\begin{cases} \widehat{\Omega} : \Upsilon \rightarrow \Upsilon, \\ \widehat{\Omega}(\mathcal{J}_{sf}(s_1)) = 3\mathcal{J}_{sf}\left(\frac{s_1}{3}\right). \end{cases}$$

$\widehat{\Omega}$ maintains orthogonality. Therefore, for any $\mathcal{J}_{sf}, \acute{\mathcal{J}}_{sf} \in \Upsilon$ with $\mathcal{J}_{sf} \perp \acute{\mathcal{J}}_{sf}$ and it is given that $d(\acute{\mathcal{J}}_{sf}, \mathcal{J}_{sf}) = \varepsilon$, then,

$$\|\mathcal{J}_{sf}(s_1) - \acute{\mathcal{J}}_{sf}(s_1)\| \leq \varepsilon \mathbb{k}_d(s_1, 0, 0), \quad \forall s_1 \in V_\perp.$$

So

$$\begin{aligned} \|\widehat{\Omega}(\mathcal{J}_{sf}(s_1)) - \widehat{\Omega}(\acute{\mathcal{J}}_{sf}(s_1))\| &= \left\| 3\mathcal{J}_{sf}\left(\frac{s_1}{3}\right) - 3\acute{\mathcal{J}}_{sf}\left(\frac{s_1}{3}\right) \right\| \\ &\leq 3\varepsilon \mathbb{k}_d\left(\frac{s_1}{3}, 0, 0\right) \leq 3\varepsilon \frac{\lambda}{3} \mathbb{k}_d(s_1, 0, 0) = \lambda \varepsilon \mathbb{k}_d(s_1, 0, 0), \end{aligned}$$

and

$$d(\widehat{\Omega}(\mathcal{J}_{sf}), \widehat{\Omega}(\acute{\mathcal{J}}_{sf})) \leq \lambda \varepsilon, \quad \forall s_1 \in V_\perp.$$

This means that

$$d(\widehat{\Omega}(\mathcal{J}_{sf}), \widehat{\Omega}(\acute{\mathcal{J}}_{sf})) \leq \lambda d(\mathcal{J}_{sf}, \acute{\mathcal{J}}_{sf}),$$

and so Ω acts as a strictly \perp -contractive self-mapping on Υ , characterized by a Lipschitz constant λ . Moreover, $\widehat{\Omega}$ exhibits \perp -continuity. Specifically, if $\{\mathcal{J}_{sf}^j\}$ represents an O -sequence within Υ converging to

some $\mathcal{J}_{sf} \in \Upsilon$, then given any $\varepsilon > 0$, there is a $\lambda > 0$, where $\lambda < \varepsilon$, and an integer $\mathring{j} \in \mathbb{N}$ such that,

$$\left\| \mathcal{J}_{sf}^{\mathring{j}}(s_1) - \mathcal{J}_{sf}(s_1) \right\| \leq \lambda \mathbb{k}_d(s_1, 0, 0).$$

According to Theorem 1.11, a mapping $\mathcal{T} : \mathfrak{B} \rightarrow \mathfrak{B}$ exists that meets the criteria below.

I) \mathcal{T} serves as a F.P for $\widehat{\Omega}$, meaning,

$$\mathcal{T}(s_1) = 3\mathcal{T}\left(\frac{s_1}{3}\right), \quad \forall s_1 \in V_{\perp}. \quad (11)$$

The mapping \mathcal{T} represents a singular F.P of Ω . This denotes the existence of a unique \perp -mapping that fulfills (11), with a certain $\lambda > 0$ meeting the condition,

$$\|\mathcal{J}_{sf}(s_1) - \mathcal{T}(s_1)\| \leq \lambda \mathbb{k}_d(s_1, 0, 0), \quad \forall s_1 \in V_{\perp}.$$

II) $d(\lambda^{\mathring{j}} \mathcal{J}_{sf}, \mathcal{T}) \rightarrow 0$ as $\mathring{j} \rightarrow \infty$. This implies,

$$\lim_{\mathring{j} \rightarrow \infty} 3^{\mathring{j}} \mathcal{J}_{sf}\left(\frac{s_1}{3^{\mathring{j}}}\right) = \mathcal{T}(s_1), \quad \forall s_1 \in V_{\perp}.$$

III) $d(\mathcal{J}_{sf}, \mathcal{T}) \leq \frac{1}{1-\lambda} d(\mathcal{J}_{sf}, \widehat{\Omega}(\mathcal{T}))$, which implies,

$$\|\mathcal{J}_{sf}(s_1) - \mathcal{T}(s_1)\| \leq \frac{1}{1-\lambda} \mathbb{k}_d(s_1, 0, 0), \quad \forall s_1 \in V_{\perp}.$$

It follows from (7) and (8) that,

$$\begin{aligned} & \left\| 3\mathcal{T}\left(\frac{s_1+s_2+s_3}{3}\right) - \mathcal{T}(s_1) - \mathcal{T}(s_2) - \mathcal{T}(s_3) \right. \\ & \quad \left. - s \left[\mathcal{T}(s_1 + s_2 + s_3) - \mathcal{T}(s_1) \right. \right. \\ & \quad \left. \left. - \mathcal{T}(s_1 + s_2) - \mathcal{T}(s_1 + s_3) \right] \right\| \\ &= \lim_{\mathring{j} \rightarrow \infty} 3^{\mathring{j}} \left\| 3\mathcal{J}_{sf}\left(\frac{s_1+s_2+s_3}{3^{\mathring{j}+1}}\right) - \mathcal{J}_{sf}\left(\frac{s_1}{3^{\mathring{j}}}\right) - \mathcal{J}_{sf}\left(\frac{s_2}{3^{\mathring{j}}}\right) - \mathcal{J}_{sf}\left(\frac{s_3}{3^{\mathring{j}}}\right) \right. \\ & \quad \left. - s \left[\mathcal{J}_{sf}\left(\frac{s_1+s_2+s_3}{3^{\mathring{j}}}\right) + \mathcal{J}_{sf}\left(\frac{s_1}{3^{\mathring{j}}}\right) \right. \right. \\ & \quad \left. \left. - \mathcal{J}_{sf}\left(\frac{s_1+s_2}{3^{\mathring{j}}}\right) - \mathcal{J}_{sf}\left(\frac{s_1+s_3}{3^{\mathring{j}}}\right) \right] \right\| \\ &\leq \lim_{\mathring{j} \rightarrow \infty} 3^{\mathring{j}} \mathbb{k}_d\left(\frac{s_1}{3^{\mathring{j}}}, \frac{s_2}{3^{\mathring{j}}}, \frac{s_3}{3^{\mathring{j}}}\right) = 0, \quad \forall s_i \in V_{\perp}, i = 1, 2, 3. \end{aligned}$$

So

$$\begin{aligned} 3\mathcal{T}\left(\frac{s_1+s_2+s_3}{3}\right) - \mathcal{T}(s_1) - \mathcal{T}(s_2) - \mathcal{T}(s_3) \\ = s \left[\mathcal{T}(s_1 + s_2 + s_3) + \mathcal{T}(s_1) \right. \\ \left. - \mathcal{T}(s_1 + s_2) - \mathcal{T}(s_1 + s_3) \right], \end{aligned}$$

for $s_i \in V_\perp$, $i = 1, 2, 3$.

By proposition 2.2, $\mathcal{T} : \mathfrak{B} \rightarrow \mathfrak{B}$ is an \mathbb{C} -linear \perp -mapping. \square

In the next theorem, we apply Găvruta's control function to establish the stability of triple Lie hom-derivations while \perp -preserving through the orthogonal F.P theorem approach.

Theorem 2.4. *Let $\psi : \mathfrak{B}^3 \rightarrow \mathbb{R}^{\geq 0}$ be functions such that, there exists an $0 < \lambda < 1$ satisfying in inequality (6) and $\acute{\mathcal{J}}_{\text{sf}}, \mathcal{J}_{\text{sf}} : \mathfrak{B} \rightarrow \mathfrak{B}$ are mappings satisfying (8) and*

$$\left\| \acute{\mathcal{J}}_{\text{sf}}([s_1, s_2], s_3) - [[\acute{\mathcal{J}}_{\text{sf}}(s_1), \acute{\mathcal{J}}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3)] \right\| \leq \psi(s_1, s_2, s_3), \quad (12)$$

$$\begin{aligned} \left\| \mathcal{J}_{\text{sf}}([s_1, s_2], s_3) - [[\mathcal{J}_{\text{sf}}(s_1), \mathcal{J}_{\text{sf}}(s_2)], \mathcal{J}_{\text{sf}}(s_3)] \right. \\ \left. - [[\acute{\mathcal{J}}_{\text{sf}}(s_1), \mathcal{J}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3)] \right. \\ \left. - [[\mathcal{J}_{\text{sf}}(s_1), \acute{\mathcal{J}}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3)] \right\| \leq \psi(s_1, s_2, s_3), \quad (13) \end{aligned}$$

for $s_i \in V_\perp$, $i = 1, 2, 3$. Under these conditions, there exist unique orthogonally triple Lie homomorphism, hom-derivation $\mathbb{k}_h, \mathbb{k}_{\text{hd}} : \mathfrak{B} \rightarrow \mathfrak{B}$, respectively, which

$$\begin{aligned} \left\| \acute{\mathcal{J}}_{\text{sf}}(s_1) - \mathbb{k}_h(s_1) \right\| &\leq \frac{1}{1-\lambda} \psi(s_1, 0, 0), \\ \left\| \mathcal{J}_{\text{sf}}(s_1) - \mathbb{k}_{\text{hd}}(s_1) \right\| &\leq \frac{1}{1-\lambda} \psi(s_1, 0, 0). \end{aligned} \quad (14)$$

Proof. Following the same logic as in the proof of Theorem 2.3, for $s_1 \in V_\perp$, can define the mappings,

$$\begin{cases} \mathbb{k}_h : \mathfrak{B} \rightarrow \mathfrak{B}, \\ \mathbb{k}_h(s_1) = \lim_{j \rightarrow \infty} 3^j \acute{\mathcal{J}}_{\text{sf}}\left(\frac{s_1}{3^j}\right), \end{cases} \quad \begin{cases} \mathbb{k}_{\text{hd}} : \mathfrak{B} \rightarrow \mathfrak{B}, \\ \mathbb{k}_{\text{hd}}(s_1) = \lim_{j \rightarrow \infty} 3^j \mathcal{J}_{\text{sf}}\left(\frac{s_1}{3^j}\right). \end{cases} \quad (15)$$

Eqs. (12) and (15) imply,

$$\begin{aligned}
& \left\| \mathbb{k}_h([s_1, s_2], s_3) - [\mathbb{k}_h(s_1), \mathbb{k}_h(s_2)], \mathbb{k}_h(s_3) \right\| \\
&= \lim_{j \rightarrow \infty} 3^{3j} \left\| \mathcal{J}_{sf} \left(\left[\left[\frac{s_1}{3^j}, \frac{s_2}{3^j} \right], \frac{s_3}{3^j} \right] \right) \right. \\
&\quad \left. - \left[\left[\mathcal{J}_{sf} \left(\frac{s_1}{3^j} \right), \mathcal{J}_{sf} \left(\frac{s_2}{3^j} \right) \right], \mathcal{J}_{sf} \left(\frac{s_3}{3^j} \right) \right] \right\| \\
&\leq \lim_{j \rightarrow \infty} 3^{3j} \psi \left(\frac{s_1}{3^j}, \frac{s_2}{3^j}, \frac{s_3}{3^j} \right) = 0.
\end{aligned}$$

Hence, the mapping \mathbb{k}_h is a triple Lie homomorphism with \perp -preserving. It follows from (13) and (15), we have

$$\begin{aligned}
& \left\| \mathbb{k}_{hd}([s_1, s_2], s_3) - [\mathbb{k}_{hd}(s_1), s_2], s_3 \right\| + \left\| [s_1, \mathbb{k}_{hd}(s_2)], s_3 \right\| \\
&\quad + \left\| [s_1, s_2], \mathbb{k}_{hd}(s_3) \right\| \\
&= \lim_{j \rightarrow \infty} 3^{3j} \left\| \mathcal{J}_{sf} \left(\left[\left[\frac{s_1}{3^j}, \frac{s_2}{3^j} \right], \frac{s_3}{3^j} \right] \right) \right. \\
&\quad - \left[\left[\mathcal{J}_{sf} \left(\frac{s_1}{3^j} \right), \mathcal{J}_{sf} \left(\frac{s_2}{3^j} \right) \right], \mathcal{J}_{sf} \left(\frac{s_3}{3^j} \right) \right] \\
&\quad - \left[\left[\mathcal{J}_{sf} \left(\frac{s_1}{3^j} \right), \mathcal{J}_{sf} \left(\frac{s_2}{3^j} \right) \right], \mathcal{J}_{sf} \left(\frac{s_3}{3^j} \right) \right] \\
&\quad \left. - \left[\left[\mathcal{J}_{sf} \left(\frac{s_1}{3^j} \right), \mathcal{J}_{sf} \left(\frac{s_2}{3^j} \right) \right], \mathcal{J}_{sf} \left(\frac{s_3}{3^j} \right) \right] \right\| \\
&\leq \lim_{j \rightarrow \infty} 3^{3j} \psi \left(\frac{s_1}{3^j}, \frac{s_2}{3^j}, \frac{s_3}{3^j} \right) = 0.
\end{aligned}$$

Hence, the mapping $\mathbb{k}_{hd} : \mathfrak{B} \rightarrow \mathfrak{B}$ is an orthogonally triple Lie hom-derivations. \square

Theorem 2.3 generalized the result of Rassias' theorem, by the following definition

$$\mathbb{k}_d(s_1, s_2, s_3) := \theta \left[\|s_1\|^r + \|s_2\|^r + \|s_3\|^r \right], \quad \forall \theta, r \in \mathbb{R}^{>0}, r \neq 1.$$

We have the following corollary.

Corollary 2.5. *Let $r \neq 1$ and θ be nonnegative real numbers and $\mathcal{J}_{sf} :$*

$\mathfrak{B} \rightarrow \mathfrak{B}$ be a mapping satisfying $\mathcal{J}_{\text{sf}}(0) = 0$ and

$$\begin{aligned} & \left\| 3\mathcal{J}_{\text{sf}}\left(\frac{\gamma[s_1+s_2+s_3]}{3}\right) - \gamma\mathcal{J}_{\text{sf}}(s_1) - \gamma\mathcal{J}_{\text{sf}}(s_2) - \mathbb{k}_{\text{d}}(\mathcal{J}_{\text{sf}}(s_3)) \right. \\ & \quad - s\left[\mathcal{J}_{\text{sf}}(\gamma[s_1+s_2+s_3]) + \gamma\mathcal{J}_{\text{sf}}(s_1) \right. \\ & \quad \left. \left. - \gamma\mathcal{J}_{\text{sf}}(s_1+s_2) - \gamma\mathcal{J}_{\text{sf}}(s_1+s_3)\right] \right\| \\ & \leq \theta\left[\|s_1\|^r + \|s_2\|^r + \|s_3\|^r\right], \end{aligned}$$

for each $s_i \in V_{\perp}$, $i = 1, 2, 3$, $\gamma \in \mathbb{T}^1$. Then, under these condition, there exists a unique \mathbb{C} -linear \perp -mapping $\mathcal{T} : \mathfrak{B} \rightarrow \mathfrak{B}$ with \perp -preserving, such that

$$\|\mathcal{J}_{\text{sf}}(s_1) - \mathcal{T}(s_1)\| \leq \begin{cases} \frac{2\theta}{2^r-2}\|s_1\|^r, & r > 1, \\ \frac{2\theta}{2-2^r}\|s_1\|^r, & r < 1, \end{cases} \quad (16)$$

for $s_i \in V_{\perp}$, $i = 1, 2, 3$.

Proof. By putting $\lambda = 2^{r-1}$, $r > 1$ and $\lambda = 2^{1-r}$, $r < 1$ in relation (16), the desired results are obtained. \square

In Theorem 2.4, by taking $\lambda = 2^{r-1}$ and $\lambda = 2^{1-r}$, whenever $r > 1$, $r < 1$, respectively and

$$\psi(s_1, s_2, s_3) := \theta[\|s_1\|^r + \|s_2\|^r + \|s_3\|^r], \quad \forall s_i \in V_{\perp}, i = 1, 2, 3,$$

where r and θ are nonnegative real numbers, we obtain the following result.

Corollary 2.6. Consider two elements $r \neq 1$ and θ of $\mathbb{R}^{>0}$, let

$$\psi(s_1, s_2, s_3) = \theta[\|s_1\|^r + \|s_2\|^r + \|s_3\|^r], \quad (17)$$

and assume $\mathcal{J}_{\text{sf}}, \mathcal{J}_{\text{sf}} : \mathfrak{B} \rightarrow \mathfrak{B}$, are functions satisfying (8), (12) and (13). Then there exist orthogonally triple Lie homomorphism \mathbb{k}_{h} and hom-derivations \mathbb{k}_{hd} such that,

$$\begin{cases} \|\mathcal{J}_{\text{sf}}(s_1) - \mathbb{k}_{\text{h}}(s_1)\| \leq \frac{2\theta}{2^r-2}\|s_1\|^r, \\ \|\mathcal{J}_{\text{sf}}(s_1) - \mathbb{k}_{\text{hd}}(s_1)\| \leq \frac{2\theta}{2-2^r}\|s_1\|^r, \end{cases}$$

and

$$\begin{cases} \|\mathcal{J}_{\text{sf}}(s_1) - \mathbb{k}_{\text{h}}(s_1)\| \leq \frac{2\theta}{2-2^r} \|s_1\|^r, \\ \|\mathcal{J}_{\text{sf}}(s_1) - \mathbb{k}_{\text{hd}}(s_1)\| \leq \frac{2\theta}{2-2^r} \|s_1\|^r, \end{cases}$$

for $r > 1$ and $r < 1$, respectively.

2.2 On the Hyperstability of Orthogonally Triple Lie Hom-Derivations

The initial result on hyperstability was originally presented in [9] and focused on particular ring homomorphisms. The concept of hyperstability refers to a functional equation that is considered hyperstable if every approximate solution of the equation is, in fact, an exact solution. On the other hand, the Ulam stability problem questions the existence of an exact solution. For further details on hyperstability, see [26, 31].

In this subsection, by leveraging the orthogonally F.P technique, we investigate hyperstability of FE (2).

Theorem 2.7. *Let there is $\psi : \mathfrak{B}^3 \rightarrow \mathbb{R}^{\geq 0}$ such that,*

$$\lim_{j \rightarrow \infty} \frac{1}{3^j} \mathbb{k}_{\text{d}}(0, 3^j s_2, 3^j s_3) = 0. \quad (18)$$

Moreover, suppose that $\mathcal{J}_{\text{sf}} : \mathfrak{B} \rightarrow \mathfrak{B}$ is mapping such that,

$$\begin{aligned} & \left\| 3 \mathcal{J}_{\text{sf}} \left(\frac{\gamma[s_1 + s_2 + s_3]}{3} \right) - \gamma \mathcal{J}_{\text{sf}}(s_1) - \gamma \mathcal{J}_{\text{sf}}(s_2) - \mathbb{k}_{\text{d}}(\mathcal{J}_{\text{sf}}(s_3)) \right. \\ & \quad - s \left[\mathcal{J}_{\text{sf}}(\gamma[s_1 + s_2 + s_3]) + \gamma \mathcal{J}_{\text{sf}}(s_1) \right. \\ & \quad \left. \left. - \gamma \mathcal{J}_{\text{sf}}(s_1 + s_2) - \gamma \mathcal{J}_{\text{sf}}(s_1 + s_3) \right] \right\| \\ & \leq \mathbb{k}_{\text{d}}(0, s_2, s_3). \end{aligned} \quad (19)$$

Under these conditions, it is a \mathbb{C} -linear \perp -mapping $\mathcal{T} : \mathfrak{B} \rightarrow \mathfrak{B}$ with \perp -preserving.

Proof. Letting $\gamma = 1$ and $s_2 = s_3 = 0$ in (19), we get $\mathcal{J}_{\text{sf}}(3s_1) = 3\mathcal{J}_{\text{sf}}(s_1)$ and by using induction on $j \in \mathbb{N}$ we get,

$$\mathcal{J}_{\text{sf}}(s_1) = 3^j \mathcal{J}_{\text{sf}} \left(\frac{s_1}{3^j} \right).$$

Hence, we have

$$\begin{aligned} 3^j \left\| 3 \mathcal{J}_{\text{sf}} \left(\frac{s_1 + s_2 + s_3}{3^{j+1}} \right) - \mathcal{J}_{\text{sf}} \left(\frac{s_1}{3^j} \right) - \mathcal{J}_{\text{sf}} \left(\frac{s_2}{3^j} \right) - \mathcal{J}_{\text{sf}} \left(\frac{s_3}{3^j} \right) \right. \\ \left. - s \left[\mathcal{J}_{\text{sf}} \left(\frac{s_1 + s_2 + s_3}{3^j} \right) + \mathcal{J}_{\text{sf}} \left(\frac{s_1}{3^j} \right) \right. \right. \\ \left. \left. - \mathcal{J}_{\text{sf}} \left(\frac{s_1 + s_2}{3^j} \right) - \mathcal{J}_{\text{sf}} \left(\frac{s_1 + s_3}{3^j} \right) \right] \right\| \\ \leq 3^i \mathbb{k}_d \left(0, \frac{s_2}{3^j}, \frac{s_3}{3^j} \right) = 0, \end{aligned}$$

for every $s_i \in V_\perp$, $i = 1, 2, 3$, $j \in \mathbb{N}$. So, by $j \rightarrow \infty$ and using (18), we get

$$\begin{aligned} 3 \mathcal{J}_{\text{sf}} \left(\frac{(s_1 + s_2 + s_3)}{3} \right) - \mathcal{J}_{\text{sf}}(s_1) - \mathcal{J}_{\text{sf}}(s_2) - \mathbb{k}_d(\mathcal{J}_{\text{sf}}(s_3)) \\ - s \left[\mathcal{J}_{\text{sf}}(s_1 + s_2 + s_3) + \mathcal{J}_{\text{sf}}(s_1) \right. \\ \left. - \mathcal{J}_{\text{sf}}(s_1 + s_2) - \mathcal{J}_{\text{sf}}(s_1 + s_3) \right] = 0, \end{aligned}$$

for $s_i \in V_\perp$, $i = 1, 2, 3$. Therefore, \mathcal{J}_{sf} is an \mathbb{C} -linear \perp -mapping. \square

In the subsequent theorem, our focus shifts to examining the hyperstability of orthogonally triple Lie hom-derivation. This investigation is conducted through Găvruta's control function, which serves as a pivotal tool in our analysis. By leveraging the orthogonally F.P technique, we delve into the intricacies of the equation's hyperstability properties.

Theorem 2.8. *Suppose there exist function $\psi : \mathfrak{B}^3 \rightarrow \mathbb{R}^{\geq 0}$ such that,*

$$\lim_{j \rightarrow \infty} \frac{1}{3^{3j}} \psi \left(0, 3^j s_2, 3^j s_3 \right) = 0, \quad \forall s_i \in V_\perp, i = 1, 2, 3.$$

Moreover, suppose that $\acute{\mathcal{J}}_{\text{sf}}, \mathcal{J}_{\text{sf}} : \mathfrak{B} \rightarrow \mathfrak{B}$ are mappings satisfying (19) and

$$\begin{aligned} \left\| \acute{\mathcal{J}}_{\text{sf}}([s_1, s_2], s_3) - [\zeta(s_1), \acute{\mathcal{J}}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3) \right\| &\leq \psi(0, s_2, s_3), \\ \left\| \mathcal{J}_{\text{sf}}([s_1, s_2], s_3) - [\mathcal{J}_{\text{sf}}(s_1), \acute{\mathcal{J}}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3) \right. \\ &\quad \left. - [\acute{\mathcal{J}}_{\text{sf}}(s_1), \mathcal{J}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3) \right. \\ &\quad \left. - [\acute{\mathcal{J}}_{\text{sf}}(s_1), \acute{\mathcal{J}}_{\text{sf}}(s_2)], \mathcal{J}_{\text{sf}}(s_3) \right\| &\leq \psi(0, s_2, s_3), \end{aligned}$$

for $s_i \in V_\perp$, $i = 1, 2, 3$. Then $\acute{\mathcal{J}}_{\text{sf}}$ and \mathcal{J}_{sf} are orthogonally triple Lie homomorphism and orthogonally triple Lie hom-derivation, respectively.

Proof. The methodology employed in the proof bears resemblance to that of Theorem 2.7, showcasing a parallel structure in the analytical approach. \square

In the ensuing corollary, our focus shifts towards exploring the hyperstability of $\text{F}\mathbb{E}$ (2) through control function of Rassias with \perp -preserving.

Corollary 2.9. Let $\theta, r \in \mathbb{R}^{>0}$ with $r \neq 1$ and $\mathcal{J}_{\text{sf}} : \mathfrak{B} \rightarrow \mathfrak{B}$ is mapping such that,

$$\begin{aligned} & \left\| 3\mathcal{J}_{\text{sf}}\left(\frac{\gamma[s_1+s_2+s_3]}{3}\right) - \gamma\mathcal{J}_{\text{sf}}(s_1) - \gamma\mathcal{J}_{\text{sf}}(s_2) - \gamma\mathbb{k}_d(\mathcal{J}_{\text{sf}}(s_3)) \right. \\ & \quad \left. - s\left[\mathcal{J}_{\text{sf}}(\gamma[s_1+s_2+s_3]) + \gamma\mathcal{J}_{\text{sf}}(s_1) \right. \right. \\ & \quad \left. \left. - \gamma\mathcal{J}_{\text{sf}}(s_1+s_2) - \gamma\mathcal{J}_{\text{sf}}(s_1+s_3)\right] \right\| \\ & \leq \theta\left[\|s_2\|^r + \|s_3\|^r\right]. \end{aligned}$$

Under these conditions, it is a \mathbb{C} -linear \perp -mapping $\mathcal{T} : \mathfrak{B} \rightarrow \mathfrak{B}$ with \perp -preserving.

Now, in the following corollary, our focus shifts towards exploring the hyperstability of orthogonally triple Lie hom-derivation by control function of Rassias.

Corollary 2.10. Let $\theta, r \in \mathbb{R}^{>0}$ with $r \neq 1$ and $\acute{\mathcal{J}}_{\text{sf}}, \mathcal{J}_{\text{sf}} : \mathfrak{B} \rightarrow \mathfrak{B}$ are mappings such that,

$$\left\| \acute{\mathcal{J}}_{\text{sf}}([s_1, s_2], s_3) - [\acute{\mathcal{J}}_{\text{sf}}(s_1), \acute{\mathcal{J}}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3) \right\| \leq \theta\left[\|s_2\|^r + \|s_3\|^r\right],$$

and

$$\begin{aligned} & \left\| \mathcal{J}_{\text{sf}}([s_1, s_2], s_3) - [\mathcal{J}_{\text{sf}}(s_1), \mathcal{J}_{\text{sf}}(s_2)], \mathcal{J}_{\text{sf}}(s_3) \right\| \\ & \quad - [\mathcal{J}_{\text{sf}}(s_1), \mathcal{J}_{\text{sf}}(s_2)], \acute{\mathcal{J}}_{\text{sf}}(s_3) \\ & \quad - [\acute{\mathcal{J}}_{\text{sf}}(s_1), \acute{\mathcal{J}}_{\text{sf}}(s_2)], \mathcal{J}_{\text{sf}}(s_3) \right\| \\ & \leq \theta\left[\|s_2\|^r + \|s_3\|^r\right]. \end{aligned}$$

Then $\acute{\mathcal{J}}_{\text{sf}}$ and \mathcal{J}_{sf} are orthogonally triple Lie homomorphism and orthogonally triple Lie hom-derivation, respectively.

3 Conclusion

Given the significance of orthogonal sets, triple Lie and Banach algebras, triple Lie derivations, and their practical applications in various fields such as engineering and mathematical physics, mathematical finance and economics. We introduced the new concept of orthogonally Jensen equation and orthogonally triple Lie hom-derivations on orthogonally triple Lie algebras. After that, we solved some examples of the concept of O -sets that can be O -set and \perp -contraction.

Finally, through the utilization of the F.P method with \perp -preserving, we have successfully demonstrated that orthogonally triple Lie hom-derivations and the FE (2) exhibit stability and hyperstability properties when applied to orthogonally triple Lie algebras. As the orthogonally Jensen s -functional equation, on triple Lie algebras while \perp -preserving, are applicable in different fields of science, so one can extend different models and analytic techniques by means of the generalized Jensen s -functional equation for various science processes in the next research studies [4, 5, 16, 46, 47].

References

- [1] M. Abbas, R. Anjum and S. Riasat, A new type of fixed point theorem via interpolation of operators with application in homotopy theory, *Arabian Journal of Mathematics* 12 (2023), 277–288. <https://doi.org/10.1007/s40065-022-00402-z>
- [2] M. Amyari and M. S. Moslehian, Approximately ternary semigroup homomorphisms, *Letters in Mathematical Physics*, 77 (2006), 1–9. <https://doi.org/10.1007/s11005-005-0042-6>
- [3] A. Aruchamy, S. Rayappan and A. Natarajan, Global existence and stability results for a time-fractional diffusion equation with variable exponents, *Arabian Journal of Mathematics*, 13 (2024), 237–254. <https://doi.org/10.1007/s40065-024-00463-2>
- [4] S. M. Aydogan, To investigate a class of the singular fractional integro-differential quantum equations with multi-step methods,

- Journal of Mathematical Extention*, 15(SI-NTFCA, 10) (2021) , 1–18. <https://doi.org/10.30495/JME.SI.2021.2065>
- [5] S. M. Aydogan, A. Hussain and F. M. Sakar, On a nonlinear fractional order model of novel coronavirus (COVID-2019) under AB-fractional derivative, *Journal of Mathematical Extention*, 15(SI-NTFCA, 11) (2021), 1–31. <https://doi.org/10.30495/JME.SI.2021.1994>
- [6] A. Bahraimi, G. Askari, M. Eshaghi Gordji and R. Gholami, Stability and hyperstability of orthogonally \ast - m -homomorphisms in orthogonally Lie C^\ast -algebras: a fixed point approach, *Journal of Fixed Point Theory and Applications*, 20 (2018), 89. <https://doi.org/10.1007/s11784-018-0571-0>
- [7] J. Baker, The stability of the cosine equation, *Proceedings of the American Mathematical Society*, 80(3) (1979), 242–246. <https://doi.org/10.1155/2012/984160>
- [8] T. Björk and C. Landén, On the construction of finite dimensional realizations for nonlinear forward rate models, *Finance and Stochastics*, 6 (2002), 303–331. <https://doi.org/10.1007/s007800100060>
- [9] D. G. Bourgin, Approximately isometric and multiplicative transformations on continuous function rings, *Duke Mathematical Journal*, 16(2) (1949), 385–397. <https://doi.org/10.1215/S0012-7094-49-01639-7>
- [10] S. Chandok, L. Paunović and S. Radenović, Incomplete metric with the application for integral equations of Volterra type, *Mathematical Problems in Engineering*, 2022 (2022), 11 pages. <https://doi.org/10.1155/2022/5959903>
- [11] K. Das and M. E. Samei, Using ρ -cone arcwise connectedness on parametric set-valued optimization problems, *Journal of Inequalities and Applications*, 2022 (2022), 57. <https://doi.org/10.1186/s13660-022-02792-2>

- [12] J. B. Diaz and B. Margolis, A fixed point theorem of the alternative for contractions on the generalized complete metric space, *Bulletin of the American Mathematical Society*, 74(3) (1968), 305–309. <https://doi.org/10.1090/s0002-9904-1968-11933-0>
- [13] E. Elqorachi and M. T. Rassias, Generalized Hyers-Ulam stability of trigonometric functional equations, *Mathematics*, 6(5) (2018), 83. <https://doi.org/10.3390/math6050083>
- [14] M. Eshaghi Gordji, A. Ebadian, N. Ghobadipour, J. M. Rassias and M. B. Savadkouhi, Approximately ternary homomorphisms and derivations on C^* -ternary algebras, *Abstract and Applied Analysis*, 2012 (2012), 10 pages. <https://doi.org/10.1155/2012/984160>
- [15] M. Eshaghi Gordji, M. Rameani, M. De La Sen and Y. J. Cho, On orthogonal sets and Banach fixed point theorems, *Fixed Point Theory*, 18(2) (2017), 569–578. <https://doi.org/10.24193/fpt-ro.2017.2.45>
- [16] R. George, S. M. Aydogan, F. M. Sakar, M. Ghaderi and S. Rezapour, A study on the existence of numerical and analytical solutions for fractional integrodifferential equations in hilfer type with simulation, *AIMS Mathematics*, 8(5) (2023), 665–684. <https://doi.org/10.3934/math.2023541>
- [17] P. Găvruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, *Journal of Mathematical Analysis and Applications*, 184(3) (1994), 431–436. <https://doi.org/10.1006/jmaa.1994.1211>
- [18] Z. Heydarpour, J. Izadi and K. Ghafarnezhad, Twisted (α, β) -contractive mapping with preserving orthogonality and HU stability, *Journal of Mathematical Extension*, 18(2) (2024), (6)1–18. <https://doi.org/10.30495/JME.2024.3107>
- [19] M. Houas and M. E. Samei, Solvability and ULAM-HYERS-RASSIAS stability for generalized sequential quantum fractional pantograph equations, *Partial Differential Equations in Applied Mathematics*, 9 (2024), 100651. <https://doi.org/10.1016/j.padiff.2024.100651>

- [20] D. H. Hyers, On the stability of the linear functional equation, *Proceedings of the National Academy of Sciences of the United States of America*, 27 (1941), 222–224.
- [21] F. Iachello, *Lie Algebras and Applications*, Springer, Berlin, Heidelberg (2015).
- [22] J. Izadi, Orthogonally generalized lie bracket of triple derivations, *Journal of Inequalities and Special Functions*, 15(4) (2024), 15–25. <https://doi.org/10.54379/jiasf-2024-4-2>
- [23] S. Jahedi and V. Keshavarz, Approximate generalized additive-quadratic functional equations on ternary Banach algebras, *Journal of Mathematical Extension*, 16(1) (2022), (1)1–11. <https://doi.org/10.30495/JME.2022.1859>
- [24] B. Jayawardana, P. J. Morrison and T. Ohsawa, Clebsch canonization of Lie-Poisson systems, *Journal of Geometric Mechanics*, 14(4) (2022), 635–658. <https://doi.org/10.3934/jgm.2022017>
- [25] M. Kapranov, M. Gelfand and A. Zelevinskii, *Discriminants, Resultants and Multidimensional Determinants*, Springer, Berlin Birkhauser (1994).
- [26] Z. Kefayati, T. Abdeljawad, A. Mukheimer and M. A. Alqudah, Orthogonal generalized Lie bracket of triple derivations: properties and stability, *Journal of Inequalities and Applications*, 2025 (2025), 69. <https://doi.org/10.1186/s13660-025-03324-4>
- [27] R. Kerner, The cubic chessboard, *Classical Quantum Gravity*, 14(1A) (1997), A203–A225. <https://doi.org/10.1088/0264-9381/14/1A/017>
- [28] V. Keshavarz, M. T. Heydari and D. R. Anderson, Hyers-Ulam stabilities for m th differential operators on H^2_β , *Chaos, Solitons & Fractals*, 179 (2024), 114443. <https://doi.org/10.1016/j.chaos.2023.114443>
- [29] V. Keshavarz and M. T. Heydari, Hyers-Ulam stability of solution for generalized Lie bracket of derivations, *Journal of Mathematics*, 2024 (2024), 8 pages. <https://doi.org/10.1155/2024/1015443>

- [30] V. Keshavarz and S. Jahedi, Orthogonally C^* -ternary Jordan homomorphisms and Jordan derivations: solution and stability, *Journal of Mathematics*, 2022 (2022), 7 pages. <https://doi.org/10.1155/2022/3482254>
- [31] V. Keshavarz, S. Jahedi, M. Eshaghi Gordji and S. Bazeghi, Hyperstability of orthogonally Pexider Lie functional equation: an orthogonally fixed point approach, *Thai Journal of Mathematics*, 20 (2022), 235–243.
- [32] S. Lie and G. Scheffers, *Vorlesungen über Kontinuerliche Gruppen*, Springer, Leipzig (1893).
- [33] C. F. Lo and C. H. Hui, Pricing multi-asset financial derivatives with timedependent parameters-Lie algebraic approach, *International Journal of Mathematics and Mathematical Sciences*, 32 (2002), 10 pages. <https://doi.org/10.1155/S016117120211101X>
- [34] C. F. Lo, C. H. Hui and P. H. Yuen, Constant elasticity of variance option pricing model with time-dependent parameters, *International Journal of Theoretical and Applied Finance*, 03(04) (2000), 661–674. <https://doi.org/10.1142/S0219024900000814>
- [35] J. L. Martin, Generalized classical dynamics, and the 'classical analogue' of a Fermi oscillator, *Proceedings of the Royal Society A: Mathematical. Physical and Engineering Sciences*, 251 (1959), 536–542. <https://doi.org/10.1098/rspa.1959.0126>
- [36] M. M. Matar, M. E. Samei, S. Etemad, A. Amara, S. Rezapour and J. Alzabut, Stability analysis and existence criteria with numerical illustrations to fractional jerk differential system involving generalized Caputo derivative, *Qualitative Theory of Dynamical Systems*, 23 (2024), 111. <https://doi.org/10.1007/s12346-024-00970-9>
- [37] Y. Nambu, Generalized Hamiltonian dynamics, *Physical Review D*, 7 (1973), 2405–2412. <https://doi.org/10.1103/PhysRevD.7.2405>
- [38] M. Nazam, H. Aydi and A. Hussain, Existence theorems for (ψ, ϕ) -orthogonal interpolative contractions and an application to frac-

- tional differential equations, *Optimization: A Journal of Mathematical Programming and Operations Research*, 72(7) (2023), 1899–1929. <https://doi.org/10.1080/02331934.2022.2043858>
- [39] M. Nazarianpoor, J. M. Rassias and G. Sadeghi, Stability and nonstability of octadecic functional equation in multi-normed spaces, *Arabian Journal of Mathematics*, 7 (2018), 219–228. <https://doi.org/10.1007/s40065-017-0186-0>
- [40] M. Obloza, Hyers stability of the linear differential equation, *Rocznik Nauk. Dydakt. Prace Mat.*, 13 (1993), 259–270.
- [41] S. Okubo, Triple products and Yang-Baxter equation (I): octonionic and quaternionic triple systems, *Journal of Mathematical Physics*, 34(7) (1993), 3273–3291. <https://doi.org/10.1063/1.530076>
- [42] C. Park, J. R. Lee and X. Zhang, Additive s -functional inequality and hom-derivations in Banach algebras, *Journal of Fixed Point Theory and Applications*, 21 (2019), 18. <https://doi.org/10.1007/s11784-018-0652-0>
- [43] C. G. Park, Homomorphisms between poisson JC^* -algebras, *Bulletin of the Brazilian Mathematical Society*, 36 (2005), 79–97. <https://doi.org/10.1007/s00574-005-0029-z>
- [44] S. Popov and J. M. Strelcyn, The Euler-poisson equations: An elementary approach to integrability conditions, *Journal of Geometric Mechanics*, 10(3) (2018), 293–329. <https://doi.org/10.3934/jgm.2018011>
- [45] T. M. Rassias, On the stability of the linear mapping in Banach spaces, *Proceedings of the American Mathematical Society*, 72(2) (1978), 297–300. <https://doi.org/10.2307/2042795>
- [46] S. Rezapour, F. M. Sakar, S. M. Aydogan and E. Ravash, Some results on a system of multiterm fractional integro-differential equations, *Turkish Journal of Mathematics*, 44(6) (2020), 2004–2020. <https://doi.org/10.3906/mat-1903-51>

- [47] M. E. Samei, H. Zanganeh and S. M. Aydogan, To investigate a class of the singular fractional integro-differential quantum equations with multi-step methods, *Journal of Mathematical Extention*, 15(SI-NTFCA, 15) (2021), 1–54.
<https://doi.org/10.30495/JME.SI.2021.2070>
- [48] S. M. Ulam, *Problems in Modern Mathematics*, Chapter IV, Science edn. Wiley, New York (1960).

Javad Izadi

Assistant Professor

Department of Mathematics

Department of Mathematics, Payame Noor University (PNU)

P. O. Box 19395-4697, Tehran, Iran

E-mail: j.izadi@pnu.ac.ir

Mohammad Esmael Samei

Associate Professor of Mathematics

Department of Mathematics

Faculty of Science, Bu-Ali Sina University

Hamedan, Iran

E-mail: mesamei@basu.ac.ir, mesamei@gmail.com