

Journal of Mathematical Extension
 Vol. 19, No. 6 (2025) (5) 1-13
 ISSN: 1735-8299
 URL: <http://doi.org/10.30495/JME.2025.3353>
 Original Research Paper

Generalized Volterra-Type Operators from Hardy Space into Iterated Weighted-Type Spaces

E. Abbasi

Islamic Azad University, Mahabad, Iran

D. Molaei *

Islamic Azad University, Urmia, Iran

A. Ebrahimi

Islamic Azad University, Mahabad, Iran

Abstract. Let $H(\mathbb{D})$ be the set of analytic functions on \mathbb{D} and for $1 \leq p \leq \infty$, H^p be the Hardy space. For $m \in \mathbb{N}$ suppose that I^m be m th iteration. Let $\vec{g} = (g_0, \dots, g_{m-1})$ where $\{g_i\}_{i=0}^{m-1} \subset H(\mathbb{D})$ and $I(f) = \int_0^z f(w)dw$. If I^m for $m \in \mathbb{N}$ be the m th iteration, then the generalized Volterra-type operators $I_{\vec{g}}^m$ on $H(\mathbb{D})$ is defined as follows

$$I_{\vec{g}}^m(f) = I^m \left(\sum_{i=0}^{m-1} f^{(i)} g_i \right).$$

In this paper, we investigate boundedness and compactness of generalized Volterra-type operators from Hardy space into iterated weighted-type spaces, $V_n = \{f \in H(\mathbb{D}) : \sup_{z \in \mathbb{D}} (1 - |z|^2) |f^{(n)}(z)| < \infty\}$.

AMS Subject Classification: 47G10; 30H10; 30H05; 30H99

Keywords and Phrases: Boundedness, Compactness, Generalized Volterra-type operators, Hardy space, Iterated weighted-type spaces

Received: May 2025; Accepted: November 2025

*Corresponding Author

1 Introduction

Let \mathbb{D} be an open unit disc in the complex \mathbb{C} and $H(\mathbb{D})$ be the set of analytic functions on \mathbb{D} . For $1 \leq p < \infty$, the Hardy space H^p consists of all analytic functions $f \in H(\mathbb{D})$ such that

$$\|f\|_{H^p} = \sup_{0 < r < 1} \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{\frac{1}{p}} < \infty.$$

Also H^∞ is the space of bounded analytic functions on \mathbb{D} with the norm $\|f\|_{H^\infty} = \sup_{z \in \mathbb{D}} |f(z)|$. More information about such spaces can be found in [9].

Another space used in this paper is n th weighted-type space. Let μ be a weight (continuous and positive function on \mathbb{D}) and $n \in \mathbb{N}_0$. The n th weighted-type space V_n^μ , consists of all analytic functions $f \in H(\mathbb{D})$ such that $b_{V_n^\mu}(f) = \sup_{z \in \mathbb{D}} \mu(z) |f^{(n)}(z)| < \infty$. This space is a Banach with the following norm

$$\|f\|_{V_n^\mu} = \sum_{i=0}^{n-1} |f^{(i)}(0)| + b_{V_n^\mu}(f) < \infty.$$

For $\alpha > 0$ and $\mu(z) = (1 - |z|^2)^\alpha$, we use V_n^α , V_n and $\|\cdot\|_n$ instead of V_n^μ , V_n^1 and $\|\cdot\|_{V_n^1}$. The space V_n^α contains a large class of analytic functions. For example when $\alpha > 0$, $V_0^\alpha = \mathcal{A}^{-\alpha}$ (growth space), $V_1^\alpha = \mathcal{B}^\alpha$ (Bloch type space), $V_2^\alpha = \mathcal{Z}^\alpha$ (Zygmund type space), $V_1 = \mathcal{B}$ (classic Bloch space) and $V_2 = \mathcal{Z}$ (classic Zygmund space). The space V_n is called iterated weighted-type space. In [7] Colonna *et al.* considered iterated weighted-type spaces and obtained some properties for these spaces, especially they showed

$$\cdots \subset V_{n+1} \subset V_n \subset \cdots \subset V_3 \subset \mathcal{Z} \subset H^\infty \subset \mathcal{B} \subset \mathcal{A}^{-1}.$$

The closed subspace of V_n^μ containing of all $f \in V_n^\mu$ such that $\lim_{|z| \rightarrow 1} \mu(z) |f^{(n)}(z)| = 0$ is denoted by $V_{n,0}^\mu$ and is called little n th weighted-type space. For more information about (little) n th weighted-type spaces, see [1, 2, 4, 7, 12, 14].

Let $m \in \mathbb{N}$, $I^m(f) = \int_0^z \int_0^{z_1} \cdots \int_0^{z_{m-1}} f(z) dz dz_1 \cdots dz_{m-1}$, and $\vec{g} = (g_0, g_1, \dots, g_{m-1})$ where $\{g_i\}_{i=0}^{m-1} \subset H(\mathbb{D})$. The generalized Volterra-type operator on $H(\mathbb{D})$ defined as follows

$$I_{\vec{g}}^m(f) = I^m\left(\sum_{i=0}^{m-1} f^{(i)} g_i\right).$$

For $m = 1$ and $g_0 = g'$, we get Volterra type operator $(J_g f)(z) = \int_0^z f(w) g'(w) dw$ and when $m = 1$ and $g_0 = 1$, we obtain the classic Volterra operator $(If)(z) = \int_0^z f(w) dw$. Also if we set $g_i = a_{m-i-1} g^{(m-i)}$ ($0 \leq i \leq m-1$), where $g \in H(\mathbb{D})$ and $\vec{a} = (a_0, \dots, a_{m-1}) \in \mathbb{C}^m$, we have generalized integration operator $I_{g, \vec{a}}^m$ defined by Chalmoukis in [6].

Since

$$I_{\vec{g}}^m(f) = I^m\left(\sum_{i=0}^{m-1} f^{(i)} g_i\right) = \sum_{i=0}^{m-1} I^m(f^{(i)} g_i) := \sum_{i=0}^{m-1} I_{g_i}^{m,i}(f),$$

so for considering properties of operator $I_{\vec{g}}^m$, firstly we investigate properties of operators $I_{g_i}^{m,i}$ where $0 \leq i \leq m-1$.

Chalmoukis in [6] considered boundedness and compactness of $I_{g, \vec{a}}^m : H^p \rightarrow H^q$, where $(0 < q < p < \infty)$ and posed a conjecture that g must be in $H^{\frac{pq}{q-p}}$. Yang *et al.* provided a positive answer to the aforementioned conjecture in [13]. Arroussi *et al.* investigated boundedness and compactness of $I_{\vec{g}}^m : A^p \rightarrow A^q$, where A^p is Bergman space. They extended Chalmoukis' result to Bergman spaces and showed that the Bergman space version of Chalmoukis' conjecture is true (see [5]). Also some authors characterized boundedness and compactness of generalized integration operators among some other analytic function spaces [8, 10]. In [14], Zhu investigated Bloch-type spaces and uncovered numerous properties associated with these spaces. Later, Stević expanded on this concept by generalizing Bloch-type spaces and introducing the n th weighted-type spaces, as detailed in [11, 12]. In recent years, extensive research has been conducted on such spaces, with one of the most notable references in this area being [7].

In this paper, firstly we investigate boundedness and compactness of the operators $I_{g_i}^{m,i}$ ($0 \leq i \leq m$) from Hardy space into iterated weighted-type spaces and we find some characterizations for boundedness and

compactness of such operators. Then we consider boundedness and compactness of the operator $I_{\overline{g}}^m : H^p \rightarrow V_n$ and we show that the operator $I_{\overline{g}}^m : H^p \rightarrow V_n$ is bounded (compact) if and only if each operator $I_{g_i}^{m,i} : H^p \rightarrow V_n (0 \leq i \leq m-1)$ is bounded (compact).

In this work, we shall use the notation $A \preceq B$ to mean that for some $c > 0$, $A \leq cB$, whereas $A \asymp B$ means $A \preceq B$ and $B \preceq A$.

2 Boundedness and Compactness of Operator

$$I_{g_i}^{m,i} : H^p \rightarrow V_n$$

In this section, we investigate boundedness and compactness of the operators $I_{g_i}^{m,i}$ from Hardy space $H^p (1 \leq p \leq \infty)$ into iterated weighted-type spaces and we obtain some characterizations for boundedness and compactness of such operators. We begin with the following lemma.

Lemma 2.1. *Let $n, k \in \mathbb{N}$. For any $f \in V_n$, we have*

$$\|f\|_n \asymp \sum_{i=0}^{n+k-2} |f^{(i)}(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^k |f^{(n+k-1)}(z)|.$$

Proof. For $n = 1$, $V_1 = \mathcal{B}$. So by using Proposition 8 in [14], we get

$$\|f\|_1 = \|f\|_{\mathcal{B}} \asymp \sum_{i=0}^{k-1} |f^{(i)}(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^k |f^{(k)}(z)|.$$

For any $n \in \mathbb{N}$, $V_n \subset V_{n-1}$ [6, Proposition 2.1]. Hence, for any $f \in V_n$, $f^{(n-1)} \in \mathcal{B}$. By replacing f with $f^{(n-1)}$ in the above equation, we obtain

$$\begin{aligned} & |f^{(n-1)}(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^k |f^{(n)}(z)| \asymp \\ & \sum_{i=n-1}^{n+k-2} |f^{(i)}(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^k |f^{(n+k-1)}(z)|. \end{aligned}$$

Hence,

$$\begin{aligned}
\|f\|_n &= \sum_{i=0}^{n-2} |f^{(i)}(0)| + |f^{(n-1)}(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2) |f^{(n)}(z)| \\
&\asymp \sum_{i=0}^{n-2} |f^{(i)}(0)| + \sum_{i=n-1}^{n+k-2} |f^{(i)}(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^k |f^{(n+k-1)}(z)| \\
&= \sum_{i=0}^{n+k-2} |f^{(i)}(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^k |f^{(n+k-1)}(z)|.
\end{aligned}$$

The proof is complete. \square

Let $m \geq n$. It is clear $\left(I_g^m(f)\right)^{(i)}(0) = \left(I_g^{m,i}(f)\right)^{(i)}(0) = 0$, when $0 \leq i \leq m-1$, so for any $f \in H(\mathbb{D})$, by using Lemma 2.1, we have

$$\begin{aligned}
\|I_g^{m,i}(f)\|_n &\asymp \sum_{k=0}^{m-1} \left| \left(I_g^{m,i}(f)\right)^{(k)}(0) \right| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^{m-n+1} \left| f^{(i)} g(z) \right| \quad (1) \\
&= \sup_{z \in \mathbb{D}} (1 - |z|^2)^{m-n+1} \left| f^{(i)} g(z) \right|.
\end{aligned}$$

Lemma 2.2. *Let $1 \leq p \leq \infty$. Then for any $f \in H^p$ and $k \in \mathbb{N}_0$,*

$$\sup_{z \in \mathbb{D}} (1 - |z|^2)^{k+\frac{1}{p}} |f^{(k)}(z)| \leq \|f\|_{H^p}.$$

Proof. From Proposition 5.1.2 of [14], we have $H^\infty \subset \mathcal{B}$ and for any $f \in H^\infty$

$$\sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| \leq \|f\|_{H^\infty}.$$

Applying Lemma 2.1, for each $k \in \mathbb{N}_0$ and $f \in H^\infty$, we obtain

$$\sup_{z \in \mathbb{D}} (1 - |z|^2)^k |f^{(k)}(z)| \leq \|f\|_1 \leq 2\|f\|_{H^\infty} \leq \|f\|_{H^\infty}.$$

Similar results for $1 \leq p < \infty$ follow easily using results from [9]. \square

Theorem 2.3. *Let $1 \leq p \leq \infty$, $m, n \in \mathbb{N}$ such that $m \geq n$ and $g_i \in H(\mathbb{D})$. Then for each $0 \leq i \leq m$, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is bounded if and only if $\sup_{z \in \mathbb{D}} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| < \infty$. Moreover, in this case*

$$\|I_{g_i}^{m,i}\| \asymp \sup_{z \in \mathbb{D}} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)|.$$

Proof. Let the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ be bounded and $f_{i,w}(z) = \frac{(1-|w|^2)^i}{(1-\bar{w}z)^{i+\frac{1}{p}}}$, where $0 \leq i \leq m$ and $w \in \mathbb{D} - \{0\}$. For $p = \infty$,

$$\sup_{z \in \mathbb{D}} |f_{i,w}(z)| = \sup_{z \in \mathbb{D}} \left| \frac{1-|w|^2}{1-\bar{w}z} \right|^i \leq \sup_{z \in \mathbb{D}} \frac{(1-|w|^2)^i}{(1-|\bar{w}|)^i} = 2^i,$$

so, $\sup_{w \in \mathbb{D}} \|f_{i,w}\|_{H^\infty} \leq 2^i$ and when $1 \leq p < \infty$, from Lemma 2 [11], there exists positive constant $C_{i,p}$ such that $\sup_{w \in \mathbb{D}} \|f_{i,w}\|_{H^p} < C_{i,p}$. Applying (1) for $f_{i,w}$, we have

$$\begin{aligned} \|I_{g_i}^{m,i}(f_{i,w})\|_n &\asymp \sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n+1} |f_{i,w}^{(i)}(z)g_i(z)| \\ &\geq (1-|w|^2)^{m-n+1} |f_{i,w}^{(i)}(w)| |g_i(w)| \\ &= |\bar{w}|^i \prod_{l=0}^{i-1} \left(i + \frac{1}{p} + l \right) (1-|w|^2)^{m-n-i-\frac{1}{p}+1} |g_i(w)|. \end{aligned}$$

Therefore,

$$\begin{aligned} \sup_{w \in \mathbb{D}} (1-|w|^2)^{m-n-i-\frac{1}{p}+1} |g_i(w)| &\preceq \|I_{g_i}^{m,i}(f_{i,w})\|_n \leq \|I_{g_i}^{m,i}\| \sup_{w \in \mathbb{D}} \|f_{i,w}\|_{H^p} \\ &\leq C_{i,p} \|I_{g_i}^{m,i}\|. \end{aligned}$$

Conversely, we assume that $\sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| < \infty$, by using (1), Lemmas 2.1 and 2.2, for any $f \in H^p$, we obtain

$$\begin{aligned} \|I_{g_i}^{m,i}(f)\|_n &\asymp \sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n+1} |f^{(i)}(z)g_i(z)| \\ &\leq \sup_{z \in \mathbb{D}} (1-|z|^2)^{i+\frac{1}{p}} |f^{(i)}(z)| \sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| \\ &\preceq \|f\|_{H^p} \sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)|. \end{aligned}$$

Therefore, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is bounded and $\|I_{g_i}^{m,i}\| \preceq \sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)|$. The proof is complete. \square

To investigate compactness of operators $I_g^{m,i} : H^p \rightarrow V_n$ ($0 \leq i \leq m$), we need the following lemma, since the proof of it is similar to the proof of [7, proposition 3.11], so it is omitted.

Lemma 2.4. *Let $1 \leq p \leq \infty$, $m, n \in \mathbb{N}$, $\{g_i\}_{i=0}^{m-1} \subset H(\mathbb{D})$ and $T = I_{\vec{g}}^m$ or $I_{g_i}^{m,i}$ ($0 \leq i \leq m$). Then the bounded operator $T : H^p \rightarrow V_n$ is compact if and only if for any bounded sequence $\{f_k\}$ in H^p which converges to zero uniformly on compact subsets of \mathbb{D} , $\lim_{k \rightarrow \infty} \|T(f_k)\|_n = 0$.*

Theorem 2.5. *Let $1 \leq p \leq \infty$, $m, n \in \mathbb{N}$ such that $m \geq n$ and $g_i \in H(\mathbb{D})$ ($0 \leq i \leq m$). Then for each $0 \leq i \leq m$, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is compact if and only if $\lim_{|z| \rightarrow 1} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| = 0$.*

Proof. Let the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ be compact. For any $0 \leq i \leq m$ and $w \in \mathbb{D} - \{0\}$ the functions $f_{i,w}(z) = \frac{(1-|w|^2)^i}{(1-\bar{w}z)^{i+\frac{1}{p}}}$ are bounded and converge uniformly to zero on compact subsets of \mathbb{D} when $|w|$ tends to 1, so by applying Lemma 2.4 $\lim_{|w| \rightarrow 1} \|I_{g_i}^{m,i}(f_{i,w})\|_n = 0$. Now by using (1), we obtain

$$\begin{aligned} \|I_{g_i}^{m,i}(f_{i,w})\|_n &\asymp (1 - |z|^2)^{m-n+1} |f_{i,w}^{(i)}(z)| |g_i(z)| \\ &\succeq (1 - |w|^2)^{m-n-i-\frac{1}{p}+1} |\bar{w}|^i |g_i(w)|. \end{aligned}$$

In the above inequality, let $|w| \rightarrow 1$. Then, we obtain $\lim_{|w| \rightarrow 1} (1 - |w|^2)^{m-n-i-\frac{1}{p}+1} |g_i(w)| = 0$.

Conversely, suppose that $\lim_{|z| \rightarrow 1} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| = 0$. Hence, for any $\varepsilon > 0$ there exists $0 < \delta < 1$ such that for each $\delta < |z| < 1$,

$$(1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| < \varepsilon. \quad (2)$$

Now by using (1) and Lemma 2.1, for any bounded sequence $\{f_k\} \subset H^p$

which converges uniformly to zero on compact subsets of \mathbb{D} , we obtain

$$\begin{aligned}
\|I_{g_i}^{m,i}(f_k)\|_n &\asymp \sup_{z \in \mathbb{D}} (1 - |z|^2)^{m-n+1} |f_k^{(i)}(z)| |g_i(z)| \\
&\leq \sup_{|z| \leq \delta} (1 - |z|^2)^{m-n+1} |f_k^{(i)}(z)| |g_i(z)| \\
&\quad + \sup_{\delta < |z| < 1} (1 - |z|^2)^{m-n+1} |f_k^{(i)}(z)| |g_i(z)| \\
&\leq \sup_{|z| \leq \delta} (1 - |z|^2)^{i+\frac{1}{p}} |f_k^{(i)}(z)| \sup_{|z| \leq \delta} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| \\
&\quad + \sup_{\delta < |z| < 1} (1 - |z|^2)^{i+\frac{1}{p}} |f_k^{(i)}(z)| \sup_{\delta < |z| < 1} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| \\
&=: X_1 + X_2.
\end{aligned}$$

By using Cauchy's estimates, for any $i \in \mathbb{N}_0$, the sequence $\{f_k^{(i)}\}$ converges uniformly to zero on compact subsets of \mathbb{D} , therefore

$$\lim_{k \rightarrow \infty} X_1 \leq \sup_{|z| \leq \delta} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| \lim_{k \rightarrow \infty} \sup_{|z| \leq \delta} |f_k^{(i)}(z)| = 0.$$

Also applying Lemma 2.2 and (2), we get

$$\begin{aligned}
\lim_{k \rightarrow \infty} X_2 &\preceq \lim_{k \rightarrow \infty} \|f_k\|_{H^p} \sup_{\delta < |z| < 1} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| \\
&\leq \varepsilon \sup_{k \in \mathbb{N}} \|f_k\|_{H^p}.
\end{aligned}$$

So, $\lim_{k \rightarrow \infty} \|I_{g_i}^{m,i}(f_k)\|_n = 0$. By using Lemmas 2.4, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is compact. The proof is complete. \square

Putting $n = 1$ and $n = 2$ in Theorems 2.3 and 2.5, we get the following corollaries.

Corollary 2.6. *Let $1 \leq p \leq \infty$, $m \in \mathbb{N}$ and $g_i \in H(\mathbb{D})$. Then for each $0 \leq i \leq m$, the operator $I_{g_i}^{m,i} : H^p \rightarrow \mathcal{B}$ is bounded (compact) if and only if $g_i \in V_0^{m-i-\frac{1}{p}}$ ($g_i \in V_{0,0}^{m-i-\frac{1}{p}}$).*

Corollary 2.7. *Let $1 \leq p \leq \infty$, $m \in \mathbb{N}$ such that $m \geq 2$ and $g_i \in H(\mathbb{D})$. Then for each $0 \leq i \leq m$, the operator $I_{g_i}^{m,i} : H^p \rightarrow \mathcal{Z}$ is bounded (compact) if and only if $g_i \in V_0^{m-i-\frac{1}{p}-1}$ ($g_i \in V_{0,0}^{m-i-\frac{1}{p}-1}$).*

3 Boundedness and Compactness of Operator

$$I_{\overrightarrow{g}}^m : H^p \rightarrow V_n$$

In this section, we will consider boundedness and compactness of the operator $I_{\overrightarrow{g}}^m$ from Hardy spaces into iterated weighted-type spaces. Especially we show that if the operator $I_{\overrightarrow{g}}^m : H^p \rightarrow V_n$ is bounded (compact) then for each $0 \leq i \leq m-1$, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is bounded (compact). For this purpose, we need the following lemma which comes from [2, Lemma 2.5] and [3, Lemma 2.3].

Lemma 3.1. *Let $1 \leq p \leq \infty$. For any $0 \neq \xi \in \mathbb{D}$ and $i \in \{0, 1, \dots, n\}$, there exists a function $v_{i,\xi} \in H^p$ with the following conditions:*

a) $v_{i,\xi}(z) = \sum_{j=1}^{n+1} c_j^i f_{i,\xi}(z)$, where $f_{i,\xi}(z) = \frac{(1-|\xi|^2)^i}{(1-\bar{\xi}z)^{i+\frac{1}{p}}}$ and c_j^i is independent of choice ξ .

b) $\sup_{\xi \in \mathbb{D}} \|v_{i,\xi}\|_{H^p} < \infty$ and

$$v_{i,\xi}^{(k)}(\xi) = \begin{cases} \frac{\bar{\xi}^i}{(1-|\xi|^2)^{i+\frac{1}{p}}}, & k = i, \\ 0, & k \neq i. \end{cases}$$

c) For any sequence $\{\xi_k\} \subset \mathbb{D}$ such that $\lim_{k \rightarrow \infty} |\xi_k| = 1$, the sequence $\{v_{i,\xi_k}\}$ converges to zero uniformly on compact subsets of \mathbb{D} .

Let $m \geq n$. For any $f \in H(\mathbb{D})$, applying Lemma 2.1, we get

$$\begin{aligned} \|I_{\overrightarrow{g}}^m(f)\|_n &\asymp \sum_{k=0}^{m-1} \left| \left(I_{\overrightarrow{g}}^m(f) \right)^{(k)}(0) \right| + \sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n+1} \left| \left(I_{\overrightarrow{g}}^m(f) \right)^{(m)} \right| \\ &= \sup_{z \in \mathbb{D}} (1-|z|^2)^{m-n+1} \left| \sum_{j=0}^{m-1} f^{(j)}(z) g_j(z) \right|. \end{aligned} \tag{3}$$

Theorem 3.2. *Let $1 \leq p \leq \infty$ and $m, n \in \mathbb{N}$ such that $m \geq n$. Then the following conditions are equivalent:*

- a) The operator $I_{\vec{g}}^m : H^p \rightarrow V_n$ is bounded.
- b) For each $0 \leq i \leq m-1$, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is bounded.
- c) For each $0 \leq i \leq m-1$, $\sup_{z \in \mathbb{D}} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| < \infty$.

Proof. (b) \Rightarrow (a) Since $I_{\vec{g}}^m = \sum_{i=0}^{m-1} I_{g_i}^{m,i}$ and all operators $I_{g_i}^{m,i} : H^p \rightarrow V_n$ ($0 \leq i \leq m-1$) are bounded, hence the operator $I_{\vec{g}}^m : H^p \rightarrow V_n$ is bounded.

(b) \Leftrightarrow (c) Theorem 2.3.

(a) \Rightarrow (c) Let the operator $I_{\vec{g}}^m : H^p \rightarrow V_n$ be bounded. For each $i \in \{0, \dots, m-1\}$ and $\xi \in \mathbb{D} - \{0\}$, let $v_{i,\xi}$ be function found in Lemma 3.1, by using (3), we have

$$\begin{aligned}
\|I_{\vec{g}}^m(v_{i,\xi})\|_{V_n} &\asymp \sup_{z \in \mathbb{D}} (1 - |z|^2)^{m-n+1} \left| \sum_{j=0}^{m-1} v_{i,\xi}^{(j)}(z) g_j(z) \right| \\
&\geq (1 - |\xi|^2)^{m-n+1} \left| \sum_{j=0}^{m-1} v_{i,\xi}^{(j)}(\xi) g_j(\xi) \right| \\
&\geq (1 - |\xi|^2)^{m-n+1} |v_{i,\xi}^{(s)}(\xi)| |g_i(\xi)| \\
&= (1 - |\xi|^2)^{m-n+1} \frac{|\bar{\xi}|^i}{(1 - |\xi|^2)^{i+\frac{1}{p}}} |g_i(\xi)| \\
&= |\bar{\xi}|^i (1 - |\xi|^2)^{m-n-i-\frac{1}{p}+1} |g_i(\xi)|.
\end{aligned} \tag{4}$$

Applying Lemma 3.1(b), we get

$$\sup_{\xi \in \mathbb{D}} (1 - |\xi|^2)^{m-n-i-\frac{1}{p}+1} |g_i(\xi)| \leq \|I_{\vec{g}}^m(v_{i,\xi})\|_{V_n} \leq \|I_{\vec{g}}^m\| \sup_{\xi \in \mathbb{D}} \|v_{i,\xi}\|_{H^p} < \infty.$$

The proof is complete. \square

Theorem 3.3. Let $1 \leq p \leq \infty$ and $m, n \in \mathbb{N}$ such that $m \geq n$. Then the following conditions are equivalent:

- a) The operator $I_{\vec{g}}^m : H^p \rightarrow V_n$ is compact.
- b) For each $0 \leq i \leq m-1$, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is compact.

c) For each $0 \leq i \leq m-1$, $\lim_{|z| \rightarrow 1} (1 - |z|^2)^{m-n-i-\frac{1}{p}+1} |g_i(z)| = 0$.

Proof. (b) \Rightarrow (a) It is clear when for each $0 \leq i \leq m-1$, the operator $I_{g_i}^{m,i} : H^p \rightarrow V_n$ is compact, then $I_{\vec{g}}^m = \sum_{i=0}^{m-1} I_{g_i}^{m,i}$ is compact.

(b) \Leftrightarrow (c) Theorem 2.5.

(a) \Rightarrow (c) Assume that the operator $I_{\vec{g}}^m : H^p \rightarrow V_n$ is compact. For each $i \in \{0, \dots, m-1\}$ and $\xi \in \mathbb{D} - \{0\}$, the sequence $\{v_{i,\xi}\}$ is bounded and converges to zero uniformly on compact subsets of \mathbb{D} when $|\xi|$ tends to 1 (Lemma 3.1), so by using Lemma 2.4, $\lim_{|\xi| \rightarrow 1} \|I_{\vec{g}}^m(v_{i,\xi})\|_n = 0$. Now it is enough to let $|\xi|$ tends to 1 in the inequality (4), therefore $\lim_{|\xi| \rightarrow 1} (1 - |\xi|^2)^{m-n-i-\frac{1}{p}+1} |g_i(\xi)| = 0$. The proof is complete. \square

Let $m \in \mathbb{N}$, $\bar{a} = (a_0, \dots, a_{m-1}) \in \mathbb{C}^m$ and $g \in H(\mathbb{D})$. Applying Theorems 3.2 and 3.3, we obtain similar results for the Chalmoukis operator

$$I_{g,\bar{a}}^m f(z) = I^m \left(fg^{(n)} + a_1 f' g^{(n-1)} + \dots + a_{n-1} f^{(n-1)} g' \right)$$

acting from the Hardy space into iterated weighted-type spaces.

Corollary 3.4. Let $1 \leq p \leq \infty$, $m \in \mathbb{N}$, $\bar{a} = (a_0, \dots, a_{m-1}) \in \mathbb{C}^m$ and $g \in H(\mathbb{D})$. Then

a) the operator $I_{g,\bar{a}}^m : H^p \rightarrow V_n$ is bounded if and only if

$$g \in \bigcap_{i=0}^{m-1} V_{m-i}^{m-n-i-\frac{1}{p}+1}.$$

b) the operator $I_{g,\bar{a}}^m : H^p \rightarrow V_n$ is compact if and only if

$$g \in \bigcap_{i=0}^{m-1} V_{m-i,0}^{m-n-i-\frac{1}{p}+1}.$$

Remark 3.5. By choosing suitable parameters m, p, n and \vec{g} , the results obtained in this paper, can be stated for some well-known operators and spaces.

References

- [1] E. Abbasi, The product-type operators from the Besov spaces into n th weighted type spaces. *Journal of Mathematical Extension*, 16(3): 1–14, 2022.
- [2] E. Abbasi, S. Li and H. Vaezi, Weighted composition operators from the Bloch space to n th weighted-type spaces. *Turk. J. Math.*, 44(1): 108–117, 2020.
- [3] E. Abbasi, Y. Liu and M. Hassanlou, Generalized Stević—Sharma type operators from Hardy spaces into n th weighted type spaces. *Turk. J. Math.*, 45(4): 1543–1554, 2021.
- [4] E. Abbasi, H. Vaezi and S. Li, Essential norm of weighted composition operators from H^∞ to n th weighted type spaces. *Mediterr. J. Math.*, 16: 133, 2019.
- [5] H. Arroussi, H. Liu, C. Tong and Z. Yang, A new class of Carleson measures and integral operators on Bergman spaces. *Bull. Sci. Math.*, 197: 103531, 2024.
- [6] N. Chalmoukis, Generalized integration operators on Hardy spaces. *Proc. Amer. Math. Soc.*, 148(8): 3325–3337, 2020.
- [7] F. Colonna and N. Hmidouch, Weighted composition operators on iterated weighted-type Banach spaces of analytic functions. *Complex Anal. Oper. Theory*, 13: 1989–2016, 2019.
- [8] J. Du, S. Li and D. Qu, The generalized Volterra integral operator and Toeplitz operator on weighted Bergman spaces. *Mediterr. J. Math.*, 19, 263, 2022.
- [9] P. L. Duren, *Theory of H^p Spaces*. Academic Press, New York and London, 1970.
- [10] C. Shen and S. Li, Volterra integral operators from Dirichlet type spaces into Hardy spaces. *Ric. mat.*, 72: 107–118, 2023.

- [11] S. Stević, Composition operators from the Hardy space to the n th weighted-type space on the unit disk and the half-plane. *Appl. Math. Comput.*, 215(11): 3950–3955, 2010.
- [12] S. Stević, Weighted differentiation composition operators from H^∞ and Bloch spaces to n th weighted type spaces on the unit disk. *Appl. Math. Comput.*, 216: 3634–3641, 2010.
- [13] R. Yang and S. Li, On a conjecture about generalized integration operators on Hardy spaces. arXiv preprint arXiv:2405.19372.
- [14] K. Zhu, Bloch type space of analytic functions. *Rocky Mountain J. Math.*, 23: 1143–1177, 1993.

Ebrahim Abbasi

Department of Mathematics
 Associate Professor of Mathematics
 Mah. C., Islamic Azad University
 Mahabad, Iran
 E-mail: ebrahimabbasi81@aut.ac.ir, ebrahimabbasi81@gmail.com

Daryoush Molaei

Department of Mathematics
 Assistant Professor of Mathematics
 Urm. C., Islamic Azad University
 Urmia, Iran
 E-mail: daryoush.molaei@iau.ac.ir

Ali Ebrahimi

Department of Mathematics
 Assistant Professor of Mathematics
 Mah. C., Islamic Azad University
 Mahabad, Iran
 E-mail: ebrahimi.ali2719@aut.ac.ir, ebrahimi.ali2719@gmail.com