

On the Weighted Hardy Spaces

K. Jahedi

Islamic Azad University-Shiraz Branch

B. Yousefi

Shiraz University

Abstract: Let $\{\beta(n)\}_{n=0}^{\infty}$ be a sequence of positive numbers and $1 < p < \infty$. We consider the weighted Hardy space $H^p(\beta)$. We investigate the relation between the generating function and the functional of point evaluations. Also, under a sufficient condition we determine the structure of all non-zero multiplicative linear functionals on $H^p(\beta)$.

AMS Subject Classification: Primary 47B37; Secondary 47B20.

Keywords and Phrases: The Banach space of formal power series associated with a sequence β , bounded point evaluation, generating function.

1. Introduction

First in the following, we generalize the definitions coming in [3]. Let

$\{\beta(n)\}$ be a sequence of positive numbers with $\beta(0) = 1$ and $1 < p < \infty$.

We consider the space of sequences $f = \{\hat{f}(n)\}_{n=0}^{\infty}$ such that

$$\|f\|^p = \|f\|_{\beta}^p = \sum_{n=0}^{\infty} |\hat{f}(n)|^p \beta(n)^p < \infty.$$

The notation $f(z) = \sum_{n=0}^{\infty} \hat{f}(n)z^n$ shall be used whether or not the series converges for any value of z . These are called formal power series. Let $H^p(\beta)$ denotes the space of such formal power series. It is usually called as weighted Hardy spaces. These are reflexive Banach spaces with the norm $\|\cdot\|_\beta$ and the dual of $H^p(\beta)$ is $H^q(\beta^{\frac{p}{q}})$ where $\frac{1}{p} + \frac{1}{q} = 1$ and $\beta^{\frac{p}{q}} = \{\beta(n)^{\frac{p}{q}}\}_n$ ([4]). Also if

$$g(z) = \sum_{n=0}^{\infty} \hat{g}(n)z^n \in H^q(\beta^{\frac{p}{q}}),$$

then

$$\|g\|^q = \sum_{n=0}^{\infty} |\hat{g}(n)|^q \beta(n)^p.$$

The Hardy, Bergman and Dirichlet spaces can be viewed in this way when $p = 2$ and respectively $\beta(n) = 1$, $\beta(n) = (n+1)^{-1/2}$ and $\beta(n) = (n+1)^{1/2}$. It is convenient and helpful to introduce the notation $\langle f, g \rangle$ to stand for $g(f)$ where $f \in H^p(\beta)$ and $g \in H^p(\beta)^*$. Note that

$$\langle f, g \rangle = \sum_{n=0}^{\infty} \hat{f}(n) \overline{\hat{g}(n)} \beta(n)^p.$$

Let $\hat{f}_k(n) = \delta_k(n)$. So $f_k(z) = z^k$ and then $\{f_k\}_k$ is a basis such that $\|f_k\| = \beta(k)$. Clearly M_z , the multiplication operator by z on $H^p(\beta)$ shifts the basis $\{f_k\}_k$.

Remember that a complex number λ is said to be a bounded point evaluation on $H^p(\beta)$ if the functional of point evaluation at λ, e_λ , is bounded. The functional of evaluation of the j -th derivative at λ is denoted by $e_\lambda^{(j)}$. These spaces are also studied in [1, 2, 4, 5, 6, 7, 8].

If Ω is a bounded domain in the complex domain \mathcal{C} , then by $H(\Omega)$ we mean the set of analytic functions on Ω . We will denote the open unit disc by U .

2. Main Results

We will investigate the relation between the generating function and the functional of point evaluations on $H^p(\beta)$. Also we will determine the structure of all non-zero linear functionals on $H^p(\beta)$ that are multiplicative. This extends some results of [1] into Banach spaces of formal power series. The differential of functionals of point evaluations are also considered.

Definition 1. *The generating function for the weighted Hardy space $H^p(\beta)$ is the function*

$$g(z) = \sum_{n=0}^{\infty} \frac{z^n}{\beta(n)^q}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Lemma 2. *If g is the generating function for a weighted Hardy space contained in $H(U)$, then $g \in H(U)$.*

Proof. Let g be the generating function for the weighted Hardy space $H^p(\beta)$ where $H^p(\beta) \subset H(U)$. Define

$$\hat{f}(n) = \begin{cases} 0 & n = 0 \\ \frac{1}{n\beta(n)} & n \neq 0 \end{cases}$$

and let $f(z) = \sum_{n=0}^{\infty} \hat{f}(n)z^n$. Now we have

$$\sum_{n=0}^{\infty} |\hat{f}(n)|^p \beta(n)^p = \sum_{n=1}^{\infty} \frac{1}{n^p} < \infty.$$

So $f \in H^p(\beta)$ and by assumption, it is analytic in the open unit disk U .

Thus the radius of convergence of its power series, R , is at least 1. Thus

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} |\hat{f}(n)|^{\frac{1}{n}} = \limsup_{n \rightarrow \infty} \left(\frac{1}{n\beta(n)} \right)^{\frac{1}{n}} \leq 1,$$

and so

$$\limsup_n \left(\frac{1}{\beta(n)^q} \right)^{\frac{1}{n}} = \limsup_n \left[\left(\frac{1}{n\beta(n)} \right)^{\frac{1}{n}} \right]^q \leq 1,$$

which implies that $g \in H(U)$. This completes the proof. \square

The next theorem shows the principle role of g : it generates the reproducing kernels for $H^p(\beta)$.

Lemma 3. *Let $H^p(\beta)$ be a weighted Hardy space contained in $H(U)$. For each point λ in U , the functional of evaluation at λ , e_λ , is a bounded linear functional and $\|e_\lambda\|^q = g(|\lambda|^q)$.*

Proof. For $|\lambda| < 1$, the analyticity of g on U implies that e_λ is in $H^q(\beta^{\frac{p}{q}})$. Indeed,

$$\|e_\lambda\|^q = \left\| \sum_{n=0}^{\infty} \frac{\bar{\lambda}^n}{\beta(n)^p} z^n \right\|^q = \sum_{n=0}^{\infty} \frac{|\lambda|^{nq}}{\beta(n)^q} = g(|\lambda|^q) < \infty.$$

This completes the proof. \square

Theorem 4. *If g , the generating function for $H^p(\beta)$, satisfies $g(1) = \infty$, then each non-zero bounded linear functional H on $H^p(\beta)$ such that $\langle fh, H \rangle = \langle f, H \rangle \langle h, H \rangle$ whenever f, h and fh are in $H^p(\beta)$ is in the form of $H = e_\lambda$ for some point λ in U .*

Proof. Suppose $H \in H^p(\beta)^*$ such that H is non-zero and satisfies

$$\langle fh, H \rangle = \langle f, H \rangle \langle h, H \rangle$$

whenever f, h and fh are in $H^p(\beta)$. Since the polynomials are dense in $H^p(\beta)$, this holds when f and h are polynomials.

Also, we note that

$$\langle f, H \rangle = \langle f_0 f, H \rangle = \langle f_0, H \rangle \langle f, H \rangle$$

for all f in $H^p(\beta)$, hence $\langle f_0, H \rangle = 1$. Letting $\lambda = \langle f_1, H \rangle$, it follows that

$$\langle f_2, H \rangle = \langle f_1 f_1, H \rangle = \langle f_1, H \rangle^2 = \lambda^2.$$

By induction, $\langle f_n, H \rangle = \lambda^n$ for all $n \in \mathbb{N}$. Remember that if

$$f(z) = \sum_{n=0}^{\infty} \hat{f}(n) z^n \in H^p(\beta)$$

and

$$G(z) = \sum_{n=0}^{\infty} \hat{G}(n) z^n \in H^q(\beta^{\frac{p}{q}}),$$

then

$$\langle f, G \rangle = \sum \hat{f}(n) \overline{\hat{G}(n)} \beta(n)^p.$$

Now if $H(z) = \sum_{n=0}^{\infty} \hat{H}(n) z^n$, we have

$$1 = \langle f_0, H \rangle = \overline{\hat{H}(0)} \beta(0)^p = \overline{\hat{H}(0)},$$

$$\lambda = \langle f_1, H \rangle = \overline{\hat{H}(1)} \beta(1)^p,$$

$$\vdots$$

$$\lambda^n = \langle f_n, H \rangle = \overline{\hat{H}(n)} \beta(n)^p.$$

So if $|\lambda| < 1$, then

$$H(z) = 1 + \sum_{n=1}^{\infty} \frac{\overline{\lambda^n}}{\beta(n)^p} z^n = e_{\lambda}(z),$$

and so H is the linear functional of evaluation at λ . If $|\lambda| \geq 1$, then we get

$$\|H\|^q = \sum_{n=0}^{\infty} \frac{|\lambda|^{nq}}{\beta(n)^q} \geq \sum_{n=0}^{\infty} \frac{1}{\beta(n)^q} = g(1) = \infty,$$

which contradicts the boundedness of the linear functional determined by H . This completes the proof. \square

As we saw, for λ in U ,

$$e_\lambda(z) = \sum_{n=0}^{\infty} \frac{1}{\beta(n)^p} (\bar{\lambda})^n z^n \in H^q(\beta^{p/q}).$$

This is not true in general if λ is on the unit circle.

Lemma 5. *Let $j \in \mathbb{N} \cup \{0\}$. If $\sum_{n=j}^{\infty} \frac{n^{qj}}{\beta(n)^q} < \infty$, then $e_\lambda^{(j)} \in H^q(\beta^{p/q})$ and $\langle f, e_\lambda^{(j)} \rangle = f^{(j)}(\lambda)$ for all λ in \bar{U} and f in $H^p(\beta)$.*

Proof. For λ in \bar{U} we have

$$e_\lambda^{(j)}(z) = \sum_{n=j}^{\infty} \frac{n(n-1) \cdots (n-j+1)}{\beta(n)^p} (\bar{\lambda})^{n-j} z^n,$$

and so

$$\begin{aligned} \|e_\lambda^{(j)}(z)\|^q &= \sum_{n=j}^{\infty} \left(\frac{n(n-1) \cdots (n-j+1)}{\beta(n)^p} \right)^q |\lambda|^{(n-j)q} \beta(n)^p \\ &\leq \sum_{n=j}^{\infty} \frac{(n(n-1) \cdots (n-j+1))^q}{\beta(n)^q} \end{aligned}$$

$$\leq \sum_{n=j}^{\infty} \frac{n^{qj}}{\beta(n)^q} < \infty.$$

Thus $e_{\lambda}^{(j)} \in H^q(\beta^{p/q})$. Now if

$$f(z) = \sum_{n=0}^{\infty} \hat{f}(n) z^n \in H^p(\beta),$$

then by a Theorem in [4], for all λ in \bar{U} we have:

$$\begin{aligned} \langle f, e_{\lambda}^{(j)} \rangle &= \sum_n \hat{f}(n) (\overline{e_{\lambda}^{(j)}(n)}) \beta(n)^p \\ &= \sum_{n=j}^{\infty} \hat{f}(n) \frac{n(n-1) \cdots (n-j+1)}{\beta(n)^p} \lambda^{n-j} \beta(n)^p \\ &= \sum_{n=j}^{\infty} n(n-1) \cdots (n-j+1) \hat{f}(n) \lambda^{n-j} = f^{(j)}(\lambda). \end{aligned}$$

This completes the proof. \square

Note that if $\sum_{n=j}^{\infty} \frac{n^{qj}}{\beta(n)^q} < \infty$, then $H^p(\beta)$ is very small and for every function in $H^p(\beta)$, the j -th derivative exists and is continuous on the unit circle.

Corollary 6. *If $\sum_{n=0}^{\infty} \frac{1}{\beta(n)^q} < \infty$, then the generating function belongs to $H^p(\beta)$.*

Proof. Let g be the generating function. Thus $g(z) = \sum_n \frac{1}{\beta(n)^q} z^n$ and

$$\|g\|^q = \sum_n |\hat{g}(n)|^p \beta(n)^p = \sum_n \frac{1}{\beta(n)^q} < \infty.$$

So $g \in H^p(\beta)$. \square

Corollary 7. *If $\sum_n \frac{1}{\beta(n)^q} < \infty$ and $\liminf_n \beta(n)^{1/n} = 1$, then the generating function is in $H(U)$.*

Proof. Clearly we can see that $H^p(\beta) \subset H(U)$ and so by the Lemma 2, the proof is complete. \square

Theorem 8. *In the weighted Hardy space $H^p(\beta)$ for which $g(1) = \infty$ for all integer $j \geq 0$, the normalized functional of point evaluations, $\frac{e_{w_n}^{(j)}}{\|e_{w_n}^{(j)}\|}$, tends to zero weakly as $w_n \rightarrow \xi \in \partial U$.*

Proof. For $j = 0$, the norm of the functional of point evaluations are given by the generating function g and indeed $\|e_{w_n}\|^q = g(|w_n|^q)$. Since

$$\lim_n g(|w_n|^q) = g(1) = \sum_n \frac{1}{\beta(n)^q} = \infty,$$

it follows that $\|e_{w_n}\|$ tends to infinity and for every polynomial p ,

$$\lim_m |\langle p, \frac{e_{w_n}}{\|e_{w_n}\|} \rangle| = \lim_n \frac{|p(w_n)|}{\|e_{w_n}\|} = 0.$$

But the polynomials are dense in $H^p(\beta)$, thus $\frac{e_{w_n}}{\|e_{w_n}\|} \rightarrow 0$ weakly as $n \rightarrow \infty$. If $j > 0$, then

$$\lim_n \|e_{w_n}^{(j)}\|^q = \lim_m \sum_{n=j}^{\infty} \left(\frac{n(n-1) \cdots (n-j+1)}{\beta(n)} \right)^q |w_m|^{q(n-j)}$$

$$\begin{aligned}
&= \sum_{n=j}^{\infty} \left(\frac{n(n-1) \cdots (n-j+1)}{\beta(n)} \right)^q \\
&\geq \sum_{n=j}^{\infty} \frac{1}{\beta(n)^q} = \infty.
\end{aligned}$$

Thus $\|e_{w_m}^{(j)}\|$ tends to infinity and for every polynomial p ,

$$\lim_m \left| \left\langle p, \frac{e_{w_m}^{(j)}}{\|e_{w_m}^{(j)}\|} \right\rangle \right| = \lim_m \frac{|p^{(j)}(w_m)|}{\|e_{w_m}^{(j)}\|} = 0.$$

Since the polynomials are dense in $H^p(\beta)$, the proof is complete. \square

References

- [1] C. C. Cowen and B. D. MacCluer, *Composition operators on spaces of analytic functions*, CRC press, Inc. , 1995.
- [2] K. Seddighi and B. Yousefi, On the reflexivity of operators on function spaces, *Proceedings of the American Mathematical Society*, 116 (1992), 45-52.
- [3] A. L. Shields, Weighted shift operators and analytic function theory, *Mathematical Survey, A. M. S. Providence*, 13 (1974), 49-128.
- [4] B. Yousefi, On the space $l^p(\beta)$, *Rendiconti Del Circolo Matematico Di Palermo, Serie II. Tomo XLIX* (2000), 115-120.
- [5] B. Yousefi, Unicellularity of the multiplication operator on Banach spaces of formal power series, *Studia Mathematica*, 147 (2001), 201-209.
- [6] B. Yousefi and K. Jahedi, Application of the Rosenthal-Dor Theorem on Banach spaces of formal power series, *Islamic Azad University Journal of sciences*, Fall (2001), 3147-3168.
- [7] B. Yousefi and S. Jahedi, Composition operators on Banach spaces of formal power series, *Bollettino Della Unione Matematica Italiana*, (8) 6-B (2003), 481-487.

- [8] B. Yousefi, On the eighteenth question of Allen shields, *International Journal of Mathematics*, 16 (1) (2005), 37-42.

Khadijeh Jahedi

Department of Mathematics
Islamic Azad University - Shiraz Branch
Shiraz, Iran
E-mail: Mjahedi80@yahoo.com

Bahman Yousefi

Department of Mathematics
College of Sciences
Shiraz University
Shiraz 71454, Iran
E-mail: Yousefi@Math.Susc.ac.ir