

# LIE SYMMETRIES OF SCHRÖDINGER EQUATION ON A SPHERE

YADOLLAH ARYANEJAD

**ABSTRACT.** We are going to discuss an object with a mass attached to a spring and vibrating on the surface of a sphere (see Figure 1). To do this, we first reconstruct the Schrödinger equation on a sphere. In fact, the paper considers the question of a quantum system obeying the Schrodinger equation on a sphere. After a brief introduction we set up the Hamiltonian of the system and the corresponding Schrodinger equation. We provide the Lie algebra of symmetries and build the optimal system of Lie subalgebras and its adjoint presentation. Reductions of similarities related to subalgebras are obtained which are used in our study of the 3D quantum harmonic oscillator on a sphere as a special case of the new equation, and possible solutions are proposed.

## 1. INTRODUCTION

Studies in this area are in progress since such equations depict the states and properties of nonlinear phenomena, broaden vision in terms of physical aspects, and then become more practical in engineering and other sciences, so the search for accurate solutions is important in Nonlinear is in several ways like plasma laser radiation [1, 13, 14]. To obtain

---

2020 *Mathematics Subject Classification.* 70G65, 34C14, 53C50.

*Key words and phrases.* Schrödinger equation, symmetry group, invariant solution, 3D Harmonic oscillator, sphere, Lie groups.




FIGURE 1. The object oscillates on a sphere.

an equation that can describe the oscillation of an object attached to a spring on a sphere, in general, we first consider Schrödinger equation in three-dimensional space:

$$(1) \quad -\frac{h^2}{2m}\Delta\Psi + v(x, y, z)\Psi = i\hbar\Psi_t.$$

This equation reflects the wave nature of our quantum solutions  $\Psi(x, y, z)$ . A significant part of quantum mechanics is devoted to the study of solutions to the Schrödinger equation. Equation (1) governs the time dependence of the wave-function of an object moving inside a given potential,  $v(x, y, z)$ . A unique role is played by solutions to (1) that have the simple form:  $\Psi = \psi(x, y, z)\exp(\frac{-iEt}{\hbar})$  where the function  $\psi$  satisfies the Schrödinger equation of the Schrödinger eigenvalue equation:

$$(2) \quad \begin{cases} -\frac{h^2}{2m}\Delta\psi + v(x, y, z)\psi = E\psi, \\ \Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}, \end{cases}$$

which is time-independent. In both of these equations,  $h$  and  $m$  represent real constants.  $E$  is a constant that emerges during the separation of variables procedure. Sometimes quantum problems arise on a sphere. Therefore, it is necessary to examine Equation (2) on a sphere. It is well known, the metric on  $S^2 \times R$  is:

$$(3) \quad ds^2 = dz^2 - dx^2 - \sin^2 x dy^2 \quad f \in C^\infty(G).$$

Adjusting the metric (3) on  $S^2 \times R$  and rewriting Equation (2), the differential equation on the sphere would be:

$$(4) \quad u_{zz} = u_{xx} + (\cot x)u_x + (\csc^2 x)u_{yy} + (2m/\hbar^2)(E - v)u$$

where  $u$  is the function  $\psi$  defined on  $S^2 \times R$ .

The Schrodinger equation for a harmonic oscillator may be obtained by using the classical spring potential

$$(5) \quad \begin{aligned} v(x, y, z) &= \frac{1}{2}q_x x^2 + \frac{1}{2}q_y y^2 + \frac{1}{2}q_z z^2, \\ \omega_x^2 &= \frac{q_x}{m}, \quad \omega_y^2 = \frac{q_y}{m}, \quad \omega_z^2 = \frac{q_z}{m}, \end{aligned}$$

where  $\omega$  is angular frequency and  $q_x$ ,  $q_y$  and  $q_z$  are bond force constants. Adjusting the metric (3) on  $S^2 \times R$  and rewriting Equation (5), the spring potential on the sphere would be:

$$(6) \quad v(x, y, z) = \frac{1}{2}q_x x^2 + \frac{1}{2}q_y \sin^4(x)y^2 + \frac{1}{2}q_z z^2,$$

The Schrodinger equation (4) with this form of potential is

$$(7) \quad \begin{aligned} u_{zz} &= u_{xx} + (\cot x)u_x + (\csc^2 x)u_{yy} \\ &+ (2m/\hbar^2)(E - \frac{1}{2}q_x x^2 - \frac{1}{2}q_y \sin^4(x)y^2 - \frac{1}{2}q_z z^2)u, \end{aligned}$$

where the particle oscillates on a sphere.

Equation (4) is the general state of the Equation (7), which we try to solve by the method of Lie symmetry groups. Equation (7) describes an oscillating object on the surface of a sphere. This equation is reported in the last section, and possible solutions are presented. On the other hand, using the structure of Lie algebra in investigating the geometric concepts of the relevant spaces is a natural and inevitable process [6–8]. The

symmetry group approach or Lie's approach itself, which is a computational method to find invariant solutions, is crucially utilized in revealing the answer of PDEs and ODEs. Performing the Lie symmetry group procedure; the problem of symmetry classification for different equations is widely considered in various spaces [2–5, 11, 12]. Utilizing this plan of action, one finds appropriate solutions via studied ones, investigates the invariant solutions, and decreases the order of ODEs [9, 10, 15]. In this article, utilizing Lie's procedure, we get symmetries of the Schrödinger differential equation on the sphere. Next, utilizing Ibragimov's method an optimal sub-algebras structure related to the symmetry Lie algebra is presented. The article is collected as follows:

- We describe the symmetry algebra infinitesimal generators of Equation (4), and gain several outcomes.
- We build the optimal systems of sub-algebras.
- We give the Lie invariants, and some other concepts related to the infinitesimal symmetries of Equation (4).
- Possible solutions of the 3D Harmonic Oscillator on the sphere are discussed.

## 2. INFINITESIMAL GENERATORS OF EQUATION (4)

Commonly,

$$(8) \quad \begin{cases} \Delta_h(x, u^{(j)}) = 0, & h = 1, \dots, n, \\ x = (x^1, \dots, x^p), \\ u = (u^1, \dots, u^q), \end{cases}$$

define a PDE structure of order  $j$ th, where  $u$  is dependent on  $x$ , and  $u^{(i)}$  means  $\partial_i u / (\partial x)^i$ . Local infinitesimal generators of the above structure that as a Lie group acts on the manifold  $X \times U$ , is:

$$(9) \quad \tilde{x}^i = x^i + \delta \zeta^i(x, u) + \mathcal{O}(\delta^2), \quad i = 1, \dots, p,$$

$$(10) \quad \tilde{u}^j = u^j + \delta \phi_j(x, u) + \mathcal{O}(\delta^2), \quad j = 1, \dots, q,$$

where  $\zeta^i$  and  $\phi^j$  represent the infinitesimal transformations for  $\{x^1, \dots, x^p\}$  and  $\{u^1, \dots, u^q\}$ , respectively. A given local infinitesimal generators related to the all transformations (9) as a group, is

$$(11) \quad \mathfrak{X} = \sum_{i=1}^p \zeta^i(x, u) \partial_{x^i} + \sum_{j=1}^q \phi_j(x, u) \partial_{u^j}$$

Now to utilize the mentioned technique for Equation (4), infinitesimal transformations with one parameter as a Lie group is assumed: ( $x^1, x^2$  and  $x^3$  are substituted by  $x, y$  and  $z$  respectively to not use index,)

$$\begin{cases} \tilde{x} = x + \delta \zeta^1(x, y, z, u, v) + \mathcal{O}(\delta^2), \\ \tilde{y} = y + \delta \zeta^2(x, y, z, u, v) + \mathcal{O}(\delta^2), \\ \tilde{z} = z + \delta \zeta^3(x, y, z, u, v) + \mathcal{O}(\delta^2), \\ \tilde{u} = u + \delta \phi_1(x, y, z, u, v) + \mathcal{O}(\delta^2), \\ \tilde{v} = v + \delta \phi_2(x, y, z, u, v) + \mathcal{O}(\delta^2). \end{cases}$$

The related symmetry generator will be:

$$(12) \quad \mathfrak{X} = \varsigma^1(x, y, z, u, v)\partial_x + \varsigma^2(x, y, z, u, v)\partial_y + \varsigma^3(x, y, z, u, v)\partial_t + \phi_1(x, y, z, u, v)\partial_u + \phi_2(x, y, z, u, v)\partial_v.$$

The status of existence of invariance is equivalent to the following explanation:

$$\begin{aligned} Pr^{(2)}\mathfrak{X}[u_{xx} + (\cot x)u_x + (\csc^2 x)u_{yy} - u_{zz} + (2m/h^2)(E - v(x, y, z))u] &= 0, \\ \text{whenever :} \\ u_{xx} + (\cot x)u_x + (\csc^2 x)u_{yy} - u_{zz} + (2m/h^2)(E - v(x, y, z))u &= 0. \end{aligned}$$

Since,  $\varsigma^1, \varsigma^2, \varsigma^3, \phi_1$  and  $\phi_2$  are functions with variables  $x, y, z, u$  and  $v$ , vanishing the sole coefficients, we earn the following specific equations:

$$\begin{aligned} h^2 \sin(x) \varsigma_v^2 &= 0, & h^2 \sin(x) \varsigma_{uu}^2 &= 0, & h^2 \sin(x) \varsigma_v^1 &= 0, \\ h^2 \sin(x) \varsigma_{uu}^1 &= 0, & h^2 \sin(x) \varsigma_{vv}^2 &= 0, & h^2 \sin(x) \varsigma_v^3 &= 0, \\ & \dots \end{aligned}$$

The number of these equations is 89. Examining these PDEs, we have a statement as:

**Theorem 2.1.** *The point symmetry group of Equation (4) as a Lie group owns a Lie sub-algebra consists of (12) which  $\xi$ s and  $\phi$ s are the infinitesimals as follows:*

$$\begin{aligned} \varsigma^1 &= ((c_6 \sin(y) + c_7 \cos(y)) \cos(z) + \sin(z)(c_3 \sin(y) + c_4 \cos(y))) \cos(x) \\ &\quad + c_1 \sin(y) + c_2 \cos(y) + c_5 \sin(z) \sin(x) + c_8 \cos(z) \sin(x), \\ \varsigma^2 &= \frac{(c_6 \cos(y) - c_7 \sin(y)) \cos(z) + \sin(z)(c_3 \cos(y) + c_4 \sin(y))}{\sin(x)} \\ &\quad + \frac{c_1 \cos(y) - c_2 \sin(y)}{\tan(x)} + c_{10}, \\ \varsigma^3 &= (-(c_6 \sin(y) + c_7 \cos(y)) \sin(z) + \cos(z)(c_3 \sin(y) + c_4 \cos(y))) \sin(x) \\ &\quad - c_5 \cos(z) \cos(x) + c_8 \cos(x) \sin(z) + c_9, \\ \phi_1 &= \frac{1}{2}((c_6 \sin(y) + c_7 \cos(y)) \cos(z) + \sin(z)(c_3 \sin(y) + c_4 \cos(y))) \sin(x) u, \\ &\quad \frac{1}{2} c_5 \sin(z) \cos(x) u + \frac{1}{2} c_8 \cos(z) \cos(x) u + c_{11} u + \alpha(u), \\ \phi_2 &= \frac{1}{4 \sin^2(x) m u} (((c_6 \sin(y) + c_7 \cos(y)) \cos(z) + \sin(z)(c_3 \sin(y) + c_4 \cos(y))) \\ &\quad \sin^2(x) - (c_5 \sin(z) + c_8 \cos(z)) \cos(x) \sin(x)) \left( \frac{h^2}{8} + mE + mv \right) \\ &\quad (-8) \sin(x) u + 2h^2 \sin^2(x) (\alpha_{xx} - \alpha_{zz}) + 2h^2 \alpha_{yy} \\ &\quad + \left( \frac{1}{4} h^2 \cos(x) u_x + \frac{1}{2} \sin(x) m (E - v) \right) 8 \sin(x)). \end{aligned}$$

where  $c_i, i = 1, \dots, 11$  are real constant.

**Corollary 2.2.** *Every Lie group consists of symmetries with one-parameter of (4) has eleven-dimensional Lie subalgebra obtained from the following generators:*

$$\begin{aligned}
\mathfrak{X}_1 &= \partial_y, \\
\mathfrak{X}_2 &= \partial_z, \\
\mathfrak{X}_3 &= \sin(y)\partial_x + \cos(y)\cot(x)\partial_y, \\
\mathfrak{X}_4 &= \cos(y)\partial_x - \sin(y)\cot(x)\partial_y, \\
\mathfrak{X}_5 &= \sin(x)\sin(z)\partial_x - \cos(x)\cos(z)\partial_z - \frac{1}{2}u\cos(x)\sin(z)\partial_u \\
&\quad + \Omega\cos(x)\sin(z)\partial_v, \\
\mathfrak{X}_6 &= \sin(x)\cos(z)\partial_x + \sin(z)\cos(x)\partial_z - \frac{1}{2}u\cos(x)\cos(z)\partial_u \\
&\quad + \Omega\cos(x)\cos(z)\partial_v, \\
\mathfrak{X}_7 &= \cos(x)\sin(z)\sin(y)\partial_x + \frac{\cos(y)\sin(z)}{\sin(x)}\partial_y + \sin(x)\sin(y)\cos(z)\partial_z \\
&\quad + \frac{1}{2}u\sin(x)\sin(y)\sin(z)\partial_u - \Omega\sin(x)\sin(y)\sin(z)\partial_v, \\
\mathfrak{X}_8 &= \cos(x)\sin(z)\cos(y)\partial_x - \frac{\sin(y)\sin(z)}{\sin(x)}\partial_y + \sin(x)\cos(y)\cos(z)\partial_z \\
&\quad + \frac{1}{2}u\sin(x)\cos(y)\sin(z)\partial_u - \Omega\sin(x)\cos(y)\sin(z)\partial_v, \\
\mathfrak{X}_9 &= \cos(x)\cos(z)\sin(y)\partial_x + \frac{\cos(y)\cos(z)}{\sin(x)}\partial_y - \sin(x)\sin(y)\sin(z)\partial_z \\
&\quad + \frac{1}{2}u\sin(x)\sin(y)\cos(z)\partial_u - \Omega\sin(x)\sin(y)\cos(z)\partial_v, \\
\mathfrak{X}_{10} &= \cos(x)\cos(z)\cos(y)\partial_x - \frac{\sin(y)\cos(z)}{\sin(x)}\partial_y - \sin(x)\cos(y)\sin(z)\partial_z \\
&\quad + \frac{1}{2}u\sin(x)\cos(y)\cos(z)\partial_u - \Omega\sin(x)\cos(y)\cos(z)\partial_v, \\
\mathfrak{X}_{11} &= u\partial_u, \\
\mathfrak{X}_\alpha &= \alpha\partial_u + \frac{(E-v)\alpha\partial_v}{u},
\end{aligned}$$

where  $\Omega = \frac{8mE - 8mv + h^2}{4m}$  (and  $\partial_x \equiv \frac{\partial}{\partial x}, \dots$ ).

We deliver Lie bracket for Eq.(4) by Table (1). The phrase  $[\mathfrak{X}_i, \mathfrak{X}_j] = \mathfrak{X}_i\mathfrak{X}_j - \mathfrak{X}_j\mathfrak{X}_i$  characterizes the Values in row  $i^{th}$  and column  $j^{th}$ ,  $i, j = 1, \dots, 11$ .

For instance, the flow of  $\mathfrak{X}_2$  in Corollary 2.2 is expressed by

$$\Phi_\epsilon = (x, y, z + \epsilon).$$

### 3. 1D SUBALGEBRAS OF EQUATION (4)

Utilizing the symmetry technique, one can specify the one parameter optimal structure of Equation (4). Providing special subgroups that offer different sorts of solutions is essential. Thus, we want to look for an invariant solution that is not identical to a transformation from the whole symmetry group. Such an issue causes to express the sense of an optimal structure of sub-algebra. In the study of 1D sub-algebras, the classification

TABLE 1. Lie algebra for Eq.(4).

$[ , ]$	$\mathfrak{X}_1$	$\mathfrak{X}_2$	$\mathfrak{X}_3$	$\mathfrak{X}_4$	$\mathfrak{X}_5$	$\mathfrak{X}_6$	$\mathfrak{X}_7$	$\mathfrak{X}_8$	$\mathfrak{X}_9$	$\mathfrak{X}_{10}$	$\mathfrak{X}_{11}$
$\mathfrak{X}_1$	0	0	$\mathfrak{X}_4$	$-\mathfrak{X}_3$	0	0	$\mathfrak{X}_8$	$-\mathfrak{X}_7$	$\mathfrak{X}_{10}$	$-\mathfrak{X}_9$	0
$\mathfrak{X}_2$	*	0	0	0	$\mathfrak{X}_6$	$-\mathfrak{X}_5$	$\mathfrak{X}_9$	$\mathfrak{X}_{10}$	$-\mathfrak{X}_7$	$\mathfrak{X}_8$	0
$\mathfrak{X}_3$	*	*	0	$\mathfrak{X}_1$	$\mathfrak{X}_7$	$\mathfrak{X}_9$	$-\mathfrak{X}_5$	0	$-\mathfrak{X}_6$	0	0
$\mathfrak{X}_4$	*	*	*	0	$\mathfrak{X}_8$	$\mathfrak{X}_{10}$	0	$-\mathfrak{X}_5$	0	$-\mathfrak{X}_6$	0
$\mathfrak{X}_5$	*	*	*	*	0	$-\mathfrak{X}_2$	$-\mathfrak{X}_3$	$-\mathfrak{X}_4$	0	0	0
$\mathfrak{X}_6$	*	*	*	*	*	0	0	0	$-\mathfrak{X}_3$	$-\mathfrak{X}_4$	0
$\mathfrak{X}_7$	*	*	*	*	*	*	0	$-\mathfrak{X}_1$	$-\mathfrak{X}_2$	0	0
$\mathfrak{X}_8$	*	*	*	*	*	*	*	0	0	$-\mathfrak{X}_2$	0
$\mathfrak{X}_9$	*	*	*	*	*	*	*	*	0	$-\mathfrak{X}_1$	0
$\mathfrak{X}_{10}$	*	*	*	*	*	*	*	*	*	0	0
$\mathfrak{X}_{11}$	*	*	*	*	*	*	*	*	*	*	0

question turns into arranging the adjoint representation orbits. An optimal structure of the sub-algebras question is replied by presuming a candidate of any set of related sub-algebras [17] and [16]. Adjoint candidate of every  $X_i$ , for  $i = 1, \dots, 11$  is characterized as:

$$(13) \quad \text{Ad}(e^{(\mathfrak{s} \cdot \mathfrak{X}_i)} \cdot \mathfrak{X}_j) = \mathfrak{X}_j - \mathfrak{s} \cdot [\mathfrak{X}_i, \mathfrak{X}_j] + \frac{\mathfrak{s}^2}{2} \cdot [\mathfrak{X}_i, [\mathfrak{X}_i, \mathfrak{X}_j]] - \dots,$$

where  $\mathfrak{s}$  is a parameter and  $[\mathfrak{X}_i, \mathfrak{X}_j]$  has characterized in Table (1) for  $1 \leq i, j \leq 11$  ([16], p (199)). We show the Lie algebra of (13) by  $\mathfrak{g}$ , and we collect the adjoint action in Table (2). An optimal system of one-dimensional subalgebras is constructed by utilizing Ibragimov's method.

**Theorem 3.1.** *A 1D optimal structure of Eq.(4) is presented as:*

- |                                     |  |  |
|-------------------------------------|--|--|
| 1) $\mathfrak{X}_1 \pm X_{11},$     | 12) $\mathfrak{X}_1 \pm \mathfrak{X}_2 \pm X_{11},$    | 23) $\mathfrak{X}_6 \pm \mathfrak{X}_7 \pm X_{11},$    |
| 2) $\mathfrak{X}_2 \pm X_{11},$     | 13) $\mathfrak{X}_1 \pm \mathfrak{X}_5 \pm X_{11},$    | 24) $\mathfrak{X}_6 \pm \mathfrak{X}_8 \pm X_{11},$    |
| 3) $\mathfrak{X}_3 \pm X_{11},$     | 14) $\mathfrak{X}_1 \pm \mathfrak{X}_6 \pm X_{11},$    | 25) $\mathfrak{X}_7 \pm \mathfrak{X}_{10} \pm X_{11},$ |
| 4) $\mathfrak{X}_4 \pm X_{11},$     | 15) $\mathfrak{X}_2 \pm \mathfrak{X}_3 \pm X_{11},$    | 26) $\mathfrak{X}_8 \pm \mathfrak{X}_9 \pm X_{11},$    |
| 5) $\mathfrak{X}_5 \pm X_{11},$     | 16) $\mathfrak{X}_2 \pm \mathfrak{X}_4 \pm X_{11},$    |  |
| 6) $\mathfrak{X}_6 \pm X_{11},$     | 17) $\mathfrak{X}_3 \pm \mathfrak{X}_8 \pm X_{11},$    |  |
| 7) $\mathfrak{X}_7 \pm X_{11},$     | 18) $\mathfrak{X}_3 \pm \mathfrak{X}_{10} \pm X_{11},$ |  |
| 8) $\mathfrak{X}_8 \pm X_{11},$     | 19) $\mathfrak{X}_4 \pm \mathfrak{X}_7 \pm X_{11},$    |  |
| 9) $\mathfrak{X}_9 \pm X_{11},$     | 20) $\mathfrak{X}_4 \pm \mathfrak{X}_9 \pm X_{11},$    |  |
| 10) $\mathfrak{X}_{10} \pm X_{11},$ | 21) $\mathfrak{X}_5 \pm \mathfrak{X}_9 \pm X_{11},$    |  |
| 11) $\mathfrak{X}_{11},$            | 22) $\mathfrak{X}_5 \pm \mathfrak{X}_{10} \pm X_{11},$ |  |

where  $c_i \in \mathbb{R}$  are real numeral coefficients for  $i = 1, \dots, 4$ .

TABLE 2. Adjoint presentation

$Ad$	$\mathfrak{X}_1$	$\mathfrak{X}_2$	$\mathfrak{X}_3$	$\mathfrak{X}_4$
$\mathfrak{X}_1$	$\mathfrak{X}_1$	$\mathfrak{X}_2$	$\cos(\mathfrak{s})\mathfrak{X}_3 - \sin(\mathfrak{s})\mathfrak{X}_4$	$\cos(\mathfrak{s})\mathfrak{X}_4 + \sin(\mathfrak{s})\mathfrak{X}_3$
$\mathfrak{X}_2$	$\mathfrak{X}_1$	$\mathfrak{X}_2$	$\mathfrak{X}_3$	$\mathfrak{X}_4$
$\mathfrak{X}_3$	$\cos(\mathfrak{s})\mathfrak{X}_1 + \sin(\mathfrak{s})\mathfrak{X}_4$	$\mathfrak{X}_2$	$\mathfrak{X}_3$	$\cos(\mathfrak{s})\mathfrak{X}_4 - \sin(\mathfrak{s})\mathfrak{X}_1$
$\mathfrak{X}_4$	$\cos(\mathfrak{s})\mathfrak{X}_1 - \sin(\mathfrak{s})\mathfrak{X}_3$	$\mathfrak{X}_2$	$\cos(\mathfrak{s})\mathfrak{X}_3 + \sin(\mathfrak{s})\mathfrak{X}_1$	$\mathfrak{X}_4$
$\mathfrak{X}_5$	$\mathfrak{X}_1$	$\cosh(\mathfrak{s})\mathfrak{X}_2 - \sinh(\mathfrak{s})\mathfrak{X}_6$	$\cosh(\mathfrak{s})\mathfrak{X}_3 - \sinh(\mathfrak{s})\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_4 - \sinh(\mathfrak{s})\mathfrak{X}_8$
$\mathfrak{X}_6$	$\mathfrak{X}_1$	$\cosh(\mathfrak{s})\mathfrak{X}_2 + \sinh(\mathfrak{s})\mathfrak{X}_5$	$\cosh(\mathfrak{s})\mathfrak{X}_3 - \sinh(\mathfrak{s})\mathfrak{X}_9$	$\cosh(\mathfrak{s})\mathfrak{X}_4 - \sinh(\mathfrak{s})\mathfrak{X}_{10}$
$\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_1 - \sinh(\mathfrak{s})\mathfrak{X}_8$	$\cosh(\mathfrak{s})\mathfrak{X}_2 - \sinh(\mathfrak{s})\mathfrak{X}_9$	$\cosh(\mathfrak{s})\mathfrak{X}_3 + \sinh(\mathfrak{s})\mathfrak{X}_5$	$\mathfrak{X}_4$
$\mathfrak{X}_8$	$\cosh(\mathfrak{s})\mathfrak{X}_1 + \sinh(\mathfrak{s})\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_2 - \sinh(\mathfrak{s})\mathfrak{X}_{10}$	$\mathfrak{X}_3$	$\cosh(\mathfrak{s})\mathfrak{X}_4 + \sinh(\mathfrak{s})\mathfrak{X}_5$
$\mathfrak{X}_9$	$\cosh(\mathfrak{s})\mathfrak{X}_1 - \sinh(\mathfrak{s})\mathfrak{X}_{10}$	$\cosh(\mathfrak{s})\mathfrak{X}_2 + \sinh(\mathfrak{s})\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_3 + \sinh(\mathfrak{s})\mathfrak{X}_6$	$\mathfrak{X}_4$
$\mathfrak{X}_{10}$	$\cosh(\mathfrak{s})\mathfrak{X}_1 + \sinh(\mathfrak{s})\mathfrak{X}_9$	$\cosh(\mathfrak{s})\mathfrak{X}_2 + \sinh(\mathfrak{s})\mathfrak{X}_8$	$\mathfrak{X}_3$	$\cosh(\mathfrak{s})\mathfrak{X}_4 + \sinh(\mathfrak{s})\mathfrak{X}_6$
$\mathfrak{X}_{11}$	$\mathfrak{X}_1$	$\mathfrak{X}_2$	$\mathfrak{X}_3$	$\mathfrak{X}_4$
$Ad$	$\mathfrak{X}_5$	$\mathfrak{X}_6$	$\mathfrak{X}_7$	$\mathfrak{X}_8$
$\mathfrak{X}_1$	$\mathfrak{X}_5$	$\mathfrak{X}_6$	$\cos(\mathfrak{s})\mathfrak{X}_7 - \sin(\mathfrak{s})\mathfrak{X}_8$	$\cos(\mathfrak{s})\mathfrak{X}_8 + \sin(\mathfrak{s})\mathfrak{X}_7$
$\mathfrak{X}_2$	$\cos(\mathfrak{s})\mathfrak{X}_5 - \sin(\mathfrak{s})\mathfrak{X}_6$	$\cos(\mathfrak{s})\mathfrak{X}_6 + \sin(\mathfrak{s})\mathfrak{X}_5$	$\cos(\mathfrak{s})\mathfrak{X}_7 - \sin(\mathfrak{s})\mathfrak{X}_9$	$\cos(\mathfrak{s})\mathfrak{X}_8 - \sin(\mathfrak{s})\mathfrak{X}_{10}$
$\mathfrak{X}_3$	$\cos(\mathfrak{s})\mathfrak{X}_5 - \sin(\mathfrak{s})\mathfrak{X}_7$	$\cos(\mathfrak{s})\mathfrak{X}_6 - \sin(\mathfrak{s})\mathfrak{X}_9$	$\cos(\mathfrak{s})\mathfrak{X}_7 + \sin(\mathfrak{s})\mathfrak{X}_5$	$\mathfrak{X}_8$
$\mathfrak{X}_4$	$\cos(\mathfrak{s})\mathfrak{X}_5 - \sin(\mathfrak{s})\mathfrak{X}_8$	$\cos(\mathfrak{s})\mathfrak{X}_6 - \sin(\mathfrak{s})\mathfrak{X}_{10}$	$\mathfrak{X}_7$	$\cos(\mathfrak{s})\mathfrak{X}_8 + \sin(\mathfrak{s})\mathfrak{X}_5$
$\mathfrak{X}_5$	$\mathfrak{X}_5$	$\cosh(\mathfrak{s})\mathfrak{X}_6 - \sinh(\mathfrak{s})\mathfrak{X}_2$	$\cosh(\mathfrak{s})\mathfrak{X}_7 - \sinh(\mathfrak{s})\mathfrak{X}_3$	$\cosh(\mathfrak{s})\mathfrak{X}_8 - \sinh(\mathfrak{s})\mathfrak{X}_4$
$\mathfrak{X}_6$	$\cosh(\mathfrak{s})\mathfrak{X}_7 + \sinh(\mathfrak{s})\mathfrak{X}_2$	$\mathfrak{X}_6$	$\mathfrak{X}_7$	$\mathfrak{X}_8$
$\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_7 + \sinh(\mathfrak{s})\mathfrak{X}_3$	$\mathfrak{X}_6$	$\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_8 - \sinh(\mathfrak{s})\mathfrak{X}_1$
$\mathfrak{X}_8$	$\cosh(\mathfrak{s})\mathfrak{X}_7 + \sinh(\mathfrak{s})\mathfrak{X}_4$	$\mathfrak{X}_6$	$\cosh(\mathfrak{s})\mathfrak{X}_7 + \sinh(\mathfrak{s})\mathfrak{X}_1$	$\mathfrak{X}_8$
$\mathfrak{X}_9$	$\mathfrak{X}_5$	$\cosh(\mathfrak{s})\mathfrak{X}_6 + \sinh(\mathfrak{s})\mathfrak{X}_3$	$\cosh(\mathfrak{s})\mathfrak{X}_7 + \sinh(\mathfrak{s})\mathfrak{X}_2$	$\mathfrak{X}_8$
$\mathfrak{X}_{10}$	$\mathfrak{X}_5$	$\cosh(\mathfrak{s})\mathfrak{X}_6 + \sinh(\mathfrak{s})\mathfrak{X}_4$	$\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_8 + \sinh(\mathfrak{s})\mathfrak{X}_2$
$\mathfrak{X}_{11}$	$\mathfrak{X}_5$	$\mathfrak{X}_6$	$\mathfrak{X}_7$	$\mathfrak{X}_8$
$Ad$	$\mathfrak{X}_9$	$\mathfrak{X}_{10}$	$\mathfrak{X}_{11}$	
$\mathfrak{X}_1$	$\cos(\mathfrak{s})\mathfrak{X}_9 - \sin(\mathfrak{s})\mathfrak{X}_{10}$	$\cos(\mathfrak{s})\mathfrak{X}_{10} + \sin(\mathfrak{s})\mathfrak{X}_9$	$\mathfrak{X}_1$	
$\mathfrak{X}_2$	$\cos(\mathfrak{s})\mathfrak{X}_9 + \sin(\mathfrak{s})\mathfrak{X}_7$	$\cos(\mathfrak{s})\mathfrak{X}_{10} + \sin(\mathfrak{s})\mathfrak{X}_8$	$\mathfrak{X}_2$	
$\mathfrak{X}_3$	$\cos(\mathfrak{s})\mathfrak{X}_9 + \sin(\mathfrak{s})\mathfrak{X}_6$	$\mathfrak{X}_{10}$	$\mathfrak{X}_3$	
$\mathfrak{X}_4$	$\mathfrak{X}_9$	$\cos(\mathfrak{s})\mathfrak{X}_{10} + \sin(\mathfrak{s})\mathfrak{X}_6$	$\mathfrak{X}_4$	
$\mathfrak{X}_5$	$\mathfrak{X}_9$	$\mathfrak{X}_{10}$	$\mathfrak{X}_5$	
$\mathfrak{X}_6$	$\cosh(\mathfrak{s})\mathfrak{X}_9 - \sinh(\mathfrak{s})\mathfrak{X}_3$	$\cosh(\mathfrak{s})\mathfrak{X}_{10} - \sinh(\mathfrak{s})\mathfrak{X}_4$	$\mathfrak{X}_6$	
$\mathfrak{X}_7$	$\cosh(\mathfrak{s})\mathfrak{X}_9 - \sinh(\mathfrak{s})\mathfrak{X}_2$	$\mathfrak{X}_{10}$	$\mathfrak{X}_7$	
$\mathfrak{X}_8$	$\mathfrak{X}_9$	$\cosh(\mathfrak{s})\mathfrak{X}_{10} - \sinh(\mathfrak{s})\mathfrak{X}_2$	$\mathfrak{X}_8$	
$\mathfrak{X}_9$	$\mathfrak{X}_9$	$\cosh(\mathfrak{s})\mathfrak{X}_{10} - \sinh(\mathfrak{s})\mathfrak{X}_1$	$\mathfrak{X}_9$	
$\mathfrak{X}_{10}$	$\cosh(\mathfrak{s})\mathfrak{X}_9 + \sinh(\mathfrak{s})\mathfrak{X}_1$	$\mathfrak{X}_{10}$	$\mathfrak{X}_{10}$	
$\mathfrak{X}_{11}$	$\mathfrak{X}_9$	$\mathfrak{X}_{10}$	$\mathfrak{X}_{11}$	

*Proof.* Here, we use Ibragimov's method. Due to the Table (1),  $\langle \mathfrak{X}_{11} \rangle$  is the center of  $\mathfrak{g}$ , so we need to specify the sub-algebras of

$$\langle \mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_3, \mathfrak{X}_4, \mathfrak{X}_5, \mathfrak{X}_6, \mathfrak{X}_7, \mathfrak{X}_8, \mathfrak{X}_9, \mathfrak{X}_{10} \rangle.$$


$F_i^{\mathfrak{s}} : \mathfrak{g} \rightarrow \mathfrak{g}$  characterized as the linear map  $\mathfrak{X} \mapsto \text{Ad}(\exp(\mathfrak{s}\mathfrak{X}_i).\mathfrak{X})$ , where  $1 \leq i, j \leq 11$ . Some matrices of  $F_i^{\mathfrak{s}}$ ,  $1 \leq i, j \leq 11$ , namely  $M_1^{\mathfrak{s}}$  and  $M_5^{\mathfrak{s}}$ , according to basis  $\{\mathfrak{X}_1, \dots, \mathfrak{X}_{11}\}$

are reported as:

$$M_1^{\mathfrak{s}} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \cos(\mathfrak{s}) & -\sin(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sin(\mathfrak{s}) & \cos(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \cos(\mathfrak{s}) & -\sin(\mathfrak{s}) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \sin(\mathfrak{s}) & \cos(\mathfrak{s}) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \cos(\mathfrak{s}) & -\sin(\mathfrak{s}) & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sin(\mathfrak{s}) & \cos(\mathfrak{s}) & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$M_5^{\mathfrak{s}} = \begin{bmatrix} \cosh(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 & -\sinh(\mathfrak{s}) & 0 & 0 & 0 \\ 0 & \cosh(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 & -\sinh(\mathfrak{s}) & 0 & 0 \\ 0 & 0 & \cosh(\mathfrak{s}) & 0 & \sinh(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sinh(\mathfrak{s}) & 0 & \cosh(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ -\sinh(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 & \cosh(\mathfrak{s}) & 0 & 0 & 0 \\ 0 & -\sinh(\mathfrak{s}) & 0 & 0 & 0 & 0 & 0 & 0 & \cosh(\mathfrak{s}) & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Acting the eleven matrices mentioned above on a generator  $\mathfrak{X} = \sum_{i=1}^{11} a_i \mathfrak{X}_i$  periodically we specify  $\mathfrak{X}$ . In order to clarify the proof, the following two diagrams are given.





- Let  $a_{10} = 0$  and  $a_9 \neq 0$ , for vector (16), the coefficients of  $\mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_3, \mathfrak{X}_6$  and  $\mathfrak{X}_7$  can be disappeared by setting  $\mathfrak{s}_{10} = -\tanh^{-1}(a_1/a_9)$ ,  $\mathfrak{s}_7 = \tanh^{-1}(a_2/a_9)$ ,

$\mathfrak{s}_6 = \tanh^{-1}(a_3/a_9)$ ,  $\mathfrak{s}_3 = \tan^{-1}(a_6/a_9)$  and  $\mathfrak{s}_2 = \tan^{-1}(a_7/a_9)$  respectively. Thus, (16) is reduced to

$$(17) \quad (0, 0, 0, a_4, a_5, 0, 0, a_8, a_9, 0).$$

- Let  $a_{10} = 0$  and  $a_9 = a_4 \neq 0$ , for vector (17), the coefficient of  $\mathfrak{X}_5$  can be disappeared by setting  $\mathfrak{s}_8 = -\tanh^{-1}(a_5/a_4)$ . In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_4 = 1$  and  $a_9 = \pm 1$ . Thus,  $X$  gives rise to case (20).
- Let  $a_{10} = a_4 = 0$  and  $a_9 = a_5 \neq 0$ , for vector (17), the coefficient of  $\mathfrak{X}_8$  can be disappeared by setting  $\mathfrak{s}_4 = -\tan^{-1}(a_8/a_5)$ . In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_5 = 1$  and  $a_9 = \pm 1$ . Thus,  $X$  gives rise to case (21).
- Let  $a_{10} = a_4 = a_5 = 0$  and  $a_9 \neq 0$ , in vector (17). In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_8 = 1$  and  $a_9 = \pm 1$ . Thus,  $X$  gives rise to cases (9) and (26).
- Let  $a_{10} = a_9 = 0$  and  $a_8 \neq 0$ , for vector (16), the coefficient of  $\mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_4, \mathfrak{X}_5$  and  $\mathfrak{X}_7$  can be disappeared by setting  $\mathfrak{s}_7 = \tanh^{-1}(a_1/a_8)$ ,  $\mathfrak{s}_{10} = -\tanh^{-1}(a_2/a_8)$ ,  $\mathfrak{s}_5 = \tanh^{-1}(a_4/a_8)$ ,  $\mathfrak{s}_4 = \tan^{-1}(a_5/a_8)$  and  $\mathfrak{s}_1 = \tan^{-1}(a_7/a_8)$  respectively. Thus, (16) is reduced to

$$(18) \quad (0, 0, a_3, 0, 0, a_6, 0, a_8, 0, 0).$$

- Let  $a_{10} = a_9 = 0$  and  $a_8 = a_3 \neq 0$ , for vector (18), the coefficient of  $\mathfrak{X}_6$  can be disappeared by setting  $\mathfrak{s}_9 = -\tanh^{-1}(a_6/a_3)$ . In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_3 = 1$  and  $a_8 = \pm 1$ . Thus,  $X$  gives rise to case (17).
- Let  $a_{10} = a_9 = a_3 = 0$  and  $a_8 \neq 0$ , in vector (18). In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_6 = 1$  and  $a_8 = \pm 1$ . Thus,  $X$  gives rise to cases (8) and (24).
- Let  $a_{10} = a_9 = a_8 = 0$  and  $a_7 \neq 0$ , for vector (16), the coefficient of  $\mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_3$  and  $\mathfrak{X}_5$  can be disappeared by setting  $\mathfrak{s}_8 = -\tanh^{-1}(a_1/a_7)$ ,  $\mathfrak{s}_9 = -\tanh^{-1}(a_2/a_7)$ ,  $\mathfrak{s}_5 = \tanh^{-1}(a_3/a_7)$ , and  $\mathfrak{s}_3 = \tan^{-1}(a_5/a_7)$  respectively. Thus, (16) is reduced to

$$(19) \quad (0, 0, 0, a_4, 0, a_6, a_7, 0, 0, 0).$$

- Let  $a_{10} = a_9 = a_8 = 0$  and  $a_7 = a_4 \neq 0$ , for vector (19), the coefficient of  $\mathfrak{X}_6$  can be disappeared by setting  $\mathfrak{s}_{10} = -\tanh^{-1}(a_6/a_4)$ . In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_4 = 1$  and  $a_7 = \pm 1$ . Thus,  $X$  gives rise to case (19).
- Let  $a_{10} = a_9 = a_8 = a_4 = 0$  and  $a_7 \neq 0$ , in vector (19). In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_6 = 1$  and  $a_7 = \pm 1$ . Thus,  $X$  gives rise to cases (7) and (23).
- Let  $a_{10} = a_9 = a_8 = a_7 = 0$  and  $a_6 \neq 0$ , for vector (16), the coefficient of  $\mathfrak{X}_2, \mathfrak{X}_3, \mathfrak{X}_4$  and  $\mathfrak{X}_5$  can be disappeared by setting  $\mathfrak{s}_5 = \tanh^{-1}(a_2/a_6)$ ,  $\mathfrak{s}_9 = -\tanh^{-1}(a_3/a_6)$ ,  $\mathfrak{s}_{10} = -\tanh^{-1}(a_4/a_6)$ , and  $\mathfrak{s}_2 = \tan^{-1}(a_5/a_6)$  respectively. Thus, (16) is reduced to

$$(20) \quad (a_1, 0, 0, 0, 0, a_6, 0, 0, 0, 0).$$

In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_1 = 1$  and  $a_6 = \pm 1$ . Thus,  $X$  gives rise to cases (6) and (14).

- Let  $a_{10} = a_9 = a_8 = a_7 = a_6 = 0$  and  $a_5 \neq 0$ , for vector (16), the coefficient of  $\mathfrak{X}_2, \mathfrak{X}_3$  and  $\mathfrak{X}_4$  can be disappeared by setting  $\mathfrak{s}_6 = -\tanh^{-1}(a_2/a_5)$ ,  $\mathfrak{s}_7 = -\tanh^{-1}(a_3/a_5)$  and  $\mathfrak{s}_8 = -\tanh^{-1}(a_4/a_5)$  respectively. Thus, (16) is reduced to

$$(21) \quad (a_1, 0, 0, 0, a_5, 0, 0, 0, 0, 0).$$

In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_1 = 1$  and  $a_5 = \pm 1$ . Thus,  $X$  gives rise to cases (5) and (13).

- Let  $a_{10} = a_9 = a_8 = a_7 = a_6 = a_5 = 0$  and  $a_4 \neq 0$ , for vector (16), the coefficient of  $\mathfrak{X}_1$  and  $\mathfrak{X}_3$  can be disappeared by setting  $\mathfrak{s}_3 = -\tan^{-1}(a_1/a_4)$ , and  $\mathfrak{s}_1 = \tan^{-1}(a_3/a_4)$  respectively. Thus, (16) is reduced to

$$(22) \quad (0, a_2, 0, a_4, 0, 0, 0, 0, 0, 0).$$

In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_2 = 1$  and  $a_4 = \pm 1$ . Thus,  $X$  gives rise to cases (4) and (16).

- Let  $a_{10} = a_9 = a_8 = a_7 = a_6 = a_5 = a_4 = 0$  and  $a_3 \neq 0$ , for vector (16), the coefficient of  $\mathfrak{X}_1$  can be disappeared by setting  $\mathfrak{s}_4 = \tan^{-1}(a_1/a_3)$ . Thus, (16) is reduced to

$$(23) \quad (0, a_2, a_3, 0, 0, 0, 0, 0, 0, 0).$$

In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_2 = 0, 1$  and  $a_3 = \pm 1$ . Thus,  $X$  gives rise to cases (3) and (15).

- Let  $a_{10} = a_9 = a_8 = a_7 = a_6 = a_5 = a_4 = a_3 = 0$ . Thus, (16) is reduced to

$$(24) \quad (a_1, a_2, 0, 0, 0, 0, 0, 0, 0, 0).$$

In order to simplify the phrase, by scaling  $\mathfrak{X}$ , assume that  $a_1 = 0, 1$  and  $a_2 = 0, \pm 1$ . Thus,  $X$  gives rise to cases (1), (2) and (12).

□

#### 4. SOME REDUCED EQUATIONS OF EQ.(4)

Now, we are going to offer a classified symmetry reduction of Eq.(4) regarding sub-algebras of Theorem 3.1. For this purpose, we have to look for a new shape of Eq.(4) in particular coordinates to reduce it. A coordinate like this would be built by realizing independent invariant  $\varsigma, \eta, k, h$  corresponds to the infinitesimal solution. Therefore, representing the problem in other coordinates, utilizing the derivative will reduce the order of PDE. Every 1D sub-algebras in 3.1, the similarity variables  $\varsigma_i, \eta_i, k_i$ , and  $h_i$  are brought in Table 3, where, in cases (16) and (18), one puts  $\alpha = 1$ . Every similarity variable is utilized to reduce Eq.(4) to a new PDE which, we bring in Table 4.

TABLE 3. Lie group and similarity variable.

$i$	$H_i$	$\zeta_i$	$\eta_i$	$t_i$	$w_i$	$u_i$	$v_i$
1	$\mathfrak{X}_1$	$x$	$z$	$u$	$v$	$k(\zeta, \eta)$	$f(\zeta, \eta)$
2	$\mathfrak{X}_2$	$x$	$y$	$u$	$v$	$k(\zeta, \eta)$	$f(\zeta, \eta)$
3	$\mathfrak{X}_1 + a\mathfrak{X}_2$	$x$	$y - \frac{z}{a}$	$u$	$v$	$k(\zeta, \eta)$	$f(\zeta, \eta)$
4	$\mathfrak{X}_1 + a\mathfrak{X}_{11}$	$x$	$z$	$Ln(u) - ay$	$v$	$e^{(ay+k(\zeta, \eta))}$	$f(\zeta, \eta)$
5	$\mathfrak{X}_2 + a\mathfrak{X}_{11}$	$x$	$y$	$Ln(u) - az$	$v$	$e^{(az+k(\zeta, \eta))}$	$f(\zeta, \eta)$
6	$\mathfrak{X}_1 + a\mathfrak{X}_2 + b\mathfrak{X}_{11}$	$x$	$y - \frac{z}{a}$	$Ln(u) - \frac{b}{a}z$	$v$	$e^{(\frac{b}{a}z+k(\zeta, \eta))}$	$f(\zeta, \eta)$
7	$\mathfrak{X}_1 + a\mathfrak{X}_\alpha, \alpha = u$	$x$	$z$	$Ln(u) - ay$	$Ln(E - v) + ay$	$e^{(ay+k(\zeta, \eta))}$	$E - e^{-ay+f(\zeta, \eta)}$
8	$\mathfrak{X}_2 + a\mathfrak{X}_\alpha, \alpha = u$	$x$	$y$	$Ln(u) - az$	$Ln(E - v) + az$	$e^{(az+k(\zeta, \eta))}$	$E - e^{-az+f(\zeta, \eta)}$
9	$\mathfrak{X}_1 + a\mathfrak{X}_\alpha, \alpha = 1$	$x$	$z$	$u - ay$	$uLn(E - v) + ay$	$ay + k(\zeta, \eta)$	$E - e^{(-ay+f(\zeta, \eta))/u}$
10	$\mathfrak{X}_2 + a\mathfrak{X}_\alpha, \alpha = 1$	$x$	$y$	$u - az$	$uLn(E - v) + az$	$az + k(\zeta, \eta)$	$E - e^{(-az+f(\zeta, \eta))/u}$
11	$\mathfrak{X}_1 + a\mathfrak{X}_\alpha + b\mathfrak{X}_{11}, \alpha = u$	$x$	$z$	$Ln(u) - (a+b)y$	$Ln(E - v) + ay$	$e^{(a+b)y+k(\zeta, \eta)}$	$E - e^{-ay+f(\zeta, \eta)}$
12	$\mathfrak{X}_2 + a\mathfrak{X}_\alpha + b\mathfrak{X}_{11}, \alpha = u$	$x$	$y$	$Ln(u) - (a+b)z$	$Ln(E - v) + az$	$e^{(a+b)z+k(\zeta, \eta)}$	$E - e^{-az+f(\zeta, \eta)}$
13	$\mathfrak{X}_1 + a\mathfrak{X}_\alpha + b\mathfrak{X}_{11}, \alpha = 1$	$x$	$z$	$Ln(bu + a) - by$	$uLn(E - v) + ay$	$(e^{(by+k(\zeta, \eta))} - a)/b$	$E - e^{(-ay+f(\zeta, \eta))/u}$
14	$\mathfrak{X}_2 + a\mathfrak{X}_\alpha + b\mathfrak{X}_{11}, \alpha = 1$	$x$	$y$	$Ln(bu + a) - bz$	$uLn(E - v) + az$	$(e^{(bz+k(\zeta, \eta))} - a)/b$	$E - e^{(-az+f(\zeta, \eta))/u}$
15	$\mathfrak{X}_1 + a\mathfrak{X}_2 + b\mathfrak{X}_\alpha + c\mathfrak{X}_{11}, \alpha = u$	$x$	$y - \frac{z}{a}$	$Ln(u) - (b+c)y$	$Ln(E - v) + by$	$e^{(b+c)y+k(\zeta, \eta)}$	$E - e^{-by+f(\zeta, \eta)}$
16	$\mathfrak{X}_1 + a\mathfrak{X}_2 + b\mathfrak{X}_\alpha + c\mathfrak{X}_{11}, \alpha = 1$	$x$	$y - \frac{z}{a}$	$Ln(cu + b) - cy$	$uLn(E - v) + by$	$(e^{(cy+k(\zeta, \eta))} - b)/c$	$E - e^{(-by+f(\zeta, \eta))/u}$
$\vdots$		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$

TABLE 4. Reduced equations based on similarity variable.

$i$	Reduction of equations
1	$k_{\zeta\zeta} + \cot(\zeta)k_\zeta - k_{\eta\eta} + (2m/h^2)(E - f)k = 0,$
2	$k_{\zeta\zeta} + \cot(\zeta)k_\zeta + \csc^2(\zeta)k_{\eta\eta} + (2m/h^2)(E - f)k = 0,$
3	$k_{\zeta\zeta} + \cot(\zeta)k_\zeta + \csc^2(\zeta)k_{\eta\eta} - \frac{1}{a^2}k_{\eta\eta} + (2m/h^2)(E - f)k = 0,$
4	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + a^2\csc^2(\zeta) - k_{\eta\eta} - k_\eta^2 + (2m/h^2)(E - f)e^{k+ay} = 0,$
5	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + \csc^2(\zeta)(k_{\eta\eta} + k_\eta^2) - a^2 + (2m/h^2)(E - f)e^{k+az} = 0,$
6	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + \csc^2(\zeta)(k_{\eta\eta} + k_\eta^2) - \frac{1}{a^2}k_{\eta\eta} - (\frac{1}{a}k_\eta + b)^2 + (2m/h^2)(E - f)e^{k+\frac{b}{a}y} = 0,$
7	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + a^2\csc^2(\zeta) - k_{\eta\eta} - k_\eta^2 + (2m/h^2)e^{f-ay}e^{k+ay} = 0,$
8	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + \csc^2(\zeta)(k_{\eta\eta} + k_\eta^2) - a^2 + (2m/h^2)e^{f-az}e^{k+az} = 0,$
9	$k_{\zeta\zeta} + \cot(\zeta)k_\zeta - k_{\eta\eta} + (2m/h^2)e^{(f-ay)/u}(k + ay) = 0,$
10	$k_{\zeta\zeta} + \cot(\zeta)k_\zeta + \csc^2(\zeta)k_{\eta\eta} + (2m/h^2)e^{(f-az)/u}(k + az) = 0,$
11	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + (a+b)^2\csc^2(\zeta) - k_{\eta\eta} - k_\eta^2 + (2m/h^2)e^{f-ay}e^{k+(a+b)y} = 0,$
12	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + \csc^2(\zeta)(k_{\eta\eta} + k_\eta^2) - (a+b)^2 + (2m/h^2)e^{f-az}e^{k+(a+b)z} = 0,$
13	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + b^2\csc^2(\zeta) - k_{\eta\eta} - k_\eta^2 + (2m/h^2)e^{(f-ay)/u}(e^{k+by} - a) = 0,$
14	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + \csc^2(\zeta)(k_{\eta\eta} + k_\eta^2) - b^2 + (2m/h^2)e^{(f-az)/u}(e^{k+bz} - a) = 0,$
15	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + \csc^2(\zeta)(k_{\eta\eta} + k_\eta^2) - \frac{1}{a^2}k_{\eta\eta} - (\frac{1}{a}k_\eta + \frac{b+c}{a})^2 + (2m/h^2)e^{f-ay}e^{k+(b+c)y} = 0,$
16	$k_{\zeta\zeta} + k_\zeta^2 + \cot(\zeta)k_\zeta + \csc^2(\zeta)(k_{\eta\eta} + k_\eta^2) - \frac{1}{a^2}k_{\eta\eta} - (\frac{1}{a}k_\eta + \frac{c}{a})^2 + (2m/h^2)e^{(f-ay)/u}(e^{k+cy} - b)/c = 0.$

As sample, we calculate the invariants related to  $H_9 := \mathfrak{X}_1 + a\mathfrak{X}_\alpha$ . We integrate the following characteristic expression, assuming  $\alpha(u) = 1$ .

$$\frac{dx}{0} = \frac{dy}{1} = \frac{dz}{0} = \frac{du}{1} = u \frac{dv}{E - v}.$$

Thus, the variables are calculated as:

$$\zeta = x, \quad \eta = z, \quad t = u - ay, \quad w = uLn(E - v) + ay,$$

Putting the obtained variables in Eq.(4), and utilizing derivative yields that, the answer of Eq.(4) is as:

$$u = ay + k(\varsigma, \eta), \quad v = E - e^{(-ay+f(\varsigma,\eta))/u},$$

where  $k(\varsigma, \eta)$  and  $h(\varsigma, \eta)$  satisfies the following reduced equation with 2 variables

$$(25) \quad k_{\varsigma\varsigma} + \cot(\varsigma)k_{\varsigma} - k_{\eta\eta} + (2m/h^2)e^{(f-ay)/u}(k + ay) = 0.$$

Subalgebra  $X_1 + aX_\alpha$  and the reduced Eq.(25) are brought in Tables 3 and 4, by case (9).

### 5. 3D QUANTUM HARMONIC OSCILLATOR ON A SPHERE

As we saw in the introduction, the Schrodinger equation for a harmonic oscillator on a sphere was introduced by:

$$(26) \quad \begin{aligned} u_{zz} = u_{xx} + (\cot x)u_x + (\csc^2 x)u_{yy} \\ + (2m/h^2)(E - \frac{1}{2}q_x x^2 - \frac{1}{2}q_y \sin^4(x)y^2 - \frac{1}{2}q_z z^2)u, \end{aligned}$$

where the particle oscillates on a sphere.

To interact with this equation, it is better to work on the reduction equation of the general form of the Schrodinger equation from Table 4. For this purpose, We select Equation 1 from Table 4 and try to solve it. Obviously, this equation is in terms of two variables, and solving this equation seems simpler than the original Equation (7). So, consider:

$$k_{\varsigma\varsigma} + \cot(\varsigma)k_{\varsigma} - k_{\eta\eta} + (2m/h^2)(E - f)k = 0,$$

where considering the metric (3) on  $S^2 \times R$  and the function  $f$  will become

$$f(\varsigma, \eta) = \frac{1}{2}\hat{q}\varsigma^2 + \frac{1}{2}\bar{q}\eta^2,$$

where  $\hat{q}$  and  $\bar{q}$  are constants. Thus (7) turns into

$$(27) \quad k_{\varsigma\varsigma} + \cot(\varsigma)k_{\varsigma} - k_{\eta\eta} + (2m/h^2)(E - \frac{1}{2}\hat{q}\varsigma^2 - \frac{1}{2}\bar{q}\eta^2)k = 0.$$

**Theorem 5.1.** Assume that  $M(\varsigma)$  and  $N(\eta)$  are functions. If they satisfy the following two separate ODEs:

$$(28) \quad \begin{cases} M_{\varsigma\varsigma} + \cot(\varsigma)M_{\varsigma} - (1/h^2)(c_1 + m\hat{q}\varsigma^2)M = 0, \\ N_{\eta\eta} - (1/h^2)(c_1 - m(\bar{q}\eta^2 - 2E))N = 0, \end{cases}$$

then  $k(\varsigma, \eta) = M(\varsigma)N(\eta)$  is a solution of (27).

*Proof.* It is sufficient to show that  $k(\varsigma, \eta) = M(\varsigma)N(\eta)$  satisfies Equation (27). Evaluating the derivative gives:

$$\begin{cases} k_{\varsigma\varsigma} = M_{\varsigma\varsigma}N, \\ \cot(\varsigma)k_{\varsigma} = \cot(\varsigma)M_{\varsigma}N \\ k_{\eta\eta} = N_{\eta\eta}M, \end{cases}$$

so

$$\begin{aligned}
0 &= k_{\varsigma\varsigma} + \cot(\varsigma)k_{\varsigma} - k_{\eta\eta} + (2m/h^2)(E - \frac{1}{2}\hat{q}\varsigma^2 - \frac{1}{2}\bar{q}\eta^2)k \\
&= M_{\varsigma\varsigma}N + \cot(\varsigma)M_{\varsigma}N - N_{\eta\eta}M + (2m/h^2)(E - \frac{1}{2}\hat{q}\varsigma^2 - \frac{1}{2}\bar{q}\eta^2)MN \\
&= M_{\varsigma\varsigma}N + \cot(\varsigma)M_{\varsigma}N - N_{\eta\eta}M \\
&\quad + (1/h^2)(2mE - m\hat{q}\varsigma^2 - m\bar{q}\eta^2 + c_1 - c_1)MN \\
&= \left( M_{\varsigma\varsigma} + \cot(\varsigma)M_{\varsigma} - (1/h^2)(c_1 + m\hat{q}\varsigma^2)M \right) N \\
&\quad + \left( N_{\eta\eta} - (1/h^2)(c_1 - m(\bar{q}\eta^2 - 2E))N \right) M.
\end{aligned}$$

Because  $M$  and  $N$  are not zero, then the necessary result is obtained.  $\square$

Now to solve Equation (27), we need to consider Equations (28). In this sense, for  $c_1 = -2mE$ , the solution of the second equation of (28), using Maple, is

$$N = C_1 \sqrt{\eta} \text{Bessel}J\left(\frac{1}{4}, \frac{1}{2} \sqrt{\frac{\bar{q}m}{h^2}} \eta^2\right) + C_2 \sqrt{\eta} \text{Bessel}Y\left(\frac{1}{4}, \frac{1}{2} \sqrt{\frac{\bar{q}m}{h^2}} \eta^2\right),$$

where  $C_1, C_2 \in \mathbf{R}$  and BesselJ and BesselY are the Bessel functions of the first and second kinds, respectively. The first equation of (28) with the assumption  $y = \frac{M_{\varsigma}}{\varsigma}$  turns into

$$y_{\varsigma} = -y^2 - \cot(\varsigma)y + \frac{\hat{q}\varsigma^2 m + c_1}{h^2}.$$

This ODE is called Riccati. Indeed, we started with Equation (7) and finally reached the Riccati equation.

## CONCLUSION

We answered a question of a quantum system obeying the Schrodinger equation on a sphere. We set up the Hamiltonian of the system and the corresponding Schrodinger equation. As a result, Equation (6) on a sphere was presented. In this regard, we tried to solve the equation. Decreasing the order of the equation, we present several newer equations with only two variables in Table 4. Using one of the equations in Table 4 for an oscillator object attached to a spring on a sphere, we converted this two-variable PDE into two one-variable ODEs. One of those equations was solved and the other equation is a Riccati equation, which can be discussed in exceptional cases.

## REFERENCES

- [1] S.C. Anco, M.L. Gandarias, E. Recio, Line-solitons, line-shocks, and conservation laws of a universal KP-like equation in 2+1 dimensions, J. Math. Anal. Appl. 504 (2021), 125319.
- [2] Y. AryaNejad, *Symmetry Analysis of Wave Equation on Conformally Flat Spaces*, J. Geom. Phys. 161(2021), 104029.
- [3] Y. AryaNejad, *Exact solutions of diffusion equation on sphere*, Comput. Methods Differ. Equ., 10(3) (2022), 789-798.

- [4] Y. AryaNejad, A. Khalili, *Invariant solutions and conservation laws of time-dependent negative-order (vnCBS) equation*, J. Mahani Math. Res. 2024; 13(2): 155 - 168.
- [5] Y. AryaNejad, R. Mirzavand, *Conservation laws and invariant solutions of time-dependent Calogero-Bogoyavlenskii-Schiff equation*, J. Linear. Topological. Algebra. 11 (4) (2022), 243-252.
- [6] Y. Aryanejad, *Some geometrical properties of the Oscillator group* Caspian Journal of Mathematical Sciences (CJMS), **9**(2), (2020) 266-275.
- [7] Y. Aryanejad, *Harmonicity of left-invariant vector fields on Einstein Lorentzian Lie groups*, Iranian Journal of Mathematical Sciences and Informatics, **15**(1), (2020) 65-78.
- [8] Y. Aryanejad, *Some geometrical properties of Berger Sphere*, Caspian Journal of Mathematical Sciences (CJMS) 10.2 (2021): 183-194.
- [9] G.W. Bluman and J.D. Cole, *Similarity Methods for Differential equations*, Appl. Math. Sci., No. 13, Springer-Verlag, New York, 1974.
- [10] G.W. Bluman and S. Kumei, *Symmetries and Differential Equations*, Springer, New York, 1989.
- [11] M. Jafari, A. Zaeim, M. Gandom *On similarity reductions and conservation laws of the two non-linearity terms Benjamin-Bona-Mahoney equation*, Journal of Mathematical Extension. **17**(7), (2023) 1-22.
- [12] A. P. Márquez, M. S. Bruzón, *Lie point symmetries, traveling wave solutions and conservation laws of a non-linear viscoelastic wave equation*, Mathematics, 9 (17) (2021) 21–31.
- [13] A. P. Márquez, T. M. Garrido, E. Recio, R. de la Rosa, *Lie symmetries and exact solutions for a fourth-order nonlinear diffusion equation*, Math Meth Appl Sci. 2022;45(17):10614-10627.
- [14] M. Nadjafikhah, Y. AryaNejad, N. Zandi, *Conformal Einstein PP-Wave as Quantum Solutions*, Journal of Mathematical Extension. **16**(12)(2022), 1–19.
- [15] M. Nadjafikhah, M. Jafary, *Computation of Partially Invariant Solutions for the Einstein Walker Manifolds*, Communications in Nonlinear Science and Numerical Simulation. 18(12) (2011) 3317-3324.
- [16] P.J. Olver, *Applications of Lie Groups to Differential Equations*, Springer, New York, 1986.
- [17] L.V. Ovsiannikov, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.

YADOLLAH ARYANEJAD: DEPARTMENT OF MATHEMATICS, PAYAME NOOR UNIVERSITY, P.O. BOX 19395-3697, TEHRAN, I.R. OF IRAN.

Email address: y.aryanejad@pnu.ac.ir