

Some Properties of Entropy for the Exponentiated Pareto Distribution (EPD) Based on Order Statistics

J. Behboodian

Islamic Azad University-Shiraz Branch

S. Tahmasebi

Shiraz University

Abstract. In this paper, we derived the exact form of the entropy for Exponentiated Pareto Distribution (EPD). Some properties of the entropy and mutual information are presented for order statistics of EPD. Also, the bounds are computed for the entropies of the sample minimum and maximum for EPD.

AMS Subject Classification: 94A17; 60E05

Keywords and Phrases: Differential entropy, entropy bounds, exponentiated Pareto Distribution, order Statistics, mutual information.

1. Introduction

We will first introduce the concept of differential entropy which is the entropy of a continuous random variable. Let X be a random variable with cumulative distribution function $F_X(x) = P(X \leq x)$ and density

$f_X(x) = F'_X(x)$. The differential entropy $H(X)$ of a continuous random variable X with a density $f_X(x)$ is defined as

$$H(X) = - \int_{-\infty}^{\infty} f_X(x) \log f_X(x) dx = - \int_0^1 \log f_X(F_X^{-1}(u)) du, \quad (1)$$

where $u = F_X(x)$. Now, let us consider the exponentiated Pareto distribution (EPD) with probability density function (pdf)

$$f_X(x) = \theta \lambda \left[1 - (x+1)^{-\lambda} \right]^{\theta-1} (x+1)^{-(\lambda+1)}, \quad x > 0, \quad \lambda > 0, \quad \theta > 0, \quad (2)$$

and cumulative distribution function (cdf)

$$F_X(x) = \left[1 - (x+1)^{-\lambda} \right]^{\theta}, \quad x > 0, \quad \lambda > 0, \quad \theta > 0, \quad (3)$$

where θ and λ are two shape parameters. When $\theta = 1$, the above distribution corresponds to the standard Pareto distribution of the second kind.

Analytical expression for the entropy of univariate continuous distributions are discussed by Cover and Thomas [3], Lazo and Rathie [8], Nadarajah and Zografos [9]. Also, the information properties of order statistics have been studied by a few authors. Among them Wong and Chen [15], Park [10], Ebrahimi et al. [5] provided several results and some characterizations of shannon entropy for order statistics.

The rest of this paper is organized as follows. In Section 2, we derived the exact form of the entropy for exponentiated pareto distribution

(EPD). In Section 3, we present shannon entropy for jth order statistic of EPD, some properties of the entropy, and mutual information for order statistics of EPD.

2. Entropy for EPD

Suppose X is a random variable with $EP(\theta, \lambda)$ and density function (2). Now using (1), the log-density of (2) is

$$\log f_X(x) = \log(\theta\lambda) + (\theta-1) \log \left(1 - (x+1)^{-\lambda}\right) - (\lambda+1) \log(x+1), \quad (4)$$

and the entropy is

$$\begin{aligned} H(X) &= E(-\log f_X(x)) \\ &= - \int_0^\infty f_X(x) \log f_X(x) dx \\ &= -\log(\theta\lambda) + (1-\theta)E \left[\log \left(1 - (x+1)^{-\lambda}\right) \right] \\ &\quad + (\lambda+1)E [\log(x+1)]. \end{aligned} \quad (5)$$

So, we need to find $E [\log (1 - (x+1)^{-\lambda})]$ and $E [\log(x+1)]$ to obtain the Shannon entropy.

Derivation of these two expectations are based on the following strategy:

$$k(r) = E[(x+1)^r] = \int_0^\infty \theta\lambda(x+1)^{r-(\lambda+1)} \left[1 - (x+1)^{-\lambda}\right]^{\theta-1} dx. \quad (6)$$

By the change of variable $[1 - (x + 1)^{-\lambda}] = t$, $0 < t < 1$, we obtain:

$$\begin{aligned}
 k(r) &= E[(x + 1)^r] \\
 &= \int_0^1 \theta(t)^{\theta-1} (1-t)^{-\frac{r}{\lambda}} dx \\
 &= \theta \cdot \frac{\Gamma(\theta)\Gamma(1-\frac{r}{\lambda})}{\Gamma(\theta+1-\frac{r}{\lambda})}, \quad 1 - \frac{r}{\lambda} \neq 0, -1, -2, \dots
 \end{aligned} \tag{7}$$

Differentiating both sides of (7) with respect to r we obtain:

$$\begin{aligned}
 k' &= E[(x + 1)^r \log(x + 1)] \\
 &= \frac{\theta\Gamma(\theta) \left[\frac{-1}{\lambda}\Gamma'(1-\frac{r}{\lambda})\Gamma(\theta+1-\frac{r}{\lambda}) + \frac{1}{\lambda}\Gamma(1-\frac{r}{\lambda})\Gamma'(\theta+1-\frac{r}{\lambda}) \right]}{\left(\Gamma(\theta+1-\frac{r}{\lambda})\right)^2}
 \end{aligned} \tag{8}$$

From relation (8), at $r = 0$ we obtain

$$E[\log(x + 1)] = \frac{1}{\lambda} [\psi(\theta + 1) - \psi(1)], \tag{9}$$

where ψ is the digamma function defined by $\psi(\theta) = \frac{d}{d\theta} \ln \Gamma(\theta)$.

Now we calculate

$$\begin{aligned}
 t(r) &= E\left[\left(1 - (x + 1)^{-\lambda}\right)^r\right] \\
 &= \int_0^\infty \theta\lambda \left(1 - (x + 1)^{-\lambda}\right)^{r+\theta-1} (x + 1)^{-(\lambda+1)} dx \\
 &= \frac{\theta}{\theta + r}
 \end{aligned} \tag{10}$$

$$\left. \frac{dt(r)}{dr} \right|_{r=0} = E\left[\log\left(1 - (x + 1)^{-\lambda}\right)\right] = \frac{-1}{\theta}. \tag{11}$$

Putting (9) and (11) in relation (5) we have:

$$H(X) = -\log(\lambda\theta) + \frac{\lambda + 1}{\lambda} [\psi(\theta + 1) - \psi(1)] + \frac{-1 + \theta}{\theta}, \tag{12}$$

where $-\psi(1) = 0.5772\dots$ is the Euler constant.

3. Some Properties of Entropy Based on Order Statistics EPD

Let X_1, \dots, X_n be a random sample from a distribution $F_X(x)$ with density $f_X(x) > 0$. The order statistics of this sample is defined by the arrangement of X_1, \dots, X_n from the smallest to the largest, by $Y_1 < Y_2 < \dots < Y_n$. The density of Y_j , $j = 1, \dots, n$, is

$$f_{Y_j}(y) = \frac{n!}{(j-1)!(n-j)!} f_X(y) [F_X(y)]^{j-1} [1 - F_X(y)]^{n-j}. \quad (13)$$

Now, let U_1, U_2, \dots, U_n be a random sample from $U(0, 1)$ with the order statistics $W_1 < W_2 < \dots < W_n$. The density of W_j , $j = 1, \dots, n$, is

$$f_{W_j}(w) = \frac{1}{B(j, n-j+1)} w^{j-1} [1-w]^{n-j}, \quad 0 < w < 1, \quad (14)$$

where $B(j, n-j+1) = \frac{\Gamma(j)\Gamma(n-j+1)}{\Gamma(n+1)} = \frac{(j-1)!(n-j)!}{n!}$.

The entropy of the beta distribution is

$$\begin{aligned} H_n(W_j) &= -(j-1) [\psi(j) - \psi(n+1)] - (n-j) \\ &\quad [\psi(n+1-j) - \psi(n+1)] + \log B(j, n-j+1), \end{aligned}$$

where $\psi(t) = \frac{d \log \Gamma(t)}{dt}$, $\psi(n+1) = \psi(n) + \frac{1}{n}$.

Using the fact that $W_j = F_X(Y_j)$ and $Y_j = F_X^{-1}(W_j)$, $j = 1, 2, \dots, n$, are one to one transformations, the entropies of order statistics can be computed by

$$H(Y_j) = H_n(W_j) - E_{g_j} [\log f_X(F_X^{-1}(W_j))] \quad (15)$$

$$= H_n(W_j) - \int f_j(y) \log f_X(y) dy, \quad (16)$$

Now, we can have an application of (16) for the EPD. Let X be a random variable having the EPD(θ, λ). For computing $H(Y_j)$, we have

$$F_X^{-1}(W_j) = \left[1 - (W_j)^{\frac{1}{\theta}} \right]^{-\frac{1}{\lambda}} - 1,$$

and the expectation term in (15) is obtained as follows:

$$\begin{aligned} E_{g_j} [\log f_X(F_X^{-1}(W_j))] &= E_{g_j} [\log(\theta\lambda) + \frac{\lambda+1}{\lambda} \log(1 - (W_j)^{\frac{1}{\theta}}) + \\ &\quad (\theta-1) \log((W_j)^{\frac{1}{\theta}})] \\ &= \log(\theta\lambda) + \frac{\lambda+1}{\lambda} E_{g_j} [\log(1 - (W_j)^{\frac{1}{\theta}})] \\ &\quad + \theta E_{g_j} [\log(W_j)] \\ &= \log(\theta\lambda) + \frac{\lambda+1}{\lambda} \left[\frac{n!}{(j-1)!} \right. \\ &\quad \left. \sum_{k=0}^{n-j} \frac{(-1)^k (\psi(1) - \psi(\theta k + j\theta + 1))}{k!(n-j-k)!(k+j)} \right] \\ &\quad + \frac{\theta-1}{\theta} (\psi(j) - \psi(n+1)). \end{aligned} \quad (17)$$

Therefor, by (15) and (17) the entropy of j th order statistic is

$$\begin{aligned} H(Y_j) &= H_n(W_j) - \log(\theta\lambda) + \frac{\lambda+1}{\lambda} \\ &\quad \left[\frac{n!}{(j-1)!} \cdot \sum_{k=0}^{n-j} \frac{(-1)^k (-\psi(1) + \psi(\theta k + j\theta + 1))}{k!(n-j-k)!(k+j)} \right] \\ &\quad + \frac{1-\theta}{\theta} (\psi(j) - \psi(n+1)). \end{aligned} \quad (18)$$

For the sample minimum $j = 1$, $H_n(W_1) = 1 - \log n - \frac{1}{n}$ and

$$\begin{aligned} H(Y_1) = & 1 - \log n - \frac{1}{n} - \log(\theta\lambda) + \\ & \frac{\lambda+1}{\lambda} \times \left[\sum_{u=1}^n (-1)^{u-1} \binom{n}{u} (\psi(\theta u + 1) + \gamma) \right] \\ & + \frac{\theta-1}{\theta} \left(\psi(n) + \frac{1}{n} + \gamma \right), \end{aligned} \quad (19)$$

where $\gamma = -\psi(1) = 0.5772\dots$ is the Euler constant.

The distribution function of Y_n is $F_n(y) = [1 - (y+1)^{-\lambda}]^{n\theta} I(0, \infty)^{(y)}$ and the density is $f_n(y) = n\theta\lambda [1 - (y+1)^{-\lambda}]^{n\theta-1} (y+1)^{-(\lambda+1)} I(0, \infty)^{(y)}$.

Noting that $H_n(W_n) = 1 - \log n - \frac{1}{n}$, the formula (18) gives

$$\begin{aligned} H(Y_n) = & 1 - \log n - \frac{1}{n} - \log(\theta\lambda) + \\ & \frac{\lambda+1}{\lambda} [\psi(n\theta + 1) + \gamma] + \frac{\theta-1}{\theta} \left(\frac{1}{n} \right). \end{aligned} \quad (20)$$

For any random variable X with $H(X) < \infty$, Ebrahimi et al.[10] showed that the entropy of order statistics Y_j , $j = 1, 2, \dots, n$, is bounded as follow:

$$H(Y_j) \geq H_n(W_j) - \log M, \quad (21)$$

and

$$H(Y_j) \leq H_n(W_j) - \log M + nB_j(H(X) + \log M), \quad (22)$$

where M is the mode of the distribution and B_j denotes the j th term of binomial probability $Bin(n-1, \frac{j-1}{n-1})$. Therefore, we can compute

the bounds for the entropies of the sample minimum and maximum for EPD with parameters λ and θ . We have $M = \left(\frac{\lambda\theta + 1}{\lambda + 1}\right)^{\frac{1}{\lambda}} - 1$. So,

$$1 - \log n - \frac{1}{n} - \log \left(\left(\frac{\lambda\theta + 1}{\lambda + 1} \right)^{\frac{1}{\lambda}} - 1 \right) \leq H(Y_1), \quad (23)$$

and

$$\begin{aligned} H(Y_n) \leq & 1 - \log n - \frac{1}{n} + (n-1) \log \left(\left(\frac{\lambda\theta + 1}{\lambda + 1} \right)^{\frac{1}{\lambda}} - 1 \right) \\ & + n \left(-\log(\lambda\theta) + \frac{\lambda+1}{\lambda} [\psi(\theta+1) - \psi(1)] + \frac{1+\theta}{\theta} \right). \end{aligned} \quad (24)$$

The lower bound and upper bound for the entropies of the sample minimum and maximum for EPD are useful when n is small.

Information theory provides some concepts of extensive use in statistics, one of which is mutual information of two random variables. It is a generalization of the coefficient of determination, for a bivariate random vector (X, Y) with joint density function $f(x, y)$ and marginal density functions, $f_Y(y)$ and $f_X(x)$. The mutual information is defined as

$$\begin{aligned} I(X, Y) &= \int_S f(x, y) \log \frac{f(x, y)}{f_Y(y)f_X(x)} dx dy \\ &= H(X) + H(Y) - H(X, Y), \end{aligned} \quad (25)$$

where S is the region $f(x, y) \geq 0$ and $H(X, Y)$ is the entropy of (X, Y) . Mutual information for order statistics have an important role in statis-

tical sciences. In view of Ebrahimi et al. [10], the degree of dependency among Y_1, \dots, Y_n is measured by the mutual information between consecutive order statistics, defined by

$$I_n(Y_j, Y_{j+1}) = -\log \binom{n}{j} + n\psi(n) - j\psi(j) - (n-j)\psi(n-j) - 1. \quad (26)$$

For given n , $I_n(Y_j, Y_{j+1})$ is symmetric in j and $n-j$; increases in j for $j < \frac{n}{2}$, and decreases for $j > \frac{n}{2}$. $I_n(Y_j, Y_{j+1})$ is increasing in n . Thus, $I_n(Y_j, Y_{j+1})$ is maximum at the median and is symmetric about the median. Now, suppose Y_1, \dots, Y_n denote the order statistics of a random sample X_1, \dots, X_n from EPD, Then we can calculate the mutual information between Y_1 and Y_n . Thus, we have

$$\begin{aligned} I(Y_1, Y_n) &= H(Y_1) + H(Y_n) - H(Y_1, Y_n) \\ &= -\log n(n-1) - \frac{(n-2)}{n(n-1)^2} + 4\left(1 - \frac{1}{n}\right) - 2 - 2\log(\theta\lambda) \\ &+ \frac{\lambda+1}{\lambda} \left[\sum_{u=1}^n (-1)^{u-1} \binom{n}{u} (\psi(\theta u + 1) + \gamma) + \psi(n\theta + 1) + \gamma \right] \\ &+ \frac{\theta-1}{\theta} \left(\psi(n) + \frac{2}{n} + \gamma \right). \end{aligned} \quad (27)$$

Noting that $H(Y_1, Y_n)$ can be computed by

$$\int_0^\infty \int_0^z -f_{Y_1, Y_n}(y, z) \log f_{Y_1, Y_n}(y, z) dy dz.$$

Conclusion

We have derived the exact form of shannon entropy for the Exponentiated Pareto Distribution (EPD) and its order statistics. This distribu-

tion is applied in reliability, actuarial sciences, economics, and telecommunications. We have also presented some properties of the entropy and mutual information for order statistics of EPD.

Acknowledgements

We would like to thank Executive Editor and two referees for their careful reading of our manuscript and helpful suggestions which improved the paper.

References

- [1] B. C. Arnold, N. Balakrishnan, and H. N. Nagaraja, *A first course in order statistics*, Wiley, New York, 1991.
- [2] N. Ahmed and D. V. Gokhale, Entropy expressions and their estimators for multivariate distributions. *IEEE Trans. Inf. Theory IT*, 35 (1989), 688-692.
- [3] T. M. Cover and J. A. Thomas, *Elements of information theory*, Wiley, New York, 1991.
- [4] G. A. Darbellay and I. Vajdo, Entropy expressions for multivariate continuous distribution, *IEEE inf. Theory IT*, 46 (2000), 709-712.
- [5] N. Ebrahimi, E. S. Soofi, and H. Zahedi, Information properties of order statistics and spacing, *IEEE Transactions on Information Theory IT*, 50 (2004), 177-183.
- [6] N. L. Johnson, S. Kotz, and N. Balakrishnan, *Continuous univariate distributions*, Vol. 1, 2nd Edition, Wiley, New York, 1994.
- [7] J. N. Kapur, *Measure of information and their applications*, New York, John Wiley, 1994.
- [8] A. C. Laz, P. N. Rathie, On the entropy of continuous probability distributions, *IEEE Transactions of Information Theory IT*, 24 (1978), 120-122.

- [9] S. Nadarajah and K. Zografos, Formulas for Renyi information and related measures for univariate distributions, *Information Sciences*, 155 (2003), 119-138.
- [10] S. Park, The entropy of consecutive order statistics, *IEEE Transactions of Information Theory IT*, 41 (1995), 2003-2007.
- [11] S. Park, Fisher information on order statistics, *Journal of American Statistical Association*, 91 (1996), 385-390.
- [12] A. Renyi, On measures of entropy and information, *Proc. 4th Berkeley Symposium, Stat. Probability*, 1 (1961), 547-561.
- [13] C. E. Shannon, A mathematical theory of communication, *Bell System Technical Journal*, 27 (1948), 379-432.
- [14] E. S. Soofi, Principal information theoretic approaches, *Journal of the American Statistical Association*, 95 (2000), 1349-1353.
- [15] K. M. Wong and S. Chen, The entropy of ordered sequences and order statistics, *IEEE Transactions of Information Theory IT*, 36 (1990), 276-284.

Javad Behboodian

Department of Mathematics
Islamic Azad University - Shiraz Branch
Shiraz, Iran.
E-mail: behboodian@susc.ac.ir

Saeid Tahmasebi

Department of Mathematics and Statistics
Shiraz University
Shiraz 71454, Iran
E-mail: Stahmasby@yahoo.com