

An Application of Linear Algebra over Lattices

M. Hosseinyazdi

Payame Noor University(PNU)

Abstract. In this paper, first we consider L^n as a semimodule over a complete bounded distributive lattice L . Then we define the basic concepts of module theory for L^n . After that, we proved many similar theorems in linear algebra for the space L^n . An application of linear algebra over lattices for solving linear systems, was given.

AMS Subject Classification: 06DXX; 15A03; 16D10; 06F05.

Keywords and Phrases: Lattices, semimodule, linear algebra, linear system.

1. Introduction

Fuzzy linear systems of equations and inequalities over a bounded chain have been studied by many authors [6], [8], [7]. To extend this concept to L -fuzzy linear systems over a bounded distributive lattice L , we need some basic definitions of linear algebra over lattices such as linearly independent subset, a subsemimodule generated by a set and so on. For more details see [3], [2]. By defining subsemimodule generated by a set, we can find a theoretical necessary and sufficient condition for

consistency of the linear system of equations $A * X = b$ over a bounded distributive lattice.

Definition 1.1. Let $(H, *)$ be a commutative semigroup (monoid) with a reflexive and transitive order \leq on it. $(H, *, \leq)$ is called an ordered commutative semigroup (monoid) if

$$a \leq b \implies a * c \leq b * c \quad \forall a, b, c \in H.$$

Definition 1.2. Let $(H, *)$ be a commutative group (resp. semigroup, monoid) with a partial order \leq . $(H, *, \leq)$ is called a lattice-ordered commutative group (resp. semigroup, monoid), if

$$a \leq b \implies a * c \leq b * c, \quad \forall a, b, c \in H.$$

For simplicity, we call it *l-group* (resp. *l-semigroup*, *l-monoid*).

Example 1.3. Every lattice (L, \leq) is a l-semigroup, by letting $* = \wedge$. Clearly a bounded lattice is a l-monoid in this way.

Definition 1.4. Let $Mat_{n \times m}(L)$ be the set of all $n \times m$ matrices over the lattice (L, \leq) . Define a partial order relation on $Mat_{n \times m}(L)$ as follows:
 $X \leq Y \Leftrightarrow x_{ij} \leq y_{ij};$ for all $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m$,

where $X, Y \in Mat_{n \times m}(L)$. One can see that $(Mat_{n \times m}(L), \leq)$ is a lattice where its supremum and infimum are defined componentwise on $Mat_{n \times m}(L)$ induced by the supremum and infimum of lattice L , respectively.

Definition 1.5 ([10]). Let (R, \oplus) be a commutative monoid with neutral element 0 and (R, \otimes) be a monoid with neutral element 1 where $0 \neq 1$. Then, (R, \oplus, \otimes) is called a semiring with unity 1 and zero 0, if for all $a, b, c \in R$, the following conditions hold:

- (a) $a \otimes (b \oplus c) = (a \otimes b) \oplus (a \otimes c)$,
- (b) $(b \oplus c) \otimes a = (b \otimes a) \oplus (c \otimes a)$,
- (c) $0 = a \otimes 0 = 0 \otimes a$.

Example 1.6. Let L be a bounded distributive lattice. Then, (L, \vee, \wedge) and (L, \wedge, \vee) are semirings.

Definition 1.7 ([10]). $(R, \oplus, \otimes, \leq)$ is called an ordered semiring if

- (a) (R, \oplus, \otimes) is a semiring,
- (b) (R, \oplus, \leq) is an ordered commutative monoid,
- (c) for all $a, b, c, d \in R$,
 - (i) $a \leq b$ and $c \geq 0 \implies a \otimes c \leq b \otimes c$ and $c \otimes a \leq c \otimes b$,
 - (ii) $a \leq b$ and $d \leq 0 \implies a \otimes d \geq b \otimes d$ and $d \otimes a \geq d \otimes b$.

Definition 1.8 ([10]). Let $(H, *, \leq)$ be a commutative ordered monoid with neutral element e and let (R, \oplus, \otimes) be a semiring with unity 1 and zero 0.

Moreover, suppose that $\cdot : R \times H \longrightarrow H$ is a scalar multiplication such that for all $\alpha, \beta \in R$ and for all $a, b \in H$:

- (a) $(\alpha \otimes \beta).a = \alpha.(\beta.a)$,

$$(b) (\alpha \oplus \beta).a = (\alpha.a) \oplus (\beta.a),$$

$$(c) \alpha.(a * b) = (\alpha.a) * (\alpha.b),$$

$$(d) 0.a = e,$$

$$(e) 1.a = a,$$

then, $(R, \oplus, \otimes, H, *, .)$ is called an ordered semimodule over R .

Remark 1.9. Let L be a bounded distributive lattice.

Then, $(L, \vee, \wedge, L, \vee, \wedge)$ and $(L, \wedge, \vee, L, \wedge, \vee)$ are semimodules over

(L, \vee, \wedge) and (L, \wedge, \vee) , respectively.

Upward and downward sets, as important notions in optimization (see [4], [5]), are used in [9] as in the following definition.

Definition 1.10. Let (L, \leq) be a lattice.

(i) A subset $U \subseteq L$ is called upward set if $(a \in U, x \geq a) \implies x \in U$.

(ii) A subset $D \subseteq L$ is called downward set if $(a \in D, x \leq a) \implies x \in D$.

Example 1.11. Let (L, \leq) be a lattice and $a \in L$. Then $\{x \in L | x \geq a\}$ is an upward set and $\{x \in L | x \leq a\}$ is a downward set.

We can easily prove the following proposition.

Proposition 1.12. Let (L, \leq) be a lattice and $M_i \subseteq L$ for $i \in I$. Then $\bigcup_{i \in I} M_i$ is an upward (resp. downward) set if each M_i ; $i \in I$ is upward (resp. downward) set.

2. Basis For Semimodules

In this section we need to extend some basic definition of linear algebra to concepts of lattices. In this case suppose L is a complete distributive lattice and consider L^n as $Mat_{n \times 1}(L)$, the set of all $n \times 1$ matrices over L . By Definition 1.4., L^n is a lattice. Clearly L^n is a distributive complete lattice if L is so. For every bounded distributive lattice L , (L, \vee, \wedge) is a semiring by Example 1.6. and hence (L^n, \wedge, \leq) is a lattice-ordered commutative monoid, by Example 1.3. So we can construct a semimodule as follows.

Theorem 2.1. *Let L be a distributive complete lattice. Then (L^n, \vee, \leq) is a semimodule over (L, \vee, \wedge) .*

Proof. Let L be a bounded distributive lattice. Then (L^n, \vee, \leq) is a semimodule over (L, \vee, \wedge) with scalar multiplication $\bar{\wedge}$ defined by $\bar{\wedge} : L \times L^n \longrightarrow L^n$ such that

$$\alpha \bar{\wedge} \begin{pmatrix} a_1 \\ a_2 \\ \cdot \\ \cdot \\ a_n \end{pmatrix} = \begin{pmatrix} \alpha \wedge a_1 \\ \alpha \wedge a_2 \\ \cdot \\ \cdot \\ \alpha \wedge a_n \end{pmatrix},$$

which for simplification, we write it as \wedge .

In this way (L^n, \vee, \leq) satisfies all conditions of Definition 1.8. Note that

the identity element of (L^n, \vee) is a column matrix which all of its entry are equal to 0. \square

Definition 2.2. Let $(H, *, \leq)$ be a semimodule over semiring (R, \oplus, \otimes) and K be a subset of H such that $(K, *, \leq)$ is a monoid. Then $(K, *, \leq)$ is called a subsemimodule of $(H, *, \leq)$ if it is a semimodule over (R, \oplus, \otimes) and it is denoted by $K \leq_m H$.

The following theorem can be proved easily.

Theorem 2.3. Let $(H, *, \leq)$ be a semimodule over semiring (R, \oplus, \otimes) and K be a subset of H . Then $K \leq_m H$ if and only if

- (i) $e \in K$
- (ii) $x * y \in K$ for all $x, y \in K$,
- (iii) $a.x \in K$ for all $a \in R$, and $x \in K$.

Corollary 2.4. Let L be a distributive complete lattice and K be a sublattice of L which contains 0. Then (K^n, \vee, \leq) is a semimodule over (L, \vee, \wedge) if and only if for every elements $x \in L$ and $y \in K$, we have $x \wedge y \in K$.

Example 2.5. Let $L = \{1, 2, 3, 4, 6, 9, 12, 18, 36\}$ and $x \leq y$ if x divides y . Consider the sublattice $K = \{1, 2, 3, 6\}$. Then, L and K satisfy on Corollary 2.4. Hence (K^n, \vee, \leq) is a semimodule over (L, \vee, \wedge) .

Definition 2.6. Let $(H, *, \leq)$ be a semimodule over (R, \oplus, \otimes) and X be

a subset of H .

(i) The subsemimodule hull of (or subsemimodule generated by) X is the intersection of all subsemimodules of H which contains X and denoted by $\langle X \rangle$. Hence

$$\langle X \rangle = \bigcap_{X \subseteq K \leq H} K.$$

In the other words, $\langle X \rangle$ is the smallest subsemimodule of H which contains X .

(ii) The upward hull of (or upward set generated by) X is defined as the intersection of all upward subsets of H which contains X and is denoted by $\langle X^* \rangle$. So, $\langle X^* \rangle = \bigcap \{K : X \subseteq K \text{ and } K \text{ is an upward subset of } H\}$. In the other words, $\langle X^* \rangle$ is the smallest upward subset of H which contains X .

(iii) The downward hull of (or downward set generated by) X is defined as the intersection of all downward subsets of H which contains X and is denoted by $\langle X_* \rangle$. So, $\langle X_* \rangle = \bigcap \{K : X \subseteq K \text{ and } K \text{ is a downward subset of } H\}$. In the other words, $\langle X_* \rangle$ is the smallest downward subset of H which contains X .

Lemma 2.7. Let $(H, *, \leq)$ be a semimodule over (R, \oplus, \otimes) and $x \in H$.

Then,

(i) $\langle \{x\}^* \rangle = \{a \in H : a \geq x\}$, and

(ii) $\langle \{x\}_* \rangle = \{a \in H : a \leq x\}$.

Definition 2.8. Let $(H, *, \leq)$ be a semimodule over semiring (R, \oplus, \otimes) with scalar multiplication " \cdot " and X be a subset of H . By a linear combination of elements $x_1, \dots, x_m \in X$, we mean $(a_1 \cdot x_1) * \dots * (a_m \cdot x_m)$ where $a_1, \dots, a_m \in R$ and m is a positive integer.

Theorem 2.9. Let $(H, *, \leq)$ be a semimodule over (R, \oplus, \otimes) and X be a subset of H .

(i) Consider $M = \{(a_1 \cdot x_1) * \dots * (a_m \cdot x_m) \mid x_1, \dots, x_m \in X, a_1, \dots, a_m \in R \text{ and } m \text{ is a positive integer}\}$; as the set of all finite linear combinations of elements of X . Then, $\langle X \rangle = M$.

(ii) $\langle X^* \rangle = \bigcup_{x \in X} \langle \{x\}^* \rangle$.

(iii) $\langle X_* \rangle = \bigcup_{x \in X} \langle \{x\}_* \rangle$.

Proof. The proofs of (i)-(iii) follow from Lemma 2.7, Definition 2.8, and Proposition 1.12. \square

Example 2.10. Let $L = [0, 10]$; the bounded chain of real numbers between 0 and 10. Consider semimodule (L^2, \vee, \wedge) over (L, \vee, \wedge) , where \leq is usual partial order on L . For $X_1 = \{(2, 3)^T, (5, 1)^T\}$ the subsemimodule generated by X_1 is shown in Fig. 1.




Fig. 1. Subsemimodule hull of X_1

The upward hull of X_1 is shown in Fig. 2.






Fig. 2. Upward hull of X_1

The downward hull of X_1 is shown in Fig. 3.

Fig. 3. Downward hull of X_1

Now consider $X_2 = \{(2, 4)^T, (5, 9)^T\}$. The subsemimodule hull of X_2 is shown in Fig. 4.

Fig. 4. Subsemimodule hull of X_2

The subsemimodule $\langle X_3 \rangle$, where $X_3 = \{(3, 1)^T, (5, 2)^T, (2, 4)^T\}$,

is as follows:



Fig. 5. Subsemimodule hull of X_3

Definition 2.11. Let $(H, *, \leq)$ be a semimodule over (R, \oplus, \otimes) with zero 0 . A subset X of H is called linearly independent if for all finite subset $\{x_1, \dots, x_m\} \subseteq X$, and elements $a_1, \dots, a_m \in R$; $(a_1.x_1)*\dots*(a_m.x_m) = e$ imply $a_1 = \dots = a_m = 0$.

If the subset X is not linearly independent, it is called linearly dependent.

Example 2.12. Let $L = \{1, 2, 3, 6\}$ and $x \leq y$ means that x divides y . Clearly (L, \vee, \leq) is a semimodule over (L, \vee, \wedge) with zero 1 . Since $2 \wedge 3 = 1$, the set $\{3\}$ is not linearly independent.

Remark 2.13. By the previous example, it is not true that if $x \neq 0$ then $\{x\}$ is linearly independent. But if L is a chain, then for every

non-zero element x , the set $\{x\}$ is linearly independent.

Definition 2.14. Let $(H, *, \leq)$ be a semimodule over (R, \oplus, \otimes) . A linearly independent subset B of H is called a basis for H over R , if $\langle B \rangle = H$.

Example 2.15. Let L be as in Example 2.5. (see Fig. 6).

In this lattice the following subsets of L are linearly independent:

$$K_1 = \{6\}, \quad K_2 = \{6, 12\}, \quad K_3 = \{12, 18\}$$

$$K_4 = \{6, 12, 36\}, \quad K_5 = \{6, 12, 18, 36\}$$

But the following subsets are linearly dependent:

$$K_6 = \{9\}, \quad K_7 = \{2, 3\}, \quad K_8 = \{4, 9\}, \quad K_9 = \{6, 9\}$$

Some sublattices generated by above subsets of L are as follows:


$$\langle K_9 \rangle = \{1, 2, 3, 6, 9, 18\}, \quad \langle K_3 \rangle = \langle K_4 \rangle = \langle K_5 \rangle = \langle K_8 \rangle = L,$$

$$\langle K_6 \rangle = \{1, 3, 9\}$$

Clearly K_3, K_4 and K_5 are bases of L . Also

$$\langle (K_9)_* \rangle = \{1, 2, 3, 6, 9\}, \quad \langle K_9^* \rangle = \{6, 9, 12, 18, 36\}$$

$$\langle (K_5)_* \rangle = L, \quad \langle K_5^* \rangle = K_5.$$



Theorem 3.1. *Let L be a bounded distributive lattice. Consider (L^n, \vee, \leq) as a semimodule over semiring (L, \vee, \wedge) with scalar multiplication " \wedge ". Let A, X and b be $m \times n$, $n \times 1$ and $m \times 1$ matrices over L , respectively. The linear system $A \vee X = b$ has a solution if and only if b belongs to the subsemimodule generated by columns of A .*

Proof. If we show the columns of A by A_1, A_2, \dots, A_n ; then the linear system $A \vee X = b$ can shown by

$$(x_1 \wedge A_1) \vee (x_2 \wedge A_2) \vee \dots \vee (x_n \wedge A_n) = b$$

and clearly the linear system has a solution if and only if $b \in \langle \{A_1, \dots, A_n\} \rangle$ by Theorem 2.9. \square

Example 3.2. Let L, K_9 and K_8 be as in Example 2.15. consider the linear equation

$$(6 \wedge x_1) \vee (9 \wedge x_2) = 3 \tag{1}$$

Then the set of all solutions of (1) is

$$\{(1, 3)^T, (1, 6)^T, (1, 12)^T, (3, 1)^T, (3, 3)^T, (3, 6)^T, (3, 12)^T, (3, 2)^T, (3, 4)^T, (9, 1)^T, (9, 3)^T, (9, 6)^T, (9, 12)^T, (9, 2)^T, (9, 4)^T\}.$$

Linear equation (1) has solution since $3 \in \langle K_9 \rangle$; the subsemimodule generated by $\{6, 9\}$. But if we change right hand side of (1) to 12 we have:

$$(6 \wedge x_1) \vee (9 \wedge x_2) = 12 \tag{2}$$

Clearly (2) doesn't have any solution since $12 \notin K_9$. Now consider

$$(4 \wedge x_1) \vee (9 \wedge x_2) = b \quad (3)$$

Since $\langle \{4, 9\} \rangle = \langle K_8 \rangle = L$, so (3) has solution for all $b \in L$.

Remark 3.3. *Note that Theorem 3.1. gives a theoretical necessary and sufficient condition for consistency of (*). A computational necessary and sufficient condition for consistency of (*) over a bounded chain was given in [6]. Finding such a condition(s) over a bounded distributive lattice is still an open problem.*

References

- [1] G. Grätzer, *General lattice theory*, Academic Press, New York San Francisco, 1978.
- [2] M. Hosseinyazdi, The optimization problem over a distributive lattice, *Journal of Global Optimization*, 41 (2008), 283-298.
- [3] M. Hosseinyazdi, A. Hassankhani and M. Mashinchi, Linear systems and optimization over lattices, *International Review of Fuzzy Mathematics*, 2 (1)(2007), 27-50.
- [4] H. Mohebi, Downward sets and their best simultaneous approximation properties with applications, *Numer. Funct. Anal. Optim.*, 25 (7-8) (2004), 685-705.
- [5] H. Mohebi and A. Rabinov, Best approximation by downward sets with applications, *Anal. Theory Appl*, 22 (1) (2006), 1-22.
- [6] K. Peeva, Fuzzy linear systems, *Fuzzy Set and Systems*, 49 (1992), 339-355.

- [7] K. Peeva, Universal algorithm for solving fuzzy relational equations, *Ital. J. Pure Appl. Math.* 19 (2006), 9-20.
- [8] K. Peeva and Y. Kyosev, Fuzzy relational calculus, *Advances in fuzzy systems-applications and theory*, Vol. 22, World scientific publishing Co. Pte. Ltd, Singapore, 2004.
- [9] I. Singer, *Abstract convex analysis*, John Wiley & Sons, INC. New York, 1997.
- [10] U. Zimmermann, *Linear and combinatorial optimization in ordered algebraic structures*, North-Holland Publishing Company, Amesterdam, 1981.

Mahboobeh Hosseinyazdi
Department of Mathematics
Payame Noor University(PNU)
Shiraz, Iran
E-mail: myazdi@spnu.ac.ir