Journal of Mathematical Extension Vol. 3, No. 2 (2009), 1-15

# C\*-Algebra of Cancellative Semigroupoids

### B. Tabatabaie

Shiraz University

Abstract. In this paper the definition and some properties of semigroupoids are considered. Representations, tight representations, and universal representations of a cancellative semigroupoid are discussed. Also, the  $C^*$ -algebra of a semigroupoid is introduced and it is shown that source elements transfer to zero by tight representations.

AMS Subject Classification: 46L05Keywords and Phrases:  $C^*$ -algebra, projection, partial isometry, graph theory and representation.

## 1. Introduction

Because of the immensity of the class of all  $C^*$ -algebras it has become important to identify and study special types of  $C^*$ -algebras. The theory of  $C^*$ -crossed products by group actions, specially group  $C^*$ -algebras,  $C^*(G)$ , and reduced group  $C^*$ -algebras,  $C_r^*(G)$ , are very well developed. In 1982, J. R. Wordingham proved that the left regular representation of  $\ell^1(S)$  on  $\ell^2(S)$  is faithful ([11]). Following Wordingham,  $C^*$ -algebras of an inverse semigroup, has been investigated by Duncan and Paterson as a generalization of crossed product of discrete groups ([2, 3, 8]).

1

The notion of partial crossed product of a  $C^*$ -algebra by a discrete group is introduced by R. Exel ([4]) and generalized by McClanahan ([7]). Nándor Sieben in his master thesis, at the Arizona State University, under the supervision of J. Quigg defined the  $C^*$ -crossed product by action of an inverse semigroup and published the results in ([9]).

Partial actions of groups and actions of inverse semigroups have been studied by R. Exel in ([5]), where an inverse semigroup, S(G), is associated to a given group G. R. Exel in ([5]) proved that there is a one-to-one correspondence between actions of S(G) and partial actions of G. Also, he introduced a "partial" version of the group  $C^*$ -algebra, that is, partial group  $C^*$ -algebra,  $C^*_P(G)$ . Partial inverse semigroup  $C^*$ -algebra is introduced in ([10]). Now, following ([6]) we will consider the  $C^*$ -algebra of a cancellative semigroupoid.

The organization of this paper is as follows:

Semigroupoids and its properties are considered in Section 2. Section 3 is devoted to the representations, tight representations, universal representations, the  $C^*$ -algebra of a cancellative semigroupoid; and it is shown that source elements transfer to zero by tight representation.

### 2. Semigroupoids

In this section the concepts of semigroupoid, cancellative semigroupoid, divisibility, and source element of a semigroupoid are introduced. An equivalence relation is defined on a special subset of a given semigroupoid. Also, it is shown that the disjoint union of the quotient space of this equivalence relation with the given semigroupoid is a new semigroupoid which has no source.

Let G be a non-empty set and  $G^{(2)}$  be a special subset of  $G \times G$ , that is,  $G^{(2)}$  is the set of all ordered pairs on which a kind of multiplication is meaningful. With this in mind we have the following definition.

**Definition 2.1.** By a semigroupoid G we shall mean a triple  $(G, G^{(2)}, .)$  such that

$$:: G^{(2)} \longrightarrow G$$

is an associative binary operation in the following sense:

For given  $x, y, z \in G$  if either (i) $(x, y) \in G^{(2)}$  and  $(y, z) \in G^{(2)}$ , or (ii)  $(x, y) \in G^{(2)}$  and  $(xy, z) \in G^{(2)}$ , or (iii)  $(y, z) \in G^{(2)}$  and  $(x, yz) \in G^{(2)}$ , then all of (x, y), (y, z), (xy, z) and (x, yz) are in  $G^{(2)}$  and x(yz) = (xy)z.

**Example 2.2.** Let  $E = (E^1, E^0, r, s)$  be a graph. Then the path space of  $E, F^+(E)$ , consists of all finite paths including the vertices, is a semigroupoid with a product xy if s(x) = r(y). In particular, x = xs(x) = r(x)x.

Before we give the definition of *divisibility* we need to know that:

If a semigroupoid, say G, has not a unit element it is possible to add a unit element to it. That is, to pick some element from the universe of outside of G, call it 1, and set  $\tilde{G} = G \dot{\cup} \{1\}$ . Obviously, 1x = x1 = x for every x in G.

It should be noted that  $\tilde{G}$  may not be a semigroupoid. Because if it is a semigroupoid, since for given x, y in  $\tilde{G}$  the products x1 and 1yare meaningful we have (x, 1) and (1, y) are elements of  $G^{(2)}$ . By the Definition 2.1 we conclude that xy = (x1)y is a meaningful product and we know that it is not always the case.

For given x in  $\tilde{G}$ , we would like to determine the set of all elements of G, say y, for which xy is meaningful. Therefore we have

$$G^x = \{y \in G : (x, y) \in G^{(2)}\}$$
 and  $G^1 = G$ .

**Definition 2.3.** Let  $\tilde{G}$  be a unital semigroupoid and  $x, y \in G$ . We shall say that x divids y or y is a multiple of x, in symbols x|y, if there exists z in  $\tilde{G}$  such that  $(x, z) \in G^{(2)}$  and y = xz.

**Lemma 2.4.** The divisibility relation is reflexive, transitive, and invariant under multiplication on the left.

**Proof.** Let  $x \in G$ . Since 1x = x1 = x, we see that the relation is reflexive. To prove the transitivity, if x, y, z are in G such that x|y and y|z we should show that x|z. If x = y from y|z we conclude that x|z.

Similarly if y = z, the relation x|y shows that x|z. Otherewise from x|y we have a u in G such that  $(x, u) \in G^{(2)}$  and y = xu. Also, by y|z we conclude that there exists v in G such that  $(y, v) \in G^{(2)}$  and z = yv. Since  $(x, u), (y, v) \in G^{(2)}$ , that is  $(x, u), (xu, v) \in G^{(2)}$  we see that  $(u, v) \in G^{(2)}$ . Consequently, z = yv = (xu)v = x(uv). This shows that x|z.

To prove the last part of the lemma, let  $x, y, k \in G, x|y, (k, x) \in G^{(2)}$ and  $(k, y) \in G^{(2)}$ . We should show that ky is a multiple of kx. From x|ywe conclude that there exists u in G such that  $(x, u) \in G^{(2)}$  and xu = y. Since (k, x) and (x, u) are elements of  $G^{(2)}$  we see that  $(kx, u) \in G^{(2)}$  and  $(k, y) = (k, xu) \in G^{(2)}$ . As a consequence we have (kx)u = k(xu) = ky, that is, kx|ky. This completes the proof.  $\Box$ The following important concept is pivotal in our work.

**Definition 2.5.** We shall say that an element  $x \in G$  is cancellative if for every  $y, z \in G$  the equation xy = xz implies y = z. If every element of G is cancellative, then G is called a cancellative semigroupoid. Some elements of G has special properties, that is, given  $x \in G$ , there exists  $y \in G$  such that xy is not a legal multiplicative. Here, we would like to introduce the set of all such elements.

**Definition 2.6.** An element x of G is called source if  $G^x = \phi$ . If  $G^x \neq \phi$ , then it is called the multiplicative set of x.

Here, we make an attempt to introduce a semigroupoid without sources.

**Theorem 2.7.** If G is a semigroupoid which has sources, then there exists a semigroupoid which has no source and contains G.

**Proof.** Let  $G^0 = \{x \in G : x \text{ is a source }\}$ . Also, let

$$\psi: G \longrightarrow G$$

defined by  $\psi(x) = e'_x$  be a one-to-one map, and  $E' = \psi(G)$ . For any source y and any x such that  $y \in G^x$ , we observe that if  $t \in G^y$ , that is,  $(y,t) \in G^{(2)}$  then  $(xy,t) \in G^{(2)}$ . This shows that  $G^x \subseteq G^{xy}$ . On the other hand if  $s \in G^{xy}$ , that is,  $(xy,s) \in G^{(2)}$  then  $(y,s) \in G^{(2)}$ . So,  $s \in G^y$  and  $G^{xy} \subseteq G^y$ . Consequently  $G^y = G^{xy}$ , and we conclude that xy is also a source.

Let "~" be any equivalence relation on E' such that  $e'_{xy} \sim e'_y$  for any source y and any x for which  $y \in G^x$ . Also, let  $e_x = [e'_x] = \{t \in E' : x \sim t\}$ , and the quotient space,  $\frac{E'}{\sim}$ , be denoted by E. Take  $\Gamma = G \dot{\cup} E$ ,

$$\Gamma^{(2)} = G^{(2)} \cup \{(y, e_y) : y \in G^0\} \cup \{(e_y, e_y) : y \in G^0\}, \text{ and }$$

define the multiplication

$$:: \Gamma^{(2)} \longrightarrow \Gamma$$

which is nothing but the multiplication on G when restricted to  $G^{(2)}$ , with

$$y.e_y = e_y$$
,  $e_y.e_y = e_y$   $\forall y \in G^0$ .

Now we can prove that  $(\Gamma, \Gamma^{(2)}, .)$  is a semigroupoid which contains G and has no source. To show this, let  $r, s, t \in \Gamma$ . If  $r, s, t \in G$  it is finished, otherwise  $r = e_x$ ,  $s = e_y$  and  $t = e_z$  for some  $x, y, z \in G^0$ . Case 1. If  $e_x = y$  and  $e_z = e_y$ , then

$$(r,s) = (e_x, e_y) = (y, e_y) \in \Gamma^{(2)}$$
, and  $(s,t) = (e_y, e_z) = (e_y, e_y) \in \Gamma^{(2)}$ .

That is, (r, s) and  $(s, t) \in \Gamma^{(2)}$  and by Definition 2.1 part (i) we conclude that  $\Gamma$  is a semigroupoid.

Proofs of other cases are similar to the proof of case 1 and is left to the reader.  $\Box$ 

## 3. Representations of Semigroupoids

In this section the notion of representation of a semigroupoid is introduced. Also, a universal  $C^*$ -algebra is associated to a cancellative semigroupoid. The concept of a *tight representation* and the fact that a source element transfres to the zero operator by a tight representation are discussed.

Throught this section, G is a semigroupoid and A is a unital  $C^*$ -algebra.

**Definition 3.1.** Let  $x, y \in G$ . We shall say that x and y intersect if they have a common multiple, that is, if there exists an element m of Gsuch that x|m and y|m. The fact that x and y are intersect is denoted by  $x \cap y$ . Otherwise we will say that x and y are disjoint and is denoted by  $x \perp y$ .

The next concept is crucial in understanding the definition of a tight representation.

**Definition 3.2.** If X is any subset of G and  $Z \subseteq X$ , then Z is called a covering of X if for every  $x \in X$ , there exists  $h \in Z$  such that x and h are intersect.

The next definition is the first step in bridging semigroupoids and operator algebras.

**Definition 3.3.** By a representation of G in A we mean a mapping

$$\Pi: G \to A$$

such that  $\Pi(x) = \Pi_x$  is a partial isometry and if  $x, y \in G$  then

$$\Pi_x \Pi_y = \begin{cases} \Pi_{xy}, & if(x,y) \in G^{(2)}, \\ 0, & otherwise. \end{cases}$$

Moreover the initial projections  $Q_x = \Pi_x^* \Pi_x$ , and the final projections  $P_y = \Pi_y \Pi_y^*$  should commute among themselves and satisfy to the following conditions:

- (i) $P_x P_y = 0$ , if  $x \perp y$ ;
- (ii)  $Q_x P_y = P_y$  , if  $(x, y) \in G^{(2)}$  ;
- (iii)  $Q_x P_y = 0$ , if  $(x, y) \notin G^{(2)}$ .

It should be noted that any representation extends to  $\tilde{G}$  by taking  $\Pi(1) = \Pi_1 = 1$  and  $Q_1 = P_1 = 1$ . Here we are able to present the reason why we choose cancellative semigroupoid.

If G is not a cancellative semigroupoid, that is, there exists x in G such that for a distinct pair of elements  $y, z \in G$  we have xy = xz. For given representation  $\Pi$ , since  $\Pi_x$  is a partial isometry we have

$$\Pi_y = \Pi_y \Pi_y^* \Pi_y = (\Pi_y \Pi_y^*) \Pi_y = P_y \Pi_y = Q_x P_y \Pi_y$$

$$= \Pi_x^* \Pi_x \Pi_y \Pi_y^* \Pi_y = \Pi_x^* \Pi_x \Pi_y = \Pi_x^* (\Pi_x \Pi_y) = \Pi_x^* \Pi_{xy}$$

And,

$$\Pi_z = \Pi_z \Pi_z^* \Pi_z = (\Pi_z \Pi_z^*) \Pi_z = P_z \Pi_z =$$

$$Q_x P_z \Pi_z = \Pi_x^* \Pi_x \Pi_z \Pi_z^* \Pi_z = \Pi_x^* \Pi_x \Pi_z = \Pi_x^* (\Pi_x \Pi_z) = \Pi_x^* \Pi_{xz}$$

Since xz = xy we have  $\Pi_{xy} = \Pi_{xz}$ , that is  $\Pi_y = \Pi_z$  whereas  $y \neq z$ . This shows that if G is not a cancellative semigroupoid, then we may have  $\Pi(y) = \Pi(x)$  for some x, y such that  $x \neq y$ .

Before we present the definition of a *tight* representation we need to know some more about representations.

For given  $x \in G$  and  $z \in G^x$ , since  $(x, z) \in G^{(2)}$  we know that the initial projection,  $Q_x = \Pi_x^* \Pi_x$ , and the final projection,  $P_z = \Pi_z \Pi_z^*$ , commute and  $Q_x P_z = P_z$ . Also, we know that  $Q_x P_z = P_z$  is equivalent to  $P_z \leq Q_x$ . So, if  $z_1, z_2 \in G^x$  we have  $P_{z_1} \leq Q_x$  and  $P_{z_2} \leq Q_x$ . Consequently  $P_{z_1} \vee P_{z_2} \leqslant Q_x$ , and if H is a finite subset of  $G^x$  we have

$$\bigvee_{z \in H} P_z \leqslant Q_x.$$

If  $y \in \tilde{G}$  and  $z \in G - G^y$  then  $(y, z) \notin G^{(2)}$ , hence  $Q_y P_z = P_z Q_y = 0$ . Therefore, from  $P_z = P_z$  we have  $P_z = P_z - P_z Q_y = P_z(1 - Q_y)$  which is equivalent to  $P_z \leq 1 - Q_y$ . Since z is an arbitrary element of  $G - G^y$ we conclude that if H is a finite subset of  $G - G^y$ , then we have

$$\bigvee_{z \in H} P_z \leqslant 1 - Q_y$$

Now for given finite subsets X, Y of G, let

$$G^{X,Y} = (\bigcap_{x \in X} G^x) \cap (\bigcap_{y \in Y} G - G^y).$$

If  $z \in G^{X,Y}$ , then from  $z \in \bigcap_{x \in X} G^x$  we conclude that  $P_z \leq Q_x$  for all  $x \in X$  and as a consequence

$$P_z \leqslant \prod_{x \in X} Q_x. \tag{1}$$

Also, from  $z \in \bigcap_{y \in Y} G - G^y$  we have  $P_z \leq 1 - Q_y$  for all  $y \in Y$ , and consequently

$$P_z \leqslant \prod_{y \in Y} (1 - Q_y). \tag{2}$$

From (1) and (2), for given  $z \in G^{X,Y}$  we have

$$P_z \leqslant \prod_{x \in X} Q_x \prod_{y \in Y} (1 - Q_y).$$

Also, for given finite subset H of  $G^{X,Y}$ , we conclude that

$$\bigvee_{z \in H} P_z \leqslant \prod_{x \in X} Q_x \prod_{y \in Y} (1 - Q_y)$$

With this in mind we can present the following important definition.

**Definition 3.4.** A representation  $\Pi$  of G in A is said to be tight if for every subsets X, Y of  $\tilde{G}$  and every covering H of  $G^{X,Y}$  the following equality holds

$$\bigvee_{z \in H} P_z = \prod_{x \in X} Q_x \prod_{y \in Y} (1 - Q_y).$$

It should be noted that if no such covering exists, then any representation is tight vacuously.

Before we present the definition of the  $C^*$ -algebra of a semigroupoid we need to introduce the concept of a universal  $C^*$ -algebra.

In recent years, universal constructions play a crucial role in the theory of operator algebras, specially in the theory of  $C^*$ -algebras. In other words, many important  $C^*$ -algebras can be expressed as universal  $C^*$ -algebras generated by a given set and a set of relations which satisfy in certain conditions. In what follows we will describe that, what do we mean by a universal  $C^*$ -algebra generated by a set and a set of relations.

Suppose a set  $B = \{b_i : i \in \Omega\}$  of generators and a set R of relations are given. It should be noted that the relations can be of a very general

nature. Usually, some algebraic relations between generators and their adjoints exist. The only restriction on the relation is that:

(i) they must be realizable among operators on a Hilbert space.

(ii) each generator should have an upper bound when realized as an operator.

A representation of (B|R) is a set  $\{T_i : i \in \Omega\}$  of bounded operators on a Hilbert space H which satisfying in the given relations. Each such representation of (B|R) defines a \*-representation of the free \*-algebra  $\mathcal{A}$  on the set B. For given  $x \in \mathcal{A}$ , let

 $||x|| = \sup\{||\Pi(x)|| : \Pi \text{ is a representation of } (B|R)\}.$ 

This supremum defines a  $C^*$ -seminorm on  $\mathcal{A}$  provided that it is finite. If the elements of seminorm 0 are divided out, then the completion of  $\mathcal{A}$  is called the *universal*  $C^*$ -algebra generated by B, R, or the *universal*  $C^*$ -algebra on (B|R), and is denoted by  $C^*(B|R)$ .

**Example 3.5.** Let  $B = \{x\}$  and  $R = \{x = x^*, ||x|| < 1\}$ . Then  $C^*(B|R)$  is the universal  $C^*$ -algebra generated by a single self-adjoint element of norm 1.

Note that there is no universal  $C^*$ -algebra generated by a single selfadjoint element, because there is no bound on the norm of the element. For more on universal  $C^*$ -algebras, see Chapter II of [1].

Here, we introduce a universal  $C^*$ -algebra which contains the  $C^*$ -algebra

of semigroupoid.

**Definition 3.6.** Let G be a semigroupoid,  $B = {\Pi_x}_{x\in G}$  be a family of partial isometries and R be the set of all relations such that the correspondence  $x \to \pi_x$  is a tight representation of G. The unital universal  $C^*$ -algebra generated by B, R, that is,  $C^*(B|R)$  denoted by  $\tilde{O}(G)$ . In order to give the definition of the  $C^*$ -algebra of a semigroupoid G we need to know that, what do we mean by the universal representation of G?

**Definition 3.7.** A collection of partial isometries,  $\{\Pi_x\}_{x\in G}$ , such that the correspondence  $x \to \Pi_x$  is a tight representation of G is called the universal representation of G.

Now, we are ready to present the definition of the  $C^*$ -algebra of a semigroupoid.

**Definition 3.8.** The closed \*-subalgebra of  $\tilde{O}(G)$  which is generated by the range of the universal representation of G denoted by O(G), is the  $C^*$ -algebra of G.

We close this section by the following important theorem.

**Theorem 3.9.** If  $\Pi$  is a tight representation of a semigroupoid, G and  $x \in G$  is a source element, then  $\Pi_x = 0$ .

**Proof.** Since x is a source element, we have  $G^x = \phi$ . From the fact

that empty set is a covering for  $G^x$ , we conclude that  $Q_x = 0$ . Since  $Q_x = \prod_x^* \prod_x$ , we have  $\prod_x = 0$ .  $\Box$ 

### References

- B. Blackadar, Operator Algebras, Theory of C\*-Algebras and Von Neumann Algebras, Springer-Verlag, Berlin Heidelberg, 2006.
- [2] J. Duncan and A. L. T. Paterson, C\*-algebras of inverse semigroups, Proc. Roy. Soc. Edinburgh Math. Soc., 28 (1985), 41-58.
- [3] J. Duncan and A. L. T. Paterson, C\*-algebras of Clifford semigroups, Proc. Edinburgh Math. Soc. A, 111(1989), 129-145.
- [4] R. Exel, Circle actions on C\*-algebras, Partial automorphisms and a generalized Pimsner-Voiculescu exact sequence, J. Funct. Anal., 122 (1994), 361-401.
- [5] R. Exel, Partial actions of groups and actions of inverse semigroups, Proc. Amr. Math. Soc., 126 (12) (1998), 3481-3494.
- [6] R. Exel, Inverse semigroups and combinatorial C\*-algebras, Bull. Braz. Math. Soc., 39(2) (2008), 191-313.
- [7] K. McClanahan, K-theory for partial crossed products by discrete groups, J. Funct. Anal., 130 (1995), 77-117.
- [8] A. L. Paterson, Weak containment and Clifford semigroups, Proc. Roy. Soc. Edinburgh sect. A, 81 (1987), 23-30.
- [9] N. Sieben, C<sup>\*</sup>-crossed products by partial actions and actions of inverse semigroups, J. Australian Math. Soc., 63 (1997), 32-46.
- [10] B. Tabatabaie Shourijeh, Partial inverse semigroup C\*-algebra, Taiwanese J. of Math., Vol. 1, No. 6 (2006), 1539-1548.
- [11] J. R. Wordingham, The left regular \*-rpresentation of an inverse semigroup, Proc. Amer. Math. Soc., 86 (1982), 55-58.

### Bahman Tabatabaie

Department of Mathematics College of Sciences Shiraz University Shiraz 71454, Iran. E-mail: tabataba@math.susc.ac.ir