

Eigenfunctions of the Weighted Composition Operators

H. Rezaei

Yasouj University

Abstract. In the present paper, we characterize the eigenfunctions of a weighted composition operator on space of holomorphic function on the unit disk.

AMS Subject Classification: 47B33; 47B38.

Keywords and Phrases: Denjoy-Wolff point, linear-fractional model theorem, weighted composition operator.

1. Introduction

A weighted composition operator $C_{\varphi,\psi}$ is an operator that maps $f \in H(\mathbb{U})$, the space of holomorphic functions on the unit disk \mathbb{U} , into $C_{\varphi,\psi}(f)(z) = \varphi(z)f(\psi(z))$, where φ and ψ are analytic functions defined in \mathbb{U} such that $\psi(\mathbb{U}) \subseteq \mathbb{U}$. When $\varphi \equiv 1$, we just have the composition operator C_ψ defined by $C_\psi(f) = f \circ \psi$.

The eigenfunctions of a composition operator on the classical Hardy space H^2 , induced by a hyperbolic disk automorphism, are considered in [2, 4, 5] where it has been shown that many eigenfunctions of a composition operator can be found in the doubly cyclic subspace generated by special functions in H^2 .

Studying the eigenfunctions of weighted composition operators entails a study of the iterate behavior of holomorphic self maps. The holomorphic self maps of \mathbb{U} are divided into classes of elliptic and non-elliptic type. The elliptic type is an automorphism and has a fixed point in \mathbb{U} . It is

well known that this map is conjugate to a rotation $z \rightarrow \lambda z$ for some complex number λ with $|\lambda| = 1$. The iterate of a non-elliptic map can be characterized by the Denjoy-Wolff Iteration Theorem [1, 6, 7]. In the following notation " \xrightarrow{k} " means uniformly converges on compact subsets of \mathbb{U} and ψ_n denotes the composition of ψ with itself n -times.

Denjoy-Wolff Iteration Theorem. *Suppose ψ is a holomorphic self-map of \mathbb{U} that is not an elliptic automorphism. Then*

- (i) *If ψ has a fixed point $w \in \mathbb{U}$, then $\psi_n \xrightarrow{k} w$ and $|\psi'(w)| < 1$.*
- (ii) *If ψ has no fixed point in \mathbb{U} , then there is a point $w \in \partial\mathbb{U}$ such that $\psi_n \xrightarrow{k} w$ and the angular derivative of ψ exists at w , with $0 < \psi'(w) \leq 1$.*

We call the unique attracting point w , the Denjoy-Wolff point of ψ . By the Denjoy-Wolff Iteration Theorem, a general classification of a non-elliptic holomorphic self maps of \mathbb{U} can be given: let w be the Denjoy-Wolff point of a holomorphic self-map of \mathbb{U} . We say ψ is of dilation type if $w \in \mathbb{U}$, of hyperbolic type if $w \in \partial\mathbb{U}$ and $\psi'(w) < 1$, and of parabolic type if $w \in \partial\mathbb{U}$ and $\psi'(w) = 1$.

In the present paper we characterize the eigenfunctions of a weighted composition operators on $H(\mathbb{U})$.

2. Main Result

From now on, we assume that w is the Denjoy-Wolff point of non-elliptic holomorphic self-map ψ and φ is a holomorphic function on \mathbb{U} which is continuous at w and $\varphi(w) \neq 0$.

We characterize the eigenfunctions of $C_{\varphi,\psi}$ in $H(\mathbb{U})$. In fact, if the infinite product $\prod_{n=0}^{\infty} \frac{1}{\varphi(w)} \varphi(\psi_n(z))$ converges uniformly on compact subsets \mathbb{U} then, the function

$$g(z) = \prod_{n=0}^{\infty} \frac{1}{\varphi(w)} \varphi(\psi_n(z)) \quad (1)$$

is holomorphic on \mathbb{U} and satisfies the equation $\varphi.g \circ \psi = \varphi(w)g$ and is indeed an eigenfunction of $C_{\varphi,\psi}$. That all eigenfunctions of $C_{\varphi,\psi}$ in

$H(\mathbb{U})$, continuous at w , are obtained in this way is the content of the following theorem.

Proposition 2.1. *Let $g \in H(\mathbb{U})$ be a non-zero eigenfunction of $C_{\varphi,\psi}$ which is continuous at w . Then either $g(w) = 0$ or the infinite product $\prod_{n=0}^{\infty} \frac{1}{\varphi(w)} \varphi(\psi_n(z))$ converges uniformly on compact subsets of \mathbb{U} and*

$$g(z) = g(w) \prod_{n=0}^{\infty} \frac{1}{\varphi(w)} \varphi(\psi_n(z)). \quad (2)$$

Proof. Let $g(w) \neq 0$ and $\varphi.g \circ \psi = \lambda g$ for some non-zero scalar λ . Then $\lambda = \varphi(w)$ and for integer $n \geq 1$,

$$\left(\prod_{i=0}^{n-1} \varphi(\psi_i(z)) \right) g(\psi_n(z)) = \lambda^n g(z) = \varphi(w)^n g(z)$$

and so

$$\left(\prod_{i=0}^{n-1} \frac{1}{\varphi(w)} \varphi(\psi_i(z)) \right) g(\psi_n(z)) = g(z) \quad (z \in \mathbb{U}, n \geq 1) \quad (3)$$

where ψ_0 is the identity map on \mathbb{U} . Since $g(\psi_n(z)) \rightarrow g(w)$, the infinite product $\prod_{i=0}^{+\infty} \frac{1}{\varphi(w)} \varphi(\psi_i(z))$ converges in $H(\mathbb{U})$ to $g(w)^{-1}g(z)$ and (2) is deduced. \square

The next proposition shows the iterate sequence of holomorphic self maps can exhibit a stronger form of convergence to the Denjoy-Wolff point.

Proposition 2.2. *The series*

$$\sum_{n=1}^{+\infty} |\psi_n(z) - w|^\beta \quad (4)$$

converges uniformly on compact subsets of \mathbb{U} whenever

1. ψ is not parabolic and $\beta > 0$, or
2. ψ is parabolic automorphic and $\beta = 2$.

Proof. Suppose ψ is not of parabolic type. Then it is either of dilation or hyperbolic type. Let ψ be of dilation type and $w \in \mathbb{U}$. Then zero is the Denjoy-Wolff point of the self map $\alpha_w \circ \psi \circ \alpha_w$ where $\alpha_w(z) = \frac{z-w}{1-\bar{w}z}$. Choose $\delta > 0$ with $|\psi'(w)| < \delta < 1$. So $|\alpha_w \circ \psi \circ \alpha_w(z)| < \delta|z|$ when z is sufficiency near to zero. If K is a compact subset of \mathbb{U} , then by the Denjoy-Wolff Theorem, $\alpha_w \circ \psi_n \circ \alpha_w \rightarrow 0$ uniformly on K and $|\alpha_w \circ \psi_{n+k} \circ \alpha_w(z)| < \delta^k |\alpha_w \circ \psi_n \circ \alpha_w(z)|$ for sufficiently large n , every positive integer k , and $z \in K$. Upon replacing $\alpha_w(z)$ instead of z in the previous inequality, we get

$$\frac{|\psi_{n+k}(z) - w|}{2} \leq |\alpha_w(\psi_{n+k}(z))| \leq \delta^k |\alpha_w(\psi_n(z))| \quad (5)$$

Now suppose ψ is hyperbolic and $w \in \partial\mathbb{U}$, then $0 < \psi'(w) < 1$ and by Julia-Caratheodory Inequality ([1], Theorem 3.1) we get

$$\frac{|\psi(z) - w|^2}{1 - |\psi(z)|^2} < \psi'(w) \frac{|z - w|^2}{1 - |z|^2} \quad (z \in \mathbb{U}).$$

By substituting $\psi_n(z)$ for z , we get

$$\frac{|\psi_n(z) - w|^2}{1 - |\psi_n(z)|^2} < (\psi'(w))^n \frac{|z - w|^2}{1 - |z|^2} \quad (z \in \mathbb{U}, n \geq 0)$$

Now if K is a compact subset of \mathbb{U} , then the right hand of the above inequality is bounded on K . Hence it follows that

$$|\psi_n(z) - w| < \text{const.} (\psi'(w))^{\frac{n}{2}} \quad (z \in K) \quad (6)$$

Thus the inequality (5) and (6) imply that (4) converges uniformly on compact subsets of \mathbb{U} for $\beta > 0$. For the next part let ψ be of parabolic automorphic type. The Linear-Fractional Model Theorem [2, 8, 9] then provides a function σ holomorphic on \mathbb{U} with values in the right half-plane such that $\sigma \circ \psi = \sigma + ib$ for some real $b \neq 0$. Hence more generally $\sigma \circ \psi_n = \sigma + nib$. Let K be an arbitrary compact subset of U . For $n \geq 1$, pick $z_n \in K$ such that $|\psi_n(z_n)| \leq |\psi_n(z)|$ for all $z \in K$.

The Blaschke condition for a sequence (z_n) in \mathbb{U} is equivalent, via the map $w = \frac{1+z}{1-z}$, to the condition

$$\sum_n \frac{\text{Re } w_n}{|1 + w_n|^2} < \infty \quad (7)$$

for sequences (w_n) in the right half-plane. Since the sequence $(\sigma(z_n))$ is bounded, so (7) is to be satisfied by the sequence $w_n = \sigma(z_n) + nib$, which is therefore, the zero-sequence of a bounded holomorphic function F on the right half-plane (see [3] Theorem 11.3, page 191). The function $f = F \circ \sigma$ is then a non-constant bounded holomorphic function on \mathbb{U} , and for $n \geq 1$:

$$f(\psi_n(z_n)) = F(\sigma(\psi_n(z_n))) = F(\sigma(z_n) + nib) = 0$$

Thus some nonconstant bounded holomorphic functions on \mathbb{U} vanishes at each point of the sequence $(\psi_n(z_n))$, so that sequence satisfies the Blaschke condition. On the other hand, by the Julia-Caratheodory Inequality,

$$|\psi_n(z) - w|^2 \leq \text{const}(1 - |\psi_n(z)|^2) \leq \text{const}(1 - |\psi_n(z_n)|^2)$$

on K . Thus (4) uniformly converges on K for $\beta = 2$. \square

Recall that for any $w \in \overline{\mathbb{U}}$ and positive real number β , we denote by $Lip_\beta(w)$, the class of holomorphic functions φ satisfying

$$|\varphi(z) - \varphi(w)| = O(|z - w|^\beta) \quad (z \rightarrow w) \quad (8)$$

For example if $\varphi \in H(\mathbb{U})$ is analytic at w , then $\varphi \in Lip_\beta(w)$ for $\beta \in (0, 1]$. Moreover, if $\varphi^{(i)}(w)$ exists and equal to zero for $i = 1, \dots, n$ then $\varphi \in Lip_\beta(w)$ for $\beta \in (0, n + 1]$.

Theorem 2.3. *Let $\varphi \in Lip_\beta(w)$ and $\varphi(w) \neq 0$ then the function $g(z)$ defined by equation (1) is an eigenfunction for $C_{\varphi, \psi}$, whenever*

1. ψ is of dilation type, or
2. ψ is of hyperbolic type and $\beta > 0$, or
3. ψ is of parabolic automorphism type and $\beta = 2$.

Proof. Assume that $\varphi \in Lip_\beta(w)$ for some real number β and K is a compact subset of \mathbb{U} . Since $\psi_n \rightarrow w$ uniformly on K , by substituting $\psi_n(z)$ instead of z in (8) we get

$$|\varphi(w) - \varphi(\psi_n(z))| = O(|w - \psi_n(z)|^\beta) \quad (z \in K, n \rightarrow \infty)$$

whence

$$|1 - \frac{1}{\varphi(w)}\varphi(\psi_n(z))| = O(\frac{1}{|\varphi(w)|}|w - \psi_n(z)|^\beta) \quad (z \in K, n \rightarrow \infty).$$

Now if ψ is hyperbolic and $\beta > 0$ or ψ is parabolic automorphism and $\beta = 2$ then by pervious Proposition, $\sum_{n=0}^{\infty} |1 - \frac{1}{\varphi(w)}\varphi(\psi_n(z))|$ and consequently $g(z) = \prod_{n=0}^{\infty} \frac{1}{\varphi(w)}\varphi(\psi_n(z))$ converges uniformly on K . Thus (1) is indeed an eigenfunction for $C_{\varphi,\psi}$ and the proof is complete. \square

References

- [1] P. S. Bourdon and J. H. Shapiro, Cyclic phenomena for composition operators, *Mem. Amer. Math. Soc.*, 596 (1997).
- [2] V. Chkliar, Eigenfunctions of the hyperbolic composition operator, *Integr. Equ. Oper. Theory*, 29 (1997), 264-367.
- [3] P. L. Duren, Theory of H^p spaces, *Pure and Applied Mathematics*, Academic Press, New York, 1970; reprinted by Dover, 38 (2000).
- [4] V. Matache, The eigenfunctions of a certain composition operator, *Contemp. Math.*, 213 (1998), 121-136.
- [5] J. H. Shapiro, *Eigenfunctions for hyperbolic composition operators*, 2010.
- [6] J. H. Shapiro, *Composition operators and classical function theory*, Springer-Verlag New York, 1993.
- [7] J. H. Shapiro, *Notes on dynamics of linear operator. holomorphic self map*, 2001.

Hamid Rezaei

Department of Mathematics

Assistant Professor of Mathematics

Yasouj University

Yasouj, Iran

E-mail: hrezaei81@yahoo.com